

UNIT 17

NERVOUS COORDINATION

Major Concept

- 17.1 Nervous System of Human
- 17.2 Neurons and their types
- 17.3 Impulses
- 17.4 Synapse
- 17.5 Basic Organization of Human Nervous System
- 17.6 Effects of Drugs on Nervous System
- 17.7 Disorders of Nervous System and Diagnostic Tests

Learning Outcomes

Students will be able to:

- Recognize receptors as transducers sensitive to various stimuli.
- Trace the path of a message transmitted to the CNS for processing.
- Identify the three neurons (sensory, intermediate, motor) involved in nervous transmission.
- Identify muscles and glands as the effectors.
- Describe the detailed structure of a sensory neuron, associative and a motor neuron and relate the specialization in structures with functions.
- Differentiate between myelinated and non-myelinated neurons.
- Explain the function of the three types of neurons with the help of a reflex arc.
- Define nerve impulse.
- Describe the generation and transmission of nerve impulse.
- Name the factors responsible for the resting membrane potential of neuron.
- Evaluate from a graph the phenomena of polarization, depolarization and hyperpolarisation of membrane.
- Compare the velocities of nerve impulse in the axon membrane and in the synaptic cleft.
- Describe the role of local circuits in saltatory conduction of nerve impulse.
- Describe the structure of synapse.
- Explain synaptic transmission of nerve impulse.
- Classify neurotransmitters as inhibitory and excitatory and list some common examples.
- Identify the main components of the nervous system.
- Explain briefly the functions of major divisions of brain.
- Describe the architecture of human brain and compare its sectional view with that of the spinal cord.
- Describe cranial and spinal nerves in man.

- Explain the structure, types and functions of autonomic nervous system.
- Explain the structure and functioning of the receptors for smell, taste and touch/pain.
- Define narcotic drugs as agents that interact with the normal nervous activity.
- Compare the use and abuse of drugs with respect to heroine, *Cannabis*, nicotine, alcohol and inhalants like nail polish remover and glue.
- Define and explain the terms; drug addiction and drug tolerance with reference to caffeine and nicotine.
- Associate the effects of drug addiction and tolerance with the functioning of nervous system.
- Describe withdrawal symptoms of alcohol.
- Classify nervous disorders into vascular, infectious, structural, functional and degenerative disorders.
- Describe the causes, symptoms and treatment one type of each category of disorders outlined below (e.g. stroke as vascular, meningitis as infectious, brain tumor as structural, headache as functional and Alzheimer disease as degenerative disorder).
- Explain the principles of the important diagnostic tests for nervous disorders i.e. EEG, CT scan and MRI.

Introduction

All animals show some common characteristics, one of these is to produce response to **stimuli** (i.e. any internal and external change). The activities of different body parts in response to the stimuli must be **coordinated**. The coordination makes possible the integration of functions essential to animal behavior. It is must for animals and human to survive. In human and most animals there are two types of coordination, i.e. nervous coordination and endocrine coordination. This unit deals with only nervous coordination.

17.1 Nervous System of Man

The study of nervous system is known as **neurology**. The working of different body parts with cooperation to each other and under the control of coordinator (Brain, Spinal cord or ganglia) is called **coordination**.

The system of the body that provides coordination through electric signal among different body parts during the response to a particular stimulus is called **nervous system**. The most developed, advanced and evolved nervous system among all organisms is that of human.

17.1.1 Steps Involved in Nervous Coordination

Nervous coordination involves highly specialized cells known as **neurons**, which are either connected together or via centralized nervous system to form a network that is linked to the receptors and effectors.

Nervous system in human and higher animals consists of three basic elements.

- **Receptors** are cells, tissues or organs which receive stimuli and give information to sensory neurons.
- **The neuron** has the capacity to generate and conduct impulses to central nervous system where processing / analysis of information takes place and pass responses to the effectors.
- **Effectors** are structures such as cells, tissues, muscles and glands which carry out action or make responses.

17.1.2 Receptors or Transducers

Receptors are organs, tissues, cells or nerve endings that detect changes (stimuli) in the external or internal environment of an animal (*e.g.* Human). These stimuli then transmitted to the brain or spinal cord through sensory neurons.

Classification of Receptors on the Basis of Stimuli

Receptors are classified into following five types.

- **Chemoreceptors**, which detect the concentration of certain chemicals or ions, *e.g.* CO₂ level in the blood by **medulla of brain**, O₂ level by **carotid body**. The chemoreceptors for blood glucose, amino acids, fatty acids are located in the **hypothalamus of brain**, smell (**olfaction**) in the nasal epithelium, taste (**gustation**) found in tongue and osmoreceptor (detect osmotic pressure of blood) in hypothalamus.
- **Mechanoreceptors**, which detect stimuli of pressure, body position or acceleration, include **Meissner's corpuscles** in skin for touch, **Pacinian's corpuscles** also in skin, **baroreceptors** in blood vessel for pressure and stretch also in, **ear** for hearing and equilibrium.
- **Thermoreceptors** are mostly located in the skin to detect change in temperature (cold/warmth).
- **Nociceptors** are pain receptors widely distributed in the skin and other internal organs which detect damage to body tissues.
- **Photoreceptors or electromagnetic receptors**, detect light stimuli, such as rods and cone cells in the retina of our eyes.

17.1.3 Processing / Analysis of Information

All types of sensory inputs from various receptors are conveyed to coordinator *i.e.* brain and spinal cord by sensory neurons. The information collected by them is processed and analysed, for a suitable response by special types of neurons known as inter or associated or relay neurons.

17.1.4 Effectors

Effectors respond to stimuli by impulse coming via motor neuron, such as

muscles and glands. The glands secrete some types of chemicals while muscles respond by contracting.

Stimulus Receptor Sensory neuron Inter neuron (CNS) Motor neuron Effector Response.

17.2 Neurons

Neurons are the chief structural and functional units of nervous system. In addition to neurons, nervous system contains neuroglial cells, which nourish the neurons and also protect the neurons by myelin sheath.

17.2.1 Structure of Neuron

A typical neuron consists of three basic components, *i.e.* cell body, axon and dendrites. The **cell body** or **soma** or **cyton** or **perikaryon of neuron** contain, nucleus and various cellular organelles except centrioles. The neuron cell body contains a mass of granular cytoplasm and enclosed by cell membrane (neurilemma). The nucleus is centrally placed. The cell body contains a group of ribosomes associated with rough endoplasmic reticulum and Golgi bodies known as **Nissl's granules** (these help in protein synthesis and acetyl-choline forming enzyme) These granules are absent in axon and Dendron. (Fig.17.1)

The cell body of neuron is surrounded by cytoplasmic fibres, which are of two main types, *i.e.* axon and dendrites.

Axons are cytoplasmic process that conduct impulses away from cell body. Each axon is a usually thick long fibre (few mms to more than a meter in length) with a constant diameter. The cytoplasm of the axon is called **axoplasm** and cell membrane is called **axolemma**. Each axon terminates by branching to form small extension with knob like ending to cell presynaptic terminal. The neuroglia or glial cells that nourish, protect and support the neurons. In peripheral nervous system these cells are known as **Schwann cells**, which cover the axon by repeatedly wrapping it.

Interesting Information

Nervous system also contains about 50% of neuro-glial cells.




Fig. 17.1: A Typical Neuron

These cells are also covered by a fatty substance known as **myelin sheath**, which act as an insulator. Thus axons are called **myelinated fibres**. The non-myelinated portion of axon is called **node of Ranvier**, located between Schwann cells. The impulses jump from node of Ranvier to node of Ranvier, which are known as **saltatory impulses**.

Dendrites are also cytoplasmic extensions that carry impulses towards the cell body. If they are single, then called **Dendron**. They are thin, short and mostly branched. The branches are gradually tapered from the base to their tips. The gap between dendrites of one neuron and axon terminals of another neuron is called **synapse**.

Myelinated Fibres

It is transmit impulses much rapidly than non-myelinated neuron. It has larger and thicker axon because velocity of impulse depends upon the diameter, length and myelin sheath.

17.2.2 Types of Neurons

On the basis of functions, the neurons are of three types, *i.e.* sensory neurons, inter neurons and motor neurons.

i) Sensory Neurons conduct impulses form receptors to central nervous system. These are **unipolar** because only one fibre **originating** from cell body which immediately gives rise to two branches, one towards receptor and other towards central nervous system. Both fibres are structurally axon like except their terminal portions differ that is one is like dendrites and other is like axon. One (axon) carries impulse towards cell body while other (Dendron) away from cell body. Majority of sensory




Fig.17.2: Multipolar, Bipolar and Unipolar Neurons

neuron are unipolar but some are bipolar. The unipolar neurons are found specially in dorsal root of spinal cord. (Fig.17.2)




Fig.17.3: Three types of Neurons

ii) **Associated or Interneurons** are found only in the brain and spinal cord. These conduct impulses from sensory neurons to motor neurons. They also convey messages between various parts of the CNS. Inter neurons are mostly **multipolar** because many fibres arise from cell body. Their axon is thin and non-myelinated, while many dendrites carry impulses to its cell body.

iii) **Motor Neurons** conduct impulses from CNS to effectors. These are **multipolar** neurons. Their cell body contains many branched dendrites and a single long axon runs towards effector. (Fig.17.3)

Myelinated and Non-Myelinated Neurons

Myelinated neurons (Nerve fibres) are covered by fatty layer known as myelin sheath (axons) whereas non-myelinated neurons do not have a myelin sheath (Dendrites and Cell bodies). In myelinated neurons conduction of impulses are faster than non-myelinated neurons. Inter neurons are non-myelinated while motor and sensory neuron have myelinated portion.

Myelinated Neurons

The impulse jump from node of Ranvier to node of Ranvier. This is called saltatory impulse. Nerve impulse is 20 times faster in myelinated neuron than non-myelinated neuron.

17.2.3 Reflex action and Reflex arc

Reflex action or Reflexes are involuntary, automatic, unconscious or immediate response to external or internal environmental change or stimuli. The pathway through which a reflex travel is called **reflex arc**. The direction of reflex action is from receptor to sensory neuron to CNS to inter neuron and then through motor neuron to the effectors. Reflexes have no involvement of conscious portion of brain, therefore, the response is quick than the conscious pathway. (Fig.17.4)




Fig. 17.4: Reflex Arc

Example of Reflex arc: If we unexpectedly touch a hot object or pin prick, our hands are rapidly removed from it. The receptors of our hands are activated by heat of the object. This impulse is conveyed to sensory neuron leading to spinal cord via spinal nerve then to inter neuron which lies entirely within the spinal cord. The inter neuron then passes this impulse to motor neuron then to effectors, *i.e.* muscles which causes them to contract. (Fig.17.4)

17.3 Nerve Impulse

Nerve impulse is the information about a stimulus that is transmitted from receptors to CNS and from CNS to the effectors. It is a wave of electrochemical change which runs along the length of the neuron,

Extra Information

Refractory period lasts for about four millise-conds so as a neuron can conduct 250 impulses per second.

involving chemical reactions and movement of ions across the **neurilemma**. The measurement of the capacity to do electrical work is known as **electric potential**. It represents a type of stored energy produces during separation of charge through a barrier. The charges are positive or negative, which act as separating barrier in the plasma membrane of neuron (Neurilemma).

Oscilloscope

It is an instrument with a screen which display changes in the voltage on both side of plasma membrane of neuron with time. The electric potential which exists across a cell membrane is known as **membrane potential**. The membrane potential is either resting membrane potential (RMP) or action membrane potential (AMP).

Extra Information

In human, normally the speed of flow of impulse in neuron is about 100-120m/ second.

17.3.1 Resting Membrane Potential

A typical neuron at rest or non-conducting neuron is called resting membrane potential. In this case a typical neuron is more positive electrically outside than inside the plasma membrane while inside is more negative as compared to outside the plasma membrane. It measures about **-70 millivolts (-0.07 volt)**.




Fig. 17.5: Resting Membrane Potential (RMP)

Factors Involved in Resting Membrane Potential (RMP)

Major factors that are involved in RMP are:

Sodium and Potassium ions

There are many kinds of ions which are present inside and outside the plasma

membrane of neurons but the most important is sodium (Na^+) and potassium (K^+) ions. Sodium ions are tenfold higher in concentration outside than inside the membrane, whereas potassium ions are thirty fold higher in concentration inside than outside the membrane. The membrane of all the nerve cells possesses sodium and potassium pumps to transport these ions with the help of energy (ATP) against their concentration gradients. For every two potassium that are actively transported inside the membrane, three sodium ions are pumped out. Thus inside become more negative (70 mV) than the outside of the plasma membrane of nerve cell, which is more positive electrically than inside the membrane.

Table 17.1: Ionic Concentration Inside and Outside of Resting Neuron

Ingredients	Concentration (mmol/L)	
	Inside	Outside
Sodium ions	15	145
Potassium ions	150	5
Negative ions	156	30

Negative Organic Ions

Many types of large negative organic compounds are present in the both sides of plasma membrane of neurons. These organic ions include some proteins, amino acids, RNA, SO_4 , PO_4 , etc. These negative ions are much more inside the membrane than outside (outside Cl ions are present) where they are only in negligible concentration. Thus inside is more negative than outside. (Fig. 17.6)




Fig. 17.6: Depolarized Region and Saltatory Impulse

Channel Proteins (Gates) in Plasma Membrane of Neurons

The plasma membrane is virtually impermeable to all ions except potassium ions.

It is slightly permeable to potassium, so it leaks out of the neuron by diffusion. That is why inside becomes more negative than outside of the plasma membrane of neuron. Thus non-conducting neuron is in **polarized state** (i.e. RMP – 0.07 volts or 70 mV). The resting membrane potential (RMP) will be maintained in undisturbed membrane. If it is disturbed or stimulated by a sufficient stimulus known as **threshold**, then action potential will occur.

17.3.2 Action Membrane Potential (AMP) (Depolarized State)

In action/active membrane potential, inside of neuron become more positive and outside become more negative. It is called depolarized state, which happen when appropriate stimulus receives, the positive charge sodium ion tends to move inside of the neuron. The electrochemical change is so brief (about one millisecond) that only a portion of the neuron (i.e. one node of Ranvier to another node of Ranvier) is in the action membrane potential state followed by the recovery of polarized state, thus impulses flow from one node of Ranvier another to node of Ranvier.

The major factors involved in changing the resting membrane potential to action membrane potential are:

1. Threshold Stimulus

It is capability of a stimulus to bring electrochemical charge on neuron or to excite a given tissue. It is also known as **adequate stimulus** (about – 50 to –55 mV electric membrane potential). If stimulus is not capable to excite or not appropriate, then it is called sub **threshold** or **inadequate stimulus**.

Influx of Na^+ ions

When threshold is reached, the membrane become more permeable to Na^+ than to the K^+ due to the opening of Na^+ gates. Thus influx of Na^+ ions by diffusion occurs, and electric potential of the membrane changes from –70 mV towards zero and then reach to about 50mV. This reversal of polarity




Fig. 17.7: Flow of Impulse

across two sides of membrane is called **depolarization**. This state of electropositive inside and electronegative outside lasts for about one millisecond till the Na^+ gates are not closed.

Repolarized state

After the peak of action potential, called the **spike potential**, the permeability of Na^+ decreases, and now become more permeable for K^+ by opening of K^+ gates thus K^+ rapidly diffuses out from cytoplasm to extracellular fluid. The sodium gate closes and the neuron get its original polarity *i.e.* **repolarized** (inside more electro negative and outside more electro positive). Infact, there is a slight overshoot into a more negative potential than original resting potential. This is known as **hyperpolarized** state. It is due to the slight delay in closing of all K^+ gates compared with Na^+ gates. (Fig.17.7)

Interesting Information

The continuous impulse occurs in non-myelinated neuron fibres in which K^+ and Na^+ ions can move across the membrane along the length of neuron so action potential flow as a wave.

Refractory Period (Resting State)

It is a period when after an action potential, nerve fibre undergoes a period of recovery, in which it regain the original ionic distribution and polarity, thus prepare itself for the next stimulation. The original ionic distribution is restored by a sodium – potassium exchange pump which actively transport Na^+ ions out and K^+ ions in the neuron. This return the membrane to its resting potential *i.e.* from +50 mV to –70 mV.




Fig. 17.8: Refractory Period of Neuron

The resting membrane potential (RMP) will be maintained in undisturbed membrane. If it is disturbed or stimulated by a sufficient stimulus known as threshold, then action potential will occur. (Fig.17.8)

17.3.3 The role of Local Circuits in Saltatory Conduction of Nerve Impulse

Saltatory conduction is the propagation of action potential along myelinated axon from node to node. The “local circuits” explain how the action potential (AP) is transmitted along the neuron. Basically an action potential at a point in the axon, develops

a local circuit because, the influx of sodium ions at that point makes that particular point positively charged. However, regions around that point are still negatively charged (because they are still in the “resting potential formation”). The sodium ions at the point of AP are then attracted to these negatively charged regions, hence setting up a “local circuit” at those regions. This circuit then opens the sodium channels at these points, sodium ions flow in and the whole AP circle continues, hence the AP moves along the axon.

17.4 Synapse

There is no cytoplasmic connection between successive neurons, however, impulses are transmitted and this transmission occurs at synapse. **Synapses** are microscopic gaps between successive neurons. These gaps are between axon terminal of one neuron and dendrites of another neuron. (Fig.17.9)




Fig. 17.9: Synapse

A single neuron may form synapses with many incoming fibres of different neurons. The gaps between successive neurons are cleave-shaped thus called **synaptic cleft**. A neuron carries information (impulse towards a synaptic cleft known as **transmitting neuron** or **presynaptic neuron** while a neuron which gets the impulse from synaptic cleft, is called **post synaptic neuron** or **receiving neuron**. The axon

terminals of each presynaptic neuron have expended endings called **knobs** which contain many spherical sacs, known as **synaptic vesicles**. Each vesicle contain as many as 10,000 molecules of chemical messengers, the **neurotransmitter substance**. The dendrite of post synaptic neuron lacks these vesicles. The transfer of impulse across the synapse is called **synaptic transmission**.

Steps of Mechanism of Synaptic Transmission

1. When action potential reaches the axon terminal, it is received by synaptic knob. The calcium channels, present in the presynaptic membrane open and calcium from synaptic cleft transfer into the knob. Thus Ca^{++} concentration increases, the synaptic vesicles move towards the pre-synaptic membrane.
2. The vesicles containing neurotransmitter fuse with presynaptic membrane and release their neurotransmitters molecules into the synaptic cleft.
3. The neurotransmitter diffuses across the synaptic cleft and bind to receptor molecules on postsynaptic membrane.
4. The receptor of post synaptic membrane, opens some channels and allow sodium ions to diffuse across the

Extra Information

There are 100 known types of neurotransmitters.




Fig. 17.10: Steps of Synaptic Transmission

postsynaptic membrane. As a result post synaptic membrane depolarizes *i.e.* (inside + outside -) and action potential is generated. Since this depolarization brings the membrane potential towards threshold level, it is called **Excitatory postsynaptic potential (EPSP)**.

5. After the activation of post synaptic membrane, the neurotransmitters are immediately broken down by enzymes (like acetylcholinesterase for acetylcholine and monoamine oxidases for adrenaline). (Fig.17.10)

17.4.1 Classification of Neurotransmitters

There are two major classes of neurotransmitters.

1. Excitatory Neurotransmitters

These initiate nerve impulses, cause increased membrane permeability to sodium ions. They may be **acetylcholine** for peripheral nervous system, **biogenic amine** (derived from amino acids) for central nervous system. Their types are **epinephrine** and **nor epinephrine** for heart beat rate during stress; **serotonin** and **dopamine** affect mood, sleep, attention and learning. All of these function like hormones.

2. Inhibitory Neurotransmitter

This decrease membrane permeability to Na^+ ions, which result to raise threshold of stimulus. Thus the nerve impulse does not trigger or lessens, to adjoining neuron. Examples are glycine amino acid, gamma-amino-butyric acid (GABA), while the endorphin are peptides, decreases pain reception and act both as neurotransmitters and hormones.

17.5 Human Nervous System

Human and most animals (except coelenterates and Echinoderms) have centralized nervous system. Human nervous system is most advanced and also possess some unique features, *i.e.* enable us to convey our complex ideas, information and messages in the form of language, make and use various tools, preserve information (in written and video), great learning, memory storage capacities, *etc.*

Human nervous systems consist of two primary divisions, the central nervous system (CNS) and peripheral nervous system (PNS). (Flow chart 1)

Interesting Information


There are two types of synapses.

1. **Electrical synapses**, which are rapid and synaptic cleft is only 0.2 nm thick.

2. **Chemical synapses**, in which gap is 20 nm. Thus transfer of impulses occurs by neurotransmitters (Chemicals).

Extra Information

Coelenterates and echinoderms possess diffused nervous system.



Flow Chart.1: Classification of Human Nervous System

17.5.1 Architecture of Brain and Spinal Cord and their functions

Central Nervous System of Human (CNS)

It consists of brain and spinal cord, act as coordinating centre, these lies in the skull (Brain) and above the vertebral column (spinal cord) *i.e.* in midline of the body.

CNS is hollow and fluid filled. The brain has four cavities called **ventricles** while the cavity of spinal cord is known as **central canal**. The fluid in these cavities named as **cerebrospinal fluid (CSF)** (similar to blood plasma) act as cushion and bathes the neurons of CNS.

Protection of Central Nervous system

The brain is protected by bony cranium of skull while spinal cord is protected by vertebrae of vertebral column. Both brain and spinal cord are also covered by three membranes, collectively known as **meninges** (**singular; meninx**), the outer hard **Dura matter** (next to cranium), inner **pia matter**, next to brain and spinal cord and middle **arachnoid matter**. The cerebrospinal fluid is present between the piamatter and arachnoid matter.

Interesting Information

Cerebrum is the largest part of brain and contains highest number of neurons than any other part of brain.

According to "Roger Spray"

Both cerebral hemisphere of cerebrum superficially same but right and left portion function so differently that we could almost say we have two brain in one. Left cerebrum house our language centre, logic mathematical abilities while the right hemisphere imagination, spatial perception, artistic and emotional abilities.

Brain

The brain of human can be divided into forebrain, mid brain and hind brain. (Fig.17.11)

Fore Brain

Forebrain has two subdivisions; **telencephalon** (cerebrum) and **diencephalon** (thalamus and limbic system). The **cerebrum** of human is largest among all other animals (more than half of the brain). It is divided into two cerebral hemispheres, which are connected together by a band of axons known as **corpus-callosum**. It carries memory available on one side of the brain to the other side. Cerebrum contains four lobes, frontal, parietal, temporal and occipital lobe. The left cerebral hemisphere controls the right side of the body while right cerebral hemisphere controls left side of the body. Functionally, cerebrum consist of **sensory area**, (receive impulses from receptors), **motor area** (give responses) and **associated area** (interprets or analyses the incoming information) cerebrum act as control centre for sight, smell, taste, speech and hearing. It also controls voluntary movements, thinking, learning, conscious sensations, judgment, reasoning, decision-making, dreams, emotional feeling, intelligence, analysis and interpretation of memory.

Although the motor sensory and associated areas are located in all parts of cerebrum, however motor areas are more abundant in frontal lobe. The associated areas are most occupy the anterior of frontal lobe and wide spread in lateral portion of parietal, temporal and occipital lobes.

Diencephalon (Thalamus and limbic system): **Thalamus** lies below the cerebrum, act as relay centre and carries sensory impulse to the cerebrum and limbic system. Thalamus receives all sensory impulses (except sense of smell).

Limbic System

This system is located in an arc just under cerebrum. The limbic system consists of hypothalamus, amygdala, hippocampus and some part of cerebrum.




Fig. 17.11: C.S. of Brain of Man

Extra Information

There are convolutions (groove) called gyri which are seen on outer part of cerebrum (cerebral cortex). The deep grooves are called fissure while shallow grooves are called sulcus. These grooves greatly increase the surface area of cerebrum. A longitudinal fissure is present between two parts of cerebrum.

Hypothalamus located on the ventral side of thalamus. It is both nervous and endocrine centre *i.e.* link between nervous and endocrine centre.

The function of hypothalamus is to maintain homeostasis and contains centres for regulating body temperature, water balance, menstrual cycle, blood pressure, sleep wake cycle, hunger, sexual response and fight or flight.

Amygdalae are two almond shape masses of neurons on either side of the thalamus. It produces sensation of pleasure, sexual arousal when stimulated, punishment, love, hate, fear and rage.

Hippocampus consists of two horns that curves back from the amygdala and play important role in the formation of long term and short term memory, thus required for learning.

Mid Brain

In human this portion of brain is reduced. It contains auditory relay centre and centre that control reflex movement of eyes also contains auditory reflex centre. Mid brain contains reticular formation, which is a relay centre connecting fore brain with hind brain and is very important in screening the input information before they reach higher brain centres. (Fig.17.12)




Fig.17.12: Brain Stem

Hind Brain

Hind brain includes the **pons**, **cerebellum** and **medulla**.

Pons is a group of neuron, located above the medulla; act as a bridge between cerebellum, medulla and cerebrum.

It is involved in rate and pattern of breathing, sleep and wakefulness.

Cerebellum is second largest part of the brain, bulb or leaf like in shape. It coordinates body movements and maintains body position *i.e.* equilibrium. The cerebellum guides, smooth and accurate motions and maintains body position. It also involves in the learning and memory storage for behavior. It is best developed in bird, which engages them in complex activity of flight.

Medulla is last part of brain but in evolutionary point of view, it developed first. It controls several automatic functions, such as heart beat rate, blood pressure, breathing and swallowing.

The mid brain, together with pons and medulla know as **brain stem**, which support the life.

Ventricles of Brain

Human brain possesses four ventricles or cavities, which are filled with cerebrospinal fluid. The first and second ventricles are present between limbic system and cerebrum known as **lateral ventricles**. Another ventricle is present between limbic system and thalamus called **third ventricle** while **fourth ventricle** is present in medulla. There is a tube between third and fourth ventricle known as **iter or cerebral aqueduct**, while an opening between lateral ventricles and third ventricle is called **intra ventricular foramen**. (Fig.17.13)




Fig. 17.13: Ventricles of the Brain

17.5.2 Spinal Cord

Medulla narrows down into the spinal cord. It is an elongated, hollow fluid filled and cylindrical structure, extends from the **foramen magnum** (a hole in the bottom of skull) lying in the neural canal of vertebral column (up to 3rd lumbar vertebrae). In a cross section of spinal cord exhibits inner butterfly shaped **grey matter** while peripheral **white matter**. There is a tiny central canal in the centre of grey matter, filled with CSF around the **central canal** is a single layer of cells called **ependymal layer**.

Interesting Information

Local anaesthesia is given at fourth lumbar, so as to protect the spinal cord.

The grey matter consists of non-myelinated portion *i.e.* mostly cell bodies of

neuron while white matter is composed of myelinated nerve fibres *i.e.* mostly axons. Spinal cord controls reflexes below the neck region. It also conducts impulses to and from the different part of the bodies and brain. Thus helps in better function of brain. (Fig.17.14)




Fig. 17.14: C.S. of Spinal Cord

17.5.3 Architecture of Human Brain and compare its sectional view with that of spinal cord

Both brain and spinal cord are made of white and grey matter. The difference is in brain grey matter mostly lies outside (Cerebral cortex) and white matter lies inside (Cerebral medulla) while spinal cord grey matter lies inside look like butterfly shape and white matter lies outside. (Fig.17.14)




Fig. 17.15: Cross Section view of Brain and Spinal Cord

17.5.4 Peripheral Nervous System

It is the system of nerves and ganglia. The **nerves** are cables of bundles of nerve fibres (axon and dendron fibres) while ganglion is the concentration of cell bodies of neurons with in peripheral nervous system. Nerves may be **sensory nerves** (contains sensory neurons), **Motor nerves** (contain motor neurons) and **mixed nerves** (contains both sensory and motor nerves). There are two types of nerves on the basis of their origin *i.e.* spinal nerves and cranial nerves.

Spinal Nerves

In human thirty-one pairs spinal nerves, all are mixed. Each serves those regions of the body where it is located, Cervical, 8; Thoracic, 12; Lumbar, 05; sacral, 05; coccygeal, 01. Each nerve has a dorsal root, contains sensory fibres and ventral root contains motor fibres. Both of these roots join just before a spinal nerve leave the vertebral column.

Cranial Nerves

These nerves are also called **cerebral nerves**. There are 12 pairs of cranial nerves arising from or lead to the brain. The cerebral nerves functionally may be sensory (three pair I, II VIII), motor (five pairs, III, IV, VI, XI, XII) and mixed in nature (four pair V, VII, IX and X). These are mostly concerned with head, neck and facial regions of the body, only 10th cranial nerves, named **vagus** have branches to the pharynx, larynx and most internal organs.

The peripheral nervous system functionally subdivided into the somatic and autonomic nervous system.

Somatic Nervous System (SNS)

This subdivision controls voluntary movements which are consciously controlled, involving skeletal muscles except reflex action of skeletal muscles.

Autonomic Nervous System (ANS)

This subdivision controls involuntary responses (automatic and subconscious), consists of both sensory neurons and motor neurons that runs between central nervous system and many viscera like lungs, heart and glands. Autonomic nervous system also controls the contraction of cardiac and smooth muscles.

The motor neurons of autonomic nervous system are divided into sympathetic nervous system and parasympathetic nervous. The differences between these two system are explained in the Table 17.1. (Fig.17.16)

Table 17.1: Differences between Sympathetic and Parasympathetic Nervous System

Sympathetic Nervous System	Parasympathetic Nervous System
- It prepares the body for emergency situations and associated with "flight and flight". Increase metabolism to avoid danger. - Accelerates the heart beat from set point, rise BP. - Dilates Pupils - Inhibits digestion of food - Most preganglionic fibre (nerves) arise from the middle portion of the spinal cord and almost immediately terminate in ganglia that lie near the spinal cord. - Preganglionic fibres are short and postganglionic long.	- It promotes all internal responses which are concerned with the rest situation, and maintain body homeostasis, <i>i.e.</i> returns body functions to normal position. - Retards heart beat that is maintain at set point and lowering of BP. - Contracts Pupils. - Increases digestion of food. - A few cranial nerves, including the vagus nerve, together with fibres that arise from the sacral (bottom) portion of the spinal cord. - Preganglionic fibres are long and postganglionic short.




Fig. 17.16: Parasympathetic and Sympathetic Nervous System

17.5.5 Structure and Functioning of Receptors of Smell, Taste, Touch and Pain

1) Smell or Olfactory Receptors

The receptors, which are stimulated by chemicals are called smell or olfactory receptors. In human it is not as much developed or important as vision and hearing. Although in most predators sense of smell is highly developed and important to detect preys. These receptors are located in the upper part of the nasal cavity, which are neurons. The neurons are surrounded by ciliated columnar epithelial cells. Chemicals that stimulate the olfactory receptors enter the nasal cavity as gases, which dissolve in watery fluids that surround the cilia before they can be detected. The axons of neurons carry the smell impulses to the olfactory bulb of fore brain for appropriate responses. (Fig.17.17)

Interesting Information

There are about 1,000 different types of receptor protein on receptor neurons, each is sensitive to different odors.




Fig. 17.17: Olfactory and Gustatory Receptors

2) Taste Receptors (Gustation)

These receptors are located on the throat and mouth especially in the upper surface

of tongue as many raised structure called **papillae or taste buds**. Each taste bud has a pore through which fluid in the mouth contact the surface of receptor cells.

There are thousands of taste buds which perceived tastes, all are combination of four primary sensations; **sweet** (elicited by sucrose, glucose and other simple sugars), **sour** (acids), **salty** (NaCl and other salts) and **bitter** (alkaloids and other potentially toxic plant substances). (Fig.17.18)

3) Touch (Tactile Receptor)

There are two touch receptors (also called **mechanoreceptors**) disc-shaped dendrite endings called **Merkel's disc** and egg shaped receptors called **Meissner's corpuscles**. (the word corpuscle mean "Little body"). (Fig.17.19)




Fig. 17.18: Four Receptor on the Tongue




Fig. 17.19: T.S Skin

Both these receptors are widely distributed in the skin but are most numerous in the hands (finger tips), feet, and eyelids, tip of the tongue, lips, nipples, clitoris and tip of the penis. In addition, free nerve endings wrap around the roots of hairs, detect stimulus like wind or an insects that moves body hair.

Pressure receptors called **pacinian corpuscles** are placed deeper with in skin.

1) **Pain Receptors (Nociceptors)**

There are three types of pain receptors, *i.e.* **cutaneous** (skin), **somatic** (Joints and bones), and **visceral** (all body organs except brain). These receptors are naked dendrites that respond to chemicals released from injured tissues or excess stimuli of pressure and heat.

2) **Thermoreceptors**

Which are encapsulated nerve endings located in the dermis of skin (for hot **Ruffini ending** and for cold **Krause end bulbs**).

17.6 Effects of Drugs on Nervous Coordination

Broadly speaking, a **drug** is a substance introduced into the body to provoke a specific physiological response. Some drugs help a person cope with illness or emotional stress. Other act on the brain, artificially fanning pleasure associated with sex and other self-grafting behaviors. Many drugs are habit-forming. **Habituation** and tolerance are sign of **drug addiction**.

A **narcotic** is a group of substances, bind to certain pain killing sites in the brain thus stop the perception of brain. Their constant use blocks the production of endorphin (natural pain killer hormone secreted from anterior pituitary gland). Thus narcotic acts as an agent which interacts with the normal nervous activity.

The **side effect of narcotics** is change in mood, drowsiness, nausea, vomiting, **euphoria** (an exaggerated feeling of wellbeing), *etc.* Some common narcotic drugs are discussed here.

17.6.1 Heroin

It is most commonly use addictive illegal drug in our country, processed from morphine (extracted from the seed pod of certain varieties of **poppy plant**). Heroin gives a feeling of euphoria and acts as pain killer. It is usually injected via intravenous or intramuscular, sniffed/snorted or smoked.

Their side effects are nausea, vomiting, **dysphoria** (abnormal depression and discontent), respiratory depression, severe itching, dry mouth and **spontaneous abortion** *etc.*

17.6.2 Marijuana, Hashish (*Cannabis sativa*)

Marijuana and hashish is made from crushed leaves, flowers and stems of the plant *cannabis sativa*, in low dose it is like a **depressant**. It slows down but does not impair motor activities. It relaxes the body and elicits mild euphoria. It also causes disorientation, increased anxiety bordering to panic, **delusion** (including paranoia) and hallucinations. *Cannabis* has very beneficial effects as medicine, (stimulate hunger in Acquired Immunodeficiency Syndrome (AIDS) patients, chemotherapy). It is also effective for treating **glaucoma** (eye disease) by reducing intraocular eye pressure.

17.6.3 Nicotine

It is an alkaloid found in tobacco, has powerful effects on nervous system (both CNS and PNS). It mimics acetylcholine and directly stimulates a variety of sensory receptors. In short term effects include water retention, irritability, and increased heart beat rate and high blood pressure. It also causes gastric upsets.

17.6.4 Alcohol

Alcohol directly acts on the plasma membrane to alter cell function. It is lipid soluble thus easily crosses the blood brain barrier to exert rapid effects. Alcohol is one of the most powerful psychoactive drugs and a major factor in many deaths. In short term, a small amount causes disorientation and uncoordinated motor functions, diminish judgment. Long term addiction can lead to liver **cirrhosis**.

Withdrawal symptoms of Alcohol

The symptoms are feeling of nervousness, anxiety, irritability, shakiness, depression, headache, fatigue, rapid emotional changes and hallucination, *etc.*

17.6.5 Inhalants

Inhalants are **volatile** (vapours / gases) organic chemicals, which enhance psychoactive effects (Mind alerting) when abused or misused by concentrating and intentionally inhaling these fumes. Examples are fuel gases nitrates and anaesthetic gases. These are found in **glue, thinner, paint, gasoline and nail polish remover**.

Inhalant provide and instant rush like alcohol, causes euphoria followed by central nervous system depression.

Deep breathing of inhalant may result in loss of self-control, violent behavior, nausea, vomiting, unconsciousness, giddiness, loss of appetite. Their high dose cause hallucination, slurred speech, loss of motor skills and heart palpitation.

17.6.6 Drugs Addiction

It is a state when a person highly dependence on illegal drugs or medication. Many people want to quit these drugs but they cannot do it on their own.

17.6.7 Drug Tolerance

If a person's diminished response to a drug, which occurs when drug is used repeatedly and the body adapts to the continued presence of the drug. Examples are daily users of tea, coffee and cigarette smokers.

17.6.8 Effects of Drug Addiction and Drugs Tolerance on CNS

There are three main types of drugs affecting the CNS.

i) Depressants

Those drugs that slow down the functions of the CNS. These drugs (like alcohol, heroin) don't necessary make a person feel depressed, instead feel more relaxed if use in small quantities. Although if use in large quantities then cause vomiting, unconsciousness and even may fatal. It can disturb our coordination and concentration.

ii) Stimulants

Stimulants like caffeine, nicotine, cocaine, amphetamine and many other stimulants increase alertness and body activity, then cause depression. Caffeine, a truly common stimulant, which is present in coffee, tea, chocolate and many soft drinks. Large quantities of stimulants can "over stimulate" the user, causing anxiety, panic, headache stomach cramps, aggression and paranoia.

iii) Hallucinogens

These affect perception, the user may believe they see or hear things that are not really there, or what they see may be distorted in some way. These include, *Cannabis*, *Datura* which interfering with the activities of acetylcholine, norepinephrine or serotonin.

17.7 Disorders of Nervous System and Diagnostic Tests

Diseases of nervous system are also called neurological disorders. These disorders can be classified into vascular, infectious, structural, functional, and degenerative.

17.7.1 Vascular Disorder of Central Nervous system

This disorder occurs due to abnormality in blood circulation to CNS e.g. stroke and brain haemorrhage.

Strokes

It is also called **cerebrovascular accident (CVA)** and occurs due to rupture of small perforating cerebral arteries.

Interesting Information

Amphetamine is a commonly used street drug. These include methamphetamine and phentermine. Cocaine (coke), is a strong stimulant, used as a recreational drug.

Causes: Hypertension, diabetes mellitus, thrombosis, embolism, cigarette smoking, drugs abuse, etc.

Symptoms: Sudden loss of function in one region of brain, which causes numbness of the face, arm or leg of one side of the body (paralysis occur).

- **Aphasia** i.e. inability to express through in word.
- Trouble seeing in one or both eyes.

Treatment: Mostly anticoagulants (such as Heparin) and platelet aggregation inhibitor (such as aspirin) are used.

- Blood pressure management and nursing is essential.

17.7.2 Infectious Disorders of Central Nervous System

This can be caused by almost any infectious agent, including viruses, bacteria, fungi, protozoa and platyhelminthes. Meningitis is a common example.

Meningitis

It is an inflammation of the **meninges**. It may be virally induced but more often bacterial, sometime other microorganisms.

Symptoms: The common symptoms are stiffness in the neck, photophobia, headache, nausea, vomiting and fever. In **severe cases meningitis** may also cause paralysis, coma and even death.

Treatment:

- **For viral meningitis** there is no specific treatment and the condition is usually benign and self-limiting.
- **Recovery** usually occurs within days.
- **Antibiotics** and corticosteroid are also used as general treatment.

17.7.3 Structural Disorders of CNS

Several disorders disturb the structure of brain are referred as structural disorders, such as tumors.

Tumor

The tumor is an abnormal **mass of neuroglial cells** produced as a result of **uncontrolled cell division**. It may be **benign or malignant tumors**.

Causes: It is caused by mutation and may occur at any age in brain and spinal cord.

Symptoms: The sign and symptoms vary widely, depending on the location of the tumor but may include, headache, vision problem, **neuralgia** (Pain along the distribution of a peripheral nerve). Some of the symptoms are paralysis, hearing problems, seizures, behavior change, coma and death.

Treatment: Tumor is removed Surgically.

Functional Disorders of CNS

Headache: Headache or **cephalalgia** is pain anywhere in the region of the head or neck.

Symptoms: The brain tissue itself is not sensitive to pain because it lacks pain receptors. Rather, the pain is caused by **disturbance of the pain-sensitive structures around the brain.**

There are two major categories of headaches.

1. Primary Headaches

It is due to the **headache condition itself** and not due to another cause. Primary headaches include **migraine**, tension and cluster headaches.

2. Secondary headaches

It is due to an underlying structural problem in the head or neck such as bleeding in the brain, tumor, meningitis, encephalitis, sinusitis and certain drugs.

Treatment: Headache is treated by several analgesic drugs.

17.7.4 Degenerative Disorders of CNS

Many diseases cause degeneration in different part of the nervous system without an identifiable external cause or it may be genetic, example is Alzheimer's disease.

Alzheimer's disease (AD)

It was first described by Alois Alzheimer in 1907.

Alzheimer's disease is a slowly progressive disease of the brain that is characterized by impairment of memory and eventually by disturbances in reasoning, planning, language, and perception.

Symptoms: There is decline in the brain function in aged peoples. Its symptoms are similar to those diseases that cause **dementia** (memory loss/ impairment of memory/ particularly for recent events *i.e.* short-term memory).

Causes: There is a genetic pre-disposition to the disease in some people, therefore runs in families. High levels of aluminium may contribute to the onset of this disease.

Treatment: So far no effective treatment is known for this disease.

17.7.5 Diagnostic Tests for Nervous Disorders

Some common diagnostic tests for nervous disorders.

Electroencephalogram (EEG)

The neurons of the cerebral cortex continuously generate electric activity. This activity can be recorded by electrodes attached to precise location on the scalp, producing **electroencephalogram (EEG)**. An EEG pattern is commonly called **brain waves**. (Fig.17.20)




Fig. 17.20: Electroencephalogram (EEG)

Classification of Brain Waves (EEG of brain in Sleep and fully awake person)

The EEG regular patterns are classified as alpha, beta, theta and delta waves.

During wakefulness alpha and beta activities are experienced in human brain.

Alpha wave: These waves consists of medium frequency waves.

Beta wave: These waves consists of irregular low amplitude waves found during highly alert and attentive state.

Theta waves: These waves can occur in adults who have drowsy brain. This is a transition stage between sleep and wakefulness.

Delta waves: These waves are of high altitude occurring in deep sleep, in infancy and in patient with severe brain disorders. Distinct types of EEG patterns can be detected in patients with specific brain disorders such as epilepsy.

Computer Tomography (CT) Scan

1. It is an **X-Ray procedure** that combines many X-ray images with the aid of computer to generate cross-section views and if needed **3d images** of internal organs and structures of the body.
2. Computerized tomography is more commonly known by its abbreviated names, **CT scan** or **CAT scan**. A CT scan is used to




Fig. 17.21: C.T Scan

define normal and abnormal structures in the body and / or assist in procedures by helping to accurately guide the placement of instruments or treatments. (Fig.17.21)

Tomograph and Tomogram

1. A large donut- shaped X-ray machine or scanner called tomography takes X-ray images at many different angles around the body.
2. These images are processed by a computer to produce cross-sectional pictures of body. In each of these pictures body is seen as an X-ray "slice" of body, which is recorded on a film. This recorded image is called a tomogram.

Magnetic Resonance Imaging (MRI):

An MRI (magnetic resonance imaging) scan is a radiology technique that uses magnetism, radio waves and a computer to produce images of body structures.

MRI Scanner is a tube surrounded by a giant circular magnet.

Mechanism

- Patient is placed on a moveable bed that is inserted into the magnet.
- The patient is exposed to strong magnetic field and beam of radio-waves.
- The receiver information is processed by a computer and an image is produced.
- Image and resolution produced by MRI is quite detailed and can detect tiny changes of structures within the body.




Fig. 17.22: A MRI Scanner

Science, Technology and Society (STS)

1. **Justify the way nervous system helps to coordinate complex and intricate movement of hand to play a piano or write alphabets.**

The information gathered by the sensory receptors in the form of electrical impulses is a process called transduction. The nerve impulses are then transmitted over peripheral nervous system to the central nervous system. There the signals are integrated; *i.e.* they are brought together, creating a sensation (preceptions), adding to the memory or helping to produce thoughts. As a result of this integrative function, conscious or subconscious decisions are made and then acted upon by motor functions.

The motor functions of the nervous system employ peripheral nerves. They carry impulses from the central nervous system to the effector *i.e.* muscles that contract when stimulated by impulses. Thus hand and fingers move, so that we may play a piano or write alphabets.

2. Ascertain the effect of nerve gas as an inhibitor or acetylcholinesterase.

The nerve gases are a class of organic chemicals that disrupt the mechanism by which nerve transfer messages to organs. The principle upon which nerve gases work is to inhibit a particular enzyme in the body, namely acetylcholinesterase which rapidly break down the neurotransmitter, acetylcholine. The reason is that the organophosphorous compound transfers a phosphate group to the enzyme, making it inactive. As a result acetylcholine build up causing a jam in the nervous system and muscle contraction continue uncontrollably. The person will die.

SUMMARY

- Nervous system detects, interprets, and issue calls for response to stimuli in the animal's (*e.g.* human) external and internal environment.
- Receptors are the parts of the body that receive stimuli from internal or external environment.
- Neurons are the structural and functional unit of nervous system.
- There are three basic components of each neuron, *i.e.* axon, cell body and dendrites.
- Based on the function, there are three types of neuron, sensory, associated and motor neurons.
- Reflex action is immediate, involuntary and automatic response to external and internal stimuli.
- Nerve impulse is a wave of electrochemical changes that travels along the length of neuron.
- Synapse is a microscopic junction between two successive neurons, which transmit impulse from one neuron to another neuron.
- Central nervous system consists of brain and spinal cord
- The brain is divided into three parts, fore brain, mid brain and hind brain.
- The peripheral nervous system includes cranial nerves and spinal nerves.
- Olfactory receptors are chemoreceptors, stimulated by chemicals dissolved in liquid.
- The receptors for touch and pressure are also called mechanoreceptors.

- Endorphins are the natural pain killer chemicals in the brain.
- Nicotine is an alkaloid derived from tobacco.
- Meningitis is the inflammation of the meninges.
- Tumor is an abnormal mass cells which produced as a result of uncontrolled cell division.
- A headache is also called cephalalgia.
- EEG, CT scan and MRI scan are commonly used diagnostic test for nervous disorders.
- Inhalants are volatile organic chemicals, commonly called "glue sniffing".

EXERCISE

SECTION-I: OBJECTIVE QUESTIONS

Multiple Choice Questions (MCQs)

A. Choose the correct answers.

1. Nissel's granules is the group of:

(a) Mesosomes	(b) Lysosomes
(c) Ribosomes	(d) Chromosomes
2. Main transmitter for synapses that lie outside CNS is:

(a) Serotonin	(b) Acetylcholine
(c) Dopamine	(d) Adrenalin
3. In human rely center is located in:

(a) Fore brain	(b) Mid brain
(c) Hid brain	(d) Spinal cord
4. The part of brain which controls breathing, heart rate and swallowing is:

(a) Cerebellum	(b) Medulla
(c) Thalamus	(d) Cerebrum
5. Which of the following receptors produces sensation of touch and pressure?

(a) Chemoreceptor	(b) Mechanoreceptors
(c) Nociceptors	(d) Photoreceptors
6. Chemoreceptors detect the stimulus of:

(a) Pressure	(b) Temperature
(c) Light	(d) Smell
7. The portion of neuron that conducts nerve impulse towards cyton (cell body) is:

(a) Axon	(b) Dendrites
(c) Myelin	(d) Soma

8. Neurilemma is the membrane of:
 - (a) Muscle cell
 - (b) cyton of neuron
 - (c) Axon
 - (d) Nerve fibre
9. The fibres at the end of axon are called
 - (a) Dendrites
 - (b) Dendron
 - (c) Axon fibres
 - (d) Cyton
10. Synapse is microscopic gap between two successive:
 - (a) Neuroglia
 - (b) Neuron
 - (c) Axons
 - (d) Soma
11. Brain and spinal cord together known as:
 - (a) PNS
 - (b) ANS
 - (c) SNS
 - (d) CNS
12. Second largest part of brain is:
 - (a) Fore brain
 - (b) Cerebrum
 - (c) Cerebellum
 - (d) Medulla
13. The number of cranial and spinal nerves in human are
 - (a) 24 and 62
 - (b) 12 + 31
 - (c) 20 + 68
 - (d) 20 + 62
14. The volatile organic chemicals, once commonly referred to as "glue sniffing" are
 - (a) Cannabis
 - (b) Inhalants
 - (c) Narcotics
 - (d) Caffeine

B. Fill in the blanks.

1. Neurotransmitter molecules bind to the receptors on the membrane at synapse.
2. A microscopic gap between two consecutive neurons is called
3. The collection of axons and dendrites are called
4. Muscles and glands are principal
5. An infection of the central nervous system is called
6. The structures which respond by impulse coming from motor neurons are
7. Nociceptors detect stimulus of
8. Neurons are structural and functional unit of
9. Nissl's granules are the group of in the neuron cell body.
10. Serotonin and acetylcholine are

SECTION-II: SHORT QUESTIONS

C. Give the short answers of the following questions.

1. Define nervous coordination, receptors and effectors.
2. Describe briefly about axon.
3. Differences between sensory neuron and motor neuron.
4. Difference between reflex action and reflex arc.
5. Define synapse, presynaptic neuron and post synaptic neuron.
6. Define neurotransmitter, write name of any four neurotransmitters.
7. Write note on mid brain.
8. Describe ventricles of brain.
9. Name the five fundamental parts of reflex arc.
10. Describe structure and function of spinal cord.
11. Write any three differences between sympathetic and parasympathetic nervous system.
12. Illustrate cranial nerves.
13. Write note on gustatory receptors.
14. Write short note on heroin.
15. Prove that MRI is a better diagnostic test than CT scan.

SECTION-III: EXTENSIVE QUESTIONS

D. Give detailed answers of the following questions.

1. Describe structure of a typical neuron.
2. Define reflex arc and explain it with example.
3. Write detail note on resting membrane potential.
4. Define synapse and describe structure and write steps of mechanism of synaptic transmission.
5. Write the name of largest part of the brain and explain its structure and function.
6. Describe peripheral nervous system.
7. Write note on olfactory receptors and taste receptors.
8. Write a comprehensive note on nicotine and cannabis.
9. Describe stroke and meningitis.
10. Write note on electroencephalogram.