Chapter 2
Set, Functions and Groups

Exercise 2.1

Q1. Write the following sets in set builder notation.

i. $A = \{1, \ldots, 100\}$
ii. $B = \{100, 101, \ldots, 200\}$
iii. $C = \{-100, -99, \ldots, -50\}$
iv. $D = \{1, 3, 5, \ldots, 199, \ldots, 999\}$
v. $E = \{\text{Parliament, Lahore, Karachi, Quetta}\}$
vi. $F = \{\text{Banana, Lady, Jack}\}$
vii. $G = \{\text{The set of all rational numbers}\}$
viii. $H = \{\text{The set of all integers between 1 and 3}\}$
ix. $I = \{\text{The set of all integers between - 100 and 1000}\}$

Solution

i. $A = \{1, \ldots, 100\}$
$= \{x : x \leq 1 \leq x \leq 100\}$

ii. $B = \{100, 101, \ldots, 200\}$
$= \{x : x = 100 + 1 \leq x \leq 200\}$

iii. $C = \{-100, -99, \ldots, -50\}$
$= \{x : x = -100 + 1 \leq x \leq -50\}$

iv. $D = \{1, 3, 5, \ldots, 199, \ldots, 999\}$
$= \{x : x = 2n + 1, n \in N, 0 \leq n \leq 499\}$

v. $E = \{\text{Parliament, Lahore, Karachi, Quetta}\}$
$= \{x : x \text{ is the capital of provinces of pakistan}\}$

vi. $F = \{\text{Banana, Lady, Jack}\}$
$= \{x : x \text{ is the fruit name and 1 alphabet name}\}$

vii. $G = \{\text{The set of all rational numbers}\}$
$= \{x : x = \frac{p}{q}, p, q \in Z, q \neq 0\}$
$= \{x : x = \frac{p}{q}, p \text{ and } q \text{ are integers}, q \neq 0\}$

viii. $H = \{\text{The set of all integers between 1 and 3}\}$
$= \{2\}$

$= \{x : x \in Z, 1 < x < 3\}$

ix. $I = \{\text{The set of all integers between - 100 and 1000}\}$
$= \{x : x \in Z, -100 < x < 1000\}$

Q2. Write each of the following sets in the descriptive and tabular forms:

i. $A = \{x : x \in N, x \leq 5\}$
ii. $B = \{x : x \in Z, -3 < x \leq 3\}$
iii. $C = \{x : x \in Z, x^2 = 16\}$
iv. $D = \{x : x \in N, x \text{ is even}\}$
v. $E = \{x : x \in N, x \text{ is odd}\}$
vi. $F = \{x : x \in N, x \text{ is prime}\}$
vii. $G = \{x : x \in N, x \text{ is composite}\}$
viii. $H = \{x : x \in N, x = 2n\}$
ix. $I = \{x : x \in N, x = 2n + 1\}$

Solution

i. $A = \{x : x \in N, x \leq 5\}$
Descriptive form
The set of first five natural numbers
Tabular form
$A = \{1, \ldots, 5\}$

ii. $B = \{x : x \in Z, -3 < x \leq 3\}$
Descriptive form

The set of first two natural numbers between -3 and 3
Tabular form
$B = \{-2, -1, 0, 1, 2, 3\}$

iii. $C = \{x : x \in Z, x^2 = 16\}$
Descriptive form
The set of integers between -5 and 5
Tabular form
$C = \{-4, 4\}$

iv. $D = \{x : x \in N, x \text{ is even}\}$
Descriptive form
The set of all even integers between 1 and 5
Tabular form
$D = \{2, 4\}$

v. $E = \{x : x \in N, x \text{ is odd}\}$
Descriptive form
The set of all prime numbers less than 12
Tabular form
$E = \{1, 3, 5, 7, 9, 11\}$

vi. $F = \{x : x \in N, x \text{ is prime}\}$
Descriptive form
The set of all prime numbers between 4 and 10
Tabular form
$F = \{5, 7\}$

vii. $G = \{x : x \in N, x \text{ is composite}\}$
Descriptive form
The set of composite numbers between 4 and 10
Tabular form
$G = \{6, 8, 9\}$

viii. $H = \{x : x \in N, x = 2n\}$
Descriptive form
The set of even numbers less than 12
Tabular form
$H = \{2, 4, 6, 8, 10\}$

ix. $I = \{x : x \in N, x = 2n + 1\}$
Descriptive form
The set of odd numbers between 4 and 10
Tabular form
$I = \{5, 7, 9\}$

x. $J = \{x : x \in N, x \leq 7\}$
Descriptive form
The set of natural numbers less than or equal to 7
Tabular form
$J = \{1, \ldots, 7\}$

xi. $K = \{x : x \in N, x = 2n + 1\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$K = \{3, 5, 7, \ldots\}$

xii. $L = \{x : x \in N, x \text{ is multiple of } 3\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$L = \{3, 6, 9, \ldots\}$

xiii. $M = \{x : x \in N, x \text{ is multiple of } 4\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$M = \{4, 8, 12, \ldots\}$

xiv. $N = \{x : x \in N, x = n^2\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$N = \{1, 4, 9, \ldots\}$

xv. $O = \{x : x \in N, x = n^3\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$O = \{1, 8, 27, \ldots\}$

xvi. $P = \{x : x \in N, x \text{ is multiple of } 2 \text{ and } 3\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$P = \{6, 12, 18, \ldots\}$

xvii. $Q = \{x : x \in N, x = n + 1\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$Q = \{2, 3, 4, \ldots\}$

xviii. $R = \{x : x \in N, x = n - 1\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$R = \{0, 1, 2, \ldots\}$

xix. $S = \{x : x \in N, x = n^2 + 1\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$S = \{2, 5, 10, \ldots\}$

xx. $T = \{x : x \in N, x = n^3 - 1\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$T = \{0, 7, 26, \ldots\}$

xxi. $U = \{x : x \in N, x = n + 2\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$U = \{3, 4, 5, \ldots\}$

xxii. $V = \{x : x \in N, x = n - 2\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$V = \{-1, 0, 1, \ldots\}$

xxiii. $W = \{x : x \in N, x = n^2 + 2\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$W = \{3, 6, 11, \ldots\}$

xxiv. $X = \{x : x \in N, x = n^3 - 2\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$X = \{-1, 6, 25, \ldots\}$

xxv. $Y = \{x : x \in N, x = n + 3\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$Y = \{4, 5, 6, \ldots\}$

xxvi. $Z = \{x : x \in N, x = n - 3\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$Z = \{-2, -1, 0, \ldots\}$

xxvii. $A = \{x : x \in N, x = n^2 + 3\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$A = \{4, 7, 12, \ldots\}$

xxviii. $B = \{x : x \in N, x = n^3 - 3\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$B = \{-2, 5, 24, \ldots\}$

xxix. $C = \{x : x \in N, x = n + 4\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$C = \{5, 6, 7, \ldots\}$

xxx. $D = \{x : x \in N, x = n - 4\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$D = \{-3, -2, -1, \ldots\}$

xxxi. $E = \{x : x \in N, x = n^2 + 4\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$E = \{5, 8, 13, \ldots\}$

xxxii. $F = \{x : x \in N, x = n^3 - 4\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$F = \{-3, 4, 23, \ldots\}$

xxxiii. $G = \{x : x \in N, x = n + 5\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$G = \{6, 7, 8, \ldots\}$

xxxiv. $H = \{x : x \in N, x = n - 5\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$H = \{-4, -3, -2, \ldots\}$

xxxv. $I = \{x : x \in N, x = n^2 + 5\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$I = \{6, 9, 14, \ldots\}$

xxxvi. $J = \{x : x \in N, x = n^3 - 5\}$
Descriptive form
The set of natural numbers in which $n \in N$
Tabular form
$J = \{-4, 3, 22, \ldots\}$

Q3. Which of the following sets are finite sets and which of these are infinite?

i. **The set of all integers.**
ii. **The set of all numbers in Pakistan.**
iii. **The set of all natural numbers between 2 and 10.**
iv. **The set of all natural numbers between 2 and 10 and 20.**
v. **The set of all natural numbers between 2 and 10 and 20 and 30.**
vi. **The set of all natural numbers between 2 and 10 and 20 and 30 and 40.**
vii. **The set of all natural numbers between 2 and 10 and 20 and 30 and 40 and 50.**
viii. **The set of all the provinces of Pakistan.**
ix. **{1, 2, 3, ..., 1000000}**
x. **{2, 4, 6, ..., 1000000}**
xi. **{1, 3, 5, 7, 9}**
xii. **{1, 4, 9, 16, 25}**
xiii. **{1, 8, 27, 64, 125}**

Solution

i. **The set of all integers.**
Infinite

ii. **The set of all numbers in Pakistan.**
Finite

iii. **The set of all natural numbers between 2 and 10.**
Finite

iv. **The set of all natural numbers between 2 and 10 and 20.**
Finite

v. **The set of all natural numbers between 2 and 10 and 20 and 30.**
Finite

vi. **The set of all natural numbers between 2 and 10 and 20 and 30 and 40.**
Infinite

vii. **The set of all natural numbers between 2 and 10 and 20 and 30 and 40 and 50.**
Infinite

viii. **The set of all the provinces of Pakistan.**
Finite

ix. **{1, 2, 3, ..., 1000000}**
Finite

x. **{2, 4, 6, ..., 1000000}**
Finite

xi. **{1, 3, 5, 7, 9}**
Finite

xii. **{1, 4, 9, 16, 25}**
Finite

xiii. **{1, 8, 27, 64, 125}**
Finite

i. **{a, b, c}**
Proper subsets are:
$\{\}, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}$

ii. **{a, b, c, d}**
Proper subsets are:
$\{\}, \{a\}, \{b\}, \{c\}, \{d\}, \{a, b\}, \{a, c\}, \{a, d\}, \{b, c\}, \{b, d\}, \{c, d\}, \{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}$

iii. **{1, 2, 3, 4, 5}**
Proper subsets are:
$\{\}, \{1\}, \{2\}, \{3\}, \{4\}, \{5\}, \{1, 2\}, \{1, 3\}, \{1, 4\}, \{1, 5\}, \{2, 3\}, \{2, 4\}, \{2, 5\}, \{3, 4\}, \{3, 5\}, \{4, 5\}, \{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 5\}, \{1, 3, 4\}, \{1, 3, 5\}, \{1, 4, 5\}, \{2, 3, 4\}, \{2, 3, 5\}, \{2, 4, 5\}, \{3, 4, 5\}, \{1, 2, 3, 4\}, \{1, 2, 3, 5\}, \{1, 2, 4, 5\}, \{1, 3, 4, 5\}, \{2, 3, 4, 5\}$

iv. **{a, b, c, d, e, f}**
Proper subsets are:
$\{\}, \{a\}, \{b\}, \{c\}, \{d\}, \{e\}, \{f\}, \{a, b\}, \{a, c\}, \{a, d\}, \{a, e\}, \{a, f\}, \{b, c\}, \{b, d\}, \{b, e\}, \{b, f\}, \{c, d\}, \{c, e\}, \{c, f\}, \{d, e\}, \{d, f\}, \{e, f\}, \{a, b, c\}, \{a, b, d\}, \{a, b, e\}, \{a, b, f\}, \{a, c, d\}, \{a, c, e\}, \{a, c, f\}, \{a, d, e\}, \{a, d, f\}, \{a, e, f\}, \{b, c, d\}, \{b, c, e\}, \{b, c, f\}, \{b, d, e\}, \{b, d, f\}, \{b, e, f\}, \{c, d, e\}, \{c, d, f\}, \{c, e, f\}, \{d, e, f\}, \{a, b, c, d\}, \{a, b, c, e\}, \{a, b, c, f\}, \{a, b, d, e\}, \{a, b, d, f\}, \{a, b, e, f\}, \{a, c, d, e\}, \{a, c, d, f\}, \{a, c, e, f\}, \{a, d, e, f\}, \{b, c, d, e\}, \{b, c, d, f\}, \{b, c, e, f\}, \{b, d, e, f\}, \{c, d, e, f\}, \{a, b, c, d, e\}, \{a, b, c, d, f\}, \{a, b, c, e, f\}, \{a, b, d, e, f\}, \{a, c, d, e, f\}, \{b, c, d, e, f\}$

v. **{a}**
Proper subsets are:
$\{\}$

vi. **{a, b}**
Proper subsets are:
$\{\}, \{a\}, \{b\}$

vii. **{a, b, c}**
Proper subsets are:
$\{\}, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}$

viii. **{a, b, c, d}**
Proper subsets are:
$\{\}, \{a\}, \{b\}, \{c\}, \{d\}, \{a, b\}, \{a, c\}, \{a, d\}, \{b, c\}, \{b, d\}, \{c, d\}, \{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}$

ix. **{a, b, c, d, e}**
Proper subsets are:
$\{\}, \{a\}, \{b\}, \{c\}, \{d\}, \{e\}, \{a, b\}, \{a, c\}, \{a, d\}, \{a, e\}, \{b, c\}, \{b, d\}, \{b, e\}, \{c, d\}, \{c, e\}, \{d, e\}, \{a, b, c\}, \{a, b, d\}, \{a, b, e\}, \{a, c, d\}, \{a, c, e\}, \{a, d, e\}, \{b, c, d\}, \{b, c, e\}, \{b, d, e\}, \{c, d, e\}, \{a, b, c, d\}, \{a, b, c, e\}, \{a, b, d, e\}, \{a, c, d, e\}, \{b, c, d, e\}$

Q5. Which of the following statements are true and which of them are false?

i. $\{1, 2, 3\} \subseteq \{1, 2, 4, 5\}$ **False**
ii. $\{1, 2, 3\} \subseteq \{1, 2, 3, 4, 5\}$ **True**
iii. $\{1, 2, 3\} \subseteq \{1, 2, 3\}$ **True**
iv. $\{1, 2, 3\} \subseteq \{3, 2, 1\}$ **True**
v. $\{1, 2, 3\} \subseteq \{3, 2, 1\}$ **True**
vi. $\{1, 2, 3\} \subseteq \{3, 2, 1\}$ **True**

Q6. Which is the number of elements of the power set of each of the following sets?

i. $\{\}$ ii. $\{a\}$ iii. $\{a, b\}$ iv. $\{1, 2, 3\}$ v. $\{a, b, c, d\}$
vi. $\{1, 2, 3, 4, 5\}$ vii. $\{a, b, c, d, e, f\}$ viii. $\{1, 2, 3, 4, 5, 6, 7, 8\}$

Solution

i. $\{\}$
Power set of $\{\}$ is $\{\}$
$n = 0$
$2^n = 2^0 = 1$

ii. $\{a\}$
Power set of $\{a\}$ are $\{\}, \{a\}$
$n = 1$
$2^n = 2^1 = 2$

iii. $\{a, b\}$
Power set of $\{a, b\}$ are $\{\}, \{a\}, \{b\}, \{a, b\}$
$n = 2$
$2^n = 2^2 = 4$

iv. $\{1, 2, 3\}$
Power set of $\{1, 2, 3\}$ are $\{\}, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}$
$n = 3$
$2^n = 2^3 = 8$

v. $\{a, b, c, d\}$
Power set of $\{a, b, c, d\}$ are $\{\}, \{a\}, \{b\}, \{c\}, \{d\}, \{a, b\}, \{a, c\}, \{a, d\}, \{b, c\}, \{b, d\}, \{c, d\}, \{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}, \{a, b, c, d\}$
$n = 4$
$2^n = 2^4 = 16$

vi. $\{1, 2, 3, 4, 5\}$
Power set of $\{1, 2, 3, 4, 5\}$ are $\{\}, \{1\}, \{2\}, \{3\}, \{4\}, \{5\}, \{1, 2\}, \{1, 3\}, \{1, 4\}, \{1, 5\}, \{2, 3\}, \{2, 4\}, \{2, 5\}, \{3, 4\}, \{3, 5\}, \{4, 5\}, \{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 5\}, \{1, 3, 4\}, \{1, 3, 5\}, \{1, 4, 5\}, \{2, 3, 4\}, \{2, 3, 5\}, \{2, 4, 5\}, \{3, 4, 5\}, \{1, 2, 3, 4\}, \{1, 2, 3, 5\}, \{1, 2, 4, 5\}, \{1, 3, 4, 5\}, \{2, 3, 4, 5\}, \{1, 2, 3, 4, 5\}$
$n = 5$
$2^n = 2^5 = 32$

vii. $\{a, b, c, d, e, f\}$
Power set of $\{a, b, c, d, e, f\}$ are $\{\}, \{a\}, \{b\}, \{c\}, \{d\}, \{e\}, \{f\}, \{a, b\}, \{a, c\}, \{a, d\}, \{a, e\}, \{a, f\}, \{b, c\}, \{b, d\}, \{b, e\}, \{b, f\}, \{c, d\}, \{c, e\}, \{c, f\}, \{d, e\}, \{d, f\}, \{e, f\}, \{a, b, c\}, \{a, b, d\}, \{a, b, e\}, \{a, b, f\}, \{a, c, d\}, \{a, c, e\}, \{a, c, f\}, \{a, d, e\}, \{a, d, f\}, \{a, e, f\}, \{b, c, d\}, \{b, c, e\}, \{b, c, f\}, \{b, d, e\}, \{b, d, f\}, \{b, e, f\}, \{c, d, e\}, \{c, d, f\}, \{c, e, f\}, \{d, e, f\}, \{a, b, c, d\}, \{a, b, c, e\}, \{a, b, c, f\}, \{a, b, d, e\}, \{a, b, d, f\}, \{a, b, e, f\}, \{a, c, d, e\}, \{a, c, d, f\}, \{a, c, e, f\}, \{a, d, e, f\}, \{b, c, d, e\}, \{b, c, d, f\}, \{b, c, e, f\}, \{b, d, e, f\}, \{c, d, e, f\}, \{a, b, c, d, e\}, \{a, b, c, d, f\}, \{a, b, c, e, f\}, \{a, b, d, e, f\}, \{a, c, d, e, f\}, \{b, c, d, e, f\}, \{a, b, c, d, e, f\}$
$n = 6$
$2^n = 2^6 = 64$

viii. $\{1, 2, 3, 4, 5, 6, 7, 8\}$
Power set of $\{1, 2, 3, 4, 5, 6, 7, 8\}$ are $\{\}, \{1\}, \{2\}, \{3\}, \{4\}, \{5\}, \{6\}, \{7\}, \{8\}, \{1, 2\}, \{1, 3\}, \{1, 4\}, \{1, 5\}, \{1, 6\}, \{1, 7\}, \{1, 8\}, \{2, 3\}, \{2, 4\}, \{2, 5\}, \{2, 6\}, \{2, 7\}, \{2, 8\}, \{3, 4\}, \{3, 5\}, \{3, 6\}, \{3, 7\}, \{3, 8\}, \{4, 5\}, \{4, 6\}, \{4, 7\}, \{4, 8\}, \{5, 6\}, \{5, 7\}, \{5, 8\}, \{6, 7\}, \{6, 8\}, \{7, 8\}, \{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 5\}, \{1, 2, 6\}, \{1, 2, 7\}, \{1, 2, 8\}, \{1, 3, 4\}, \{1, 3, 5\}, \{1, 3, 6\}, \{1, 3, 7\}, \{1, 3, 8\}, \{1, 4, 5\}, \{1, 4, 6\}, \{1, 4, 7\}, \{1, 4, 8\}, \{1, 5, 6\}, \{1, 5, 7\}, \{1, 5, 8\}, \{1, 6, 7\}, \{1, 6, 8\}, \{1, 7, 8\}, \{2, 3, 4\}, \{2, 3, 5\}, \{2, 3, 6\}, \{2, 3, 7\}, \{2, 3, 8\}, \{2, 4, 5\}, \{2, 4, 6\}, \{2, 4, 7\}, \{2, 4, 8\}, \{2, 5, 6\}, \{2, 5, 7\}, \{2, 5, 8\}, \{2, 6, 7\}, \{2, 6, 8\}, \{2, 7, 8\}, \{3, 4, 5\}, \{3, 4, 6\}, \{3, 4, 7\}, \{3, 4, 8\}, \{3, 5, 6\}, \{3, 5, 7\}, \{3, 5, 8\}, \{3, 6, 7\}, \{3, 6, 8\}, \{3, 7, 8\}, \{4, 5, 6\}, \{4, 5, 7\}, \{4, 5, 8\}, \{4, 6, 7\}, \{4, 6, 8\}, \{4, 7, 8\}, \{5, 6, 7\}, \{5, 6, 8\}, \{5, 7, 8\}, \{6, 7, 8\}, \{1, 2, 3, 4\}, \{1, 2, 3, 5\}, \{1, 2, 3, 6\}, \{1, 2, 3, 7\}, \{1, 2, 3, 8\}, \{1, 2, 4, 5\}, \{1, 2, 4, 6\}, \{1, 2, 4, 7\}, \{1, 2, 4, 8\}, \{1, 2, 5, 6\}, \{1, 2, 5, 7\}, \{1, 2, 5, 8\}, \{1, 2, 6, 7\}, \{1, 2, 6, 8\}, \{1, 2, 7, 8\}, \{1, 3, 4, 5\}, \{1, 3, 4, 6\}, \{1, 3, 4, 7\}, \{1, 3, 4, 8\}, \{1, 3, 5, 6\}, \{1, 3, 5, 7\}, \{1, 3, 5, 8\}, \{1, 3, 6, 7\}, \{1, 3, 6, 8\}, \{1, 3, 7, 8\}, \{1, 4, 5, 6\}, \{1, 4, 5, 7\}, \{1, 4, 5, 8\}, \{1, 4, 6, 7\}, \{1, 4, 6, 8\}, \{1, 4, 7, 8\}, \{1, 5, 6, 7\}, \{1, 5, 6, 8\}, \{1, 5, 7, 8\}, \{1, 6, 7, 8\}, \{2, 3, 4, 5\}, \{2, 3, 4, 6\}, \{2, 3, 4, 7\}, \{2, 3, 4, 8\}, \{2, 3, 5, 6\}, \{2, 3, 5, 7\}, \{2, 3, 5, 8\}, \{2, 3, 6, 7\}, \{2, 3, 6, 8\}, \{2, 3, 7, 8\}, \{2, 4, 5, 6\}, \{2, 4, 5, 7\}, \{2, 4, 5, 8\}, \{2, 4, 6, 7\}, \{2, 4, 6, 8\}, \{2, 4, 7, 8\}, \{2, 5, 6, 7\}, \{2, 5, 6, 8\}, \{2, 5, 7, 8\}, \{2, 6, 7, 8\}, \{3, 4, 5, 6\}, \{3, 4, 5, 7\}, \{3, 4, 5, 8\}, \{3, 4, 6, 7\}, \{3, 4, 6, 8\}, \{3, 4, 7, 8\}, \{3, 5, 6, 7\}, \{3, 5, 6, 8\}, \{3, 5, 7, 8\}, \{3, 6, 7, 8\}, \{4, 5, 6, 7\}, \{4, 5, 6, 8\}, \{4, 5, 7, 8\}, \{4, 6, 7, 8\}, \{5, 6, 7, 8\}, \{1, 2, 3, 4, 5\}, \{1, 2, 3, 4, 6\}, \{1, 2, 3, 4, 7\}, \{1, 2, 3, 4, 8\}, \{1, 2, 3, 5, 6\}, \{1, 2, 3, 5, 7\}, \{1, 2, 3, 5, 8\}, \{1, 2, 3, 6, 7\}, \{1, 2, 3, 6, 8\}, \{1, 2, 3, 7, 8\}, \{1, 2, 4, 5, 6\}, \{1, 2, 4, 5, 7\}, \{1, 2, 4, 5, 8\}, \{1, 2, 4, 6, 7\}, \{1, 2, 4, 6, 8\}, \{1, 2, 4, 7, 8\}, \{1, 2, 5, 6, 7\}, \{1, 2, 5, 6, 8\}, \{1, 2, 5, 7, 8\}, \{1, 2, 6, 7, 8\}, \{1, 3, 4, 5, 6\}, \{1, 3, 4, 5, 7\}, \{1, 3, 4, 5, 8\}, \{1, 3, 4, 6, 7\}, \{1, 3, 4, 6, 8\}, \{1, 3, 4, 7, 8\}, \{1, 3, 5, 6, 7\}, \{1, 3, 5, 6, 8\}, \{1, 3, 5, 7, 8\}, \{1, 3, 6, 7, 8\}, \{1, 4, 5, 6, 7\}, \{1, 4, 5, 6, 8\}, \{1, 4, 5, 7, 8\}, \{1, 4, 6, 7, 8\}, \{1, 5, 6, 7, 8\}, \{2, 3, 4, 5, 6\}, \{2, 3, 4, 5, 7\}, \{2, 3, 4, 5, 8\}, \{2, 3, 4, 6, 7\}, \{2, 3, 4, 6, 8\}, \{2, 3, 4, 7, 8\}, \{2, 3, 5, 6, 7\}, \{2, 3, 5, 6, 8\}, \{2, 3, 5, 7, 8\}, \{2, 3, 6, 7, 8\}, \{2, 4, 5, 6, 7\}, \{2, 4, 5, 6, 8\}, \{2, 4, 5, 7, 8\}, \{2, 4, 6, 7, 8\}, \{2, 5, 6, 7, 8\}, \{3, 4, 5, 6, 7\}, \{3, 4, 5, 6, 8\}, \{3, 4, 5, 7, 8\}, \{3, 4, 6, 7, 8\}, \{3, 5, 6, 7, 8\}, \{4, 5, 6, 7, 8\}, \{1, 2, 3, 4, 5, 6\}, \{1, 2, 3, 4, 5, 7\}, \{1, 2, 3, 4, 5, 8\}, \{1, 2, 3, 4, 6, 7\}, \{1, 2, 3, 4, 6, 8\}, \{1, 2, 3, 4, 7, 8\}, \{1, 2, 3, 5, 6, 7\}, \{1, 2, 3, 5, 6, 8\}, \{1, 2, 3, 5, 7, 8\}, \{1, 2, 3, 6, 7, 8\}, \{1, 2, 4, 5, 6, 7\}, \{1, 2, 4, 5, 6, 8\}, \{1, 2, 4, 5, 7, 8\}, \{1, 2, 4, 6, 7, 8\}, \{1, 2, 5, 6, 7, 8\}, \{1, 3, 4, 5, 6, 7\}, \{1, 3, 4, 5, 6, 8\}, \{1, 3, 4, 5, 7, 8\}, \{1, 3, 4, 6, 7, 8\}, \{1, 3, 5, 6, 7, 8\}, \{1, 4, 5, 6, 7, 8\}, \{2, 3, 4, 5, 6, 7\}, \{2, 3, 4, 5, 6, 8\}, \{2, 3, 4, 5, 7, 8\}, \{2, 3, 4, 6, 7, 8\}, \{2, 3, 5, 6, 7, 8\}, \{2, 4, 5, 6, 7, 8\}, \{3, 4, 5, 6, 7, 8\}, \{1, 2, 3, 4, 5, 6, 7\}, \{1, 2, 3, 4, 5, 6, 8\}, \{1, 2, 3, 4, 5, 7, 8\}, \{1, 2, 3, 4, 6, 7, 8\}, \{1, 2, 3, 5, 6, 7, 8\}, \{1, 2, 4, 5, 6, 7, 8\}, \{1, 3, 4, 5, 6, 7, 8\}, \{2, 3, 4, 5, 6, 7, 8\}, \{1, 2, 3, 4, 5, 6, 7, 8\}$
$n = 8$
$2^n = 2^8 = 256$

Q7. Which of the following statements are true and which of them are false?

i. $\{a, b, c\} = \{a, b, c\}$ **True**
ii. $\{a, b, c\} = \{a, b, c, d\}$ **False**
iii. $\{a, b, c\} = \{a, b, c, a\}$ **True**
iv. $\{a, b, c\} = \{a, b, c, a, b\}$ **True**
v. $\{a, b, c\} = \{a, b, c, a, b, c\}$ **True**
vi. $\{a, b, c\} = \{a, b, c, a, b, c, a\}$ **True**

Q8. Which sets of sets are equivalent? Which of them are not equivalent?

i. **{a, b, c}, {1, 2, 3}**
ii. **The set of all the prime numbers, {2, 3, 5, 7, ...}, {1, 2, 3, ...}**
iii. **Set of angles of a quadrilateral, {4, 5, 6, 7}**
iv. **Set of the sides of a quadrilateral, {4, 5, 6, 7}**
v. **Set of the angles of a pentagon, {4, 5, 6, 7, 8}**
vi. **Set of the sides of a pentagon, {4, 5, 6, 7, 8}**
vii. **Set of the angles of a hexagon, {4, 5, 6, 7, 8, 9}**
viii. **Set of the sides of a hexagon, {4, 5, 6, 7, 8, 9}**
ix. **{a, b, c, d, e, f, g}, {1, 2, 3, 4, 5, 6, 7}**
x. **{a, b, c, d, e, f, g}, {1, 2, 3, 4, 5, 6, 7}**

Solution

i. **{a, b, c}, {1, 2, 3}**
Equivalent sets because each set has 3 elements.

ii. **The set of the first 4 prime numbers, {2, 3, 5, 7}, {1, 2, 3, 4}**
Equivalent sets because each set have some 4 elements.

iii. **Set of angles of a quadrilateral, {4, 5, 6, 7}**
The set of the angles of the same hexagon,
Equivalent sets because each set have the same elements.

iv. **Set of the sides of a quadrilateral, {4, 5, 6, 7}**
The set of the angles of the same hexagon,
Equivalent sets because each set have the same elements.

v. **{a, b, c, d, e}, {1, 2, 3, 4, 5}**
Equivalent sets because they have one to one (1-1) correspondence between them.

vi. **{a, b, c, d, e}, {1, 2, 3, 4, 5}**
Equivalent sets because they have one to one (1-1) correspondence between them.

vii. **{a, b, c, d, e, f, g}, {1, 2, 3, 4, 5, 6, 7}**
7 am's do the 6x7 of 1st birthday, so they are not equivalent set.

Exercise 2.2

Q1. Exhibit A ∩ B and A ∩ B by Venn Diagram in the following cases:

i. $A \subset B$ ii. $A \subset B$ iii. $A \subset B$ iv. $A \subset B$

iv. A and B are disjoint sets. A and B are overlapping sets.

Solution:

i. $A \subset B$

a. $A \subset B$



b. $A \subset B$




ii. $A \subset B$

a. $A \subset B$



b. $A \subset B$



iii. $A \subset B$

$\Rightarrow A \cap B = A$

$\Rightarrow A$



iv. A and B are disjoint sets.

a. $A \cap B$



b. $A \cap B$



v. A and B are overlapping sets.

a. $A \cap B$

b. $A \cap B$

Q2 (Exhibit A ∩ B and B ∩ A by Venn Diagram when:

i. A and B are overlapping sets. ii. $A \subset B$ iii. $B \subset A$

i. A and B are overlapping sets.

a. $A \cap B$

b. $A \cap B$

ii. $A \subset B$

a. $A \cap B$

b. $A \cap B$

iii. $B \subset A$

a. $A \cap B$

b. $A \cap B$

Q3. Under what conditions on A and B are the following statement true?

i. $A \cup B = A$ ii. $A \cup B = B$ iii. $A \cap B = A$

iv. $A \cap B = B$ v. $A \setminus B = \emptyset$ vi. $A \subset B \Rightarrow A \setminus B = \emptyset$

vii. $A \setminus B = A \Rightarrow A \cap B = \emptyset$ viii. $A \subset B \Rightarrow A \setminus (B \setminus A) = A$

ix. $A \cup B = B$ x. $A \cap B = A \Rightarrow A \subset B$ xi. $A \setminus B = B \setminus A \Rightarrow A = B$

xii. $A \Delta B = A \cup B \Rightarrow A \cap B = \emptyset$ xiii. $A \setminus (B \setminus A) = A$ xiv. $A \setminus (A \setminus B) = A \cap B$

Solution:

i. $A \cup B = A$

If is true if $B \subseteq A$.

ii. $A \cup B = B$

If is true if $A \subseteq B$.

iii. $A \cap B = A$

If is true if $A \subseteq B$ or $A = B$

iv. $A \cap B = B$

If is true if $B \subseteq A$ or $A = B$

v. $A \setminus B = \emptyset$

If is true if $A \subseteq B$ or $A = B$

vi. $A \subset B \Rightarrow A \setminus B = \emptyset$

If is true because it is commutative property w.r.t union.

vii. $A \setminus B = A \Rightarrow A \cap B = \emptyset$

If is true if $A \cap B = \emptyset$ or $A = \emptyset$

viii. $A \subset B \Rightarrow A \setminus (B \setminus A) = A$

If is true if $A \cap B = \emptyset$ or $A = \emptyset$

ix. $A \cup B = B$

If is true if $A \subseteq B$.

x. $A \cap B = A \Rightarrow A \subset B$

If is true if $A \subseteq B$.

xi. $A \setminus B = B \setminus A \Rightarrow A = B$

If is true because it is commutative property w.r.t union.

xii. $A \Delta B = A \cup B \Rightarrow A \cap B = \emptyset$

If is true if $A \cap B = \emptyset$ or $A = \emptyset$

xiii. $A \setminus (B \setminus A) = A$

If is true if $A \cap B = \emptyset$ or $A = \emptyset$

xiv. $A \setminus (A \setminus B) = A \cap B$

If is true if $A \cap B = \emptyset$ or $A = \emptyset$

Q4. Let $S = \{1,2,3,4,5,6,7,8,9,10\}$, $A = \{2,4,6,8,10\}$, $B = \{1,2,3,4,5\}$ $C = \{1,3,5,7,9\}$

List the members of each of the following sets:

i. $A \cup S$ ii. $B \cap S$ iii. $A \cup S$ iv. $A \cap B$ v. $A \cup C$ vi. $A \cup C$ vii. $A \cup C$ viii. $B \cap (C \cup S)$

Solution:

i. $A \cup S$

$= \{1,2,3,.....10\} - \{2,4,6,8,10\}$

$= \{1,3,5,7,9\}$

Hence $A \cup S = \{1,3,5,7,9\}$

ii. $B \cap S$

$= \{1,2,3,.....10\} - \{1,3,5,7,9\}$

$= \{2,4,6,8,10\}$

Hence $B \cap S = \{2,4,6,8,10\}$

iii. $A \cup S$

$= \{2,4,6,8,10\} \cup \{1,3,5,7,9\}$

$= \{1,2,3,4,5,6,7,8,9,10\}$

Hence $A \cup S = \{1,2,3,4,5,6,7,8,9,10\}$

iv. $A \cap B$

$= \{2,4,6,8,10\} - \{1,3,5,7,9\}$

$= \phi$

Hence $A \cap B = \phi$

v. $A \cup C$

$= \{2,4,6,8,10\} \cup \{1,3,5,7,9\}$

$= \phi$

Hence $A \cap B = \phi$

vi. $A \cup C$

$A' = U-A$

$= \{1,2,3,4,5,6,7,8,9,10\} - \{2,4,6,8,10\}$

$= \{1,3,5,7,9\}$

$C' = U-C$

$= \{1,2,3,4,5,6,7,8,9,10\} - \{1,3,5,7,9\}$

$= \{2,4,6,8,10\}$

$A' \cup C' = A \cap C$ Hence $\{2,4,6,8,10\}$

$= \{1,2,3,4,5,6,7,8,9,10\}$

Hence $A' \cup C' = \{1,2,3,4,5,6,7,8,9,10\}$

vii. $A' \cup C$

$A' = U-A$

$= \{1,2,3,4,5,6,7,8,9,10\} - \{2,4,6,8,10\}$

$= \{1,3,5,7,9\}$

$A' \cup C = \{1,3,5,7,9\} \cup \{1,3,5,7,9\}$

Hence $A' \cup C = \{1,3,5,7,9\}$

viii. $B \cap (C \cup S)$

$= \{1,2,3,4,5\} \cap \{1,3,5,7,9,10\}$

$= \phi$

Hence $B \cap (C \cup S) = \phi$

Q5. Using the Venn diagram if necessary, find the single set equal to the following.

i. $A \cup A'$ ii. $A \cap S$ iii. $A \cup S$ iv. $A \cup \phi$ v. $A \cap \phi$

Solution:

i. $A \cup A'$

$= U-A$

ii. $A \cap S$

iii. $A \cup S$

iv. $A \cup \phi$

v. $A \cap \phi$

Q6. Use Venn diagram to verify the following:

i. $A \cap A = A$ ii. $A \cup A = A$ iii. $(A \cap B)' = A' \cup B'$

Solution:

i. $A \cap A = A$

L.H.S

A & A are overlapping sets

R.H.S

$A \cap B$

$B = A - A$

$\Rightarrow A \cap B'$

Hence Proved

ii. $(A \cap B)' = A' \cup B'$

L.H.S

Where A and B are overlapping sets

$A \cap B$

$(A \cap B)'$

$= U - (A \cap B)$

$(A \cap B)' = U$

R.H.S

Hence Proved

Exercise 2.3

Q1. Identify the following statements as simple or compound and also find the number of propositions.
(i) 2 + 3 = 5 and 5 is a prime number. (ii) 7 > 10.
(iii) Ram is a boy or Shyam is a girl. (iv) 2 is a prime number.
Answer

- **(i) 2 + 3 = 5 and 5 is a prime number.**
  Statement 1 : 2 + 3 = 5
  Statement 2 : 5 is a prime number.
  Connective used : and
  Compound statement
  Number of propositions = 2
- **(ii) 7 > 10**
  Statement : 7 > 10
  Simple statement
  Number of propositions = 1
- **(iii) Ram is a boy or Shyam is a girl.**
  Statement 1 : Ram is a boy.
  Statement 2 : Shyam is a girl.
  Connective used : or
  Compound statement
  Number of propositions = 2
- **(iv) 2 is a prime number.**
  Statement : 2 is a prime number.
  Simple statement
  Number of propositions = 1

Q2. Identify the connective in the following statements and write the component statements.

- **(i) 2 is a prime number or 2 is an even prime number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 2 is an even prime number.
  Connective used : or
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
- **(iii) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
- **(iv) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
- **(v) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or

Q3. Write the negation of the following statements.

- **(i) 2 is a prime number.**
  Negation : 2 is not a prime number.
- **(ii) 3 is an odd number.**
  Negation : 3 is not an odd number.
- **(iii) Delhi is the capital of India.**
  Negation : Delhi is not the capital of India.
- **(iv) 7 is greater than 5.**
  Negation : 7 is not greater than 5.
- **(v) 7 is less than 10.**
  Negation : 7 is not less than 10.

Q4. Write the negation of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Negation : 2 is not a prime number or 3 is not an odd number.
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Negation : Delhi is not in India and not in China.
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Negation : 7 ≤ 5 or 7 ≥ 10
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Negation : 7 ≤ 5 and 7 ≥ 10

Q5. Write the truth value of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T

Q6. Write the truth value of the negation of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
  Negation : 2 is not a prime number or 3 is not an odd number.
  Truth value of negation : F
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
  Negation : Delhi is not in India and not in China.
  Truth value of negation : F
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
  Negation : 7 ≤ 5 or 7 ≥ 10
  Truth value of negation : F
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T
  Negation : 7 ≤ 5 and 7 ≥ 10
  Truth value of negation : F

Q7. Identify the connective in each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or

Q8. Write the truth value of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T

Q9. Write the truth value of the negation of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
  Negation : 2 is not a prime number or 3 is not an odd number.
  Truth value of negation : F
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
  Negation : Delhi is not in India and not in China.
  Truth value of negation : F
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
  Negation : 7 ≤ 5 or 7 ≥ 10
  Truth value of negation : F
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T
  Negation : 7 ≤ 5 and 7 ≥ 10
  Truth value of negation : F

Q10. Identify the connective in each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or

Q11. Write the truth value of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T

Q12. Write the truth value of the negation of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
  Negation : 2 is not a prime number or 3 is not an odd number.
  Truth value of negation : F
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
  Negation : Delhi is not in India and not in China.
  Truth value of negation : F
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
  Negation : 7 ≤ 5 or 7 ≥ 10
  Truth value of negation : F
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T
  Negation : 7 ≤ 5 and 7 ≥ 10
  Truth value of negation : F

Q13. Identify the connective in each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or

Q14. Write the truth value of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T

Q15. Write the truth value of the negation of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
  Negation : 2 is not a prime number or 3 is not an odd number.
  Truth value of negation : F
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
  Negation : Delhi is not in India and not in China.
  Truth value of negation : F
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
  Negation : 7 ≤ 5 or 7 ≥ 10
  Truth value of negation : F
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T
  Negation : 7 ≤ 5 and 7 ≥ 10
  Truth value of negation : F

Q16. Identify the connective in each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or

Q17. Write the truth value of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T

Q18. Write the truth value of the negation of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
  Negation : 2 is not a prime number or 3 is not an odd number.
  Truth value of negation : F
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
  Negation : Delhi is not in India and not in China.
  Truth value of negation : F
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
  Negation : 7 ≤ 5 or 7 ≥ 10
  Truth value of negation : F
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T
  Negation : 7 ≤ 5 and 7 ≥ 10
  Truth value of negation : F

Q19. Identify the connective in each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or

Q20. Write the truth value of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T

Q21. Write the truth value of the negation of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
  Negation : 2 is not a prime number or 3 is not an odd number.
  Truth value of negation : F
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
  Negation : Delhi is not in India and not in China.
  Truth value of negation : F
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
  Negation : 7 ≤ 5 or 7 ≥ 10
  Truth value of negation : F
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T
  Negation : 7 ≤ 5 and 7 ≥ 10
  Truth value of negation : F

Q22. Identify the connective in each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or

Q23. Write the truth value of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T

Q24. Write the truth value of the negation of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
  Negation : 2 is not a prime number or 3 is not an odd number.
  Truth value of negation : F
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
  Negation : Delhi is not in India and not in China.
  Truth value of negation : F
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
  Negation : 7 ≤ 5 or 7 ≥ 10
  Truth value of negation : F
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T
  Negation : 7 ≤ 5 and 7 ≥ 10
  Truth value of negation : F

Q25. Identify the connective in each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or

Q26. Write the truth value of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T

Q27. Write the truth value of the negation of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
  Negation : 2 is not a prime number or 3 is not an odd number.
  Truth value of negation : F
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
  Negation : Delhi is not in India and not in China.
  Truth value of negation : F
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
  Negation : 7 ≤ 5 or 7 ≥ 10
  Truth value of negation : F
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T
  Negation : 7 ≤ 5 and 7 ≥ 10
  Truth value of negation : F

Q28. Identify the connective in each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or

Q29. Write the truth value of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T

Q30. Write the truth value of the negation of each of the following statements.

- **(i) 2 is a prime number and 3 is an odd number.**
  Statement 1 : 2 is a prime number.
  Statement 2 : 3 is an odd number.
  Connective used : and
  Truth value : T
  Negation : 2 is not a prime number or 3 is not an odd number.
  Truth value of negation : F
- **(ii) Delhi is in India or in China.**
  Statement 1 : Delhi is in India.
  Statement 2 : Delhi is in China.
  Connective used : or
  Truth value : T
  Negation : Delhi is not in India and not in China.
  Truth value of negation : F
- **(iii) 7 > 5 and 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : and
  Truth value : T
  Negation : 7 ≤ 5 or 7 ≥ 10
  Truth value of negation : F
- **(iv) 7 > 5 or 7 < 10.**
  Statement 1 : 7 > 5
  Statement 2 : 7 < 10
  Connective used : or
  Truth value : T
  Negation : 7 ≤ 5 and 7 ≥ 10
  Truth value of negation : F

Exercise 2.4

Q1. Write the converse, inverse and contra positive of the following conditional:

i) $\neg p \to q$ ii) $q \to p$ iii) $\neg p \to \neg q$ iv) $\neg q \to \neg p$

Solution:

i) $\neg p \to q$

		Converse	Inverse	Contra Positive
i)	$\neg p \to q$	$q \to \neg p$	$p \to \neg q$	$\neg p \to p$

ii) $q \to p$

		Converse	Inverse	Contra Positive
ii)	$q \to p$	$p \to q$	$\neg q \to \neg p$	$\neg p \to \neg q$

iii) $\neg p \to \neg q$

		Converse	Inverse	Contra Positive
iii)	$\neg p \to \neg q$	$\neg p \to \neg q$	$p \to q$	$q \to p$

iv) $\neg q \to \neg p$

		Converse	Inverse	Contra Positive
iv)	$\neg q \to \neg p$	$\neg p \to \neg q$	$q \to p$	$p \to q$

Q2. Construct truth tables for the following statements.

i. $(p \to \neg p) \vee (p \to q)$

ii. $(p \wedge \neg p) \to q$

iii. $(\neg(p \to q) \leftrightarrow (p \wedge \neg q)$

Solution:

i. $(p \to \neg p) \vee (p \to q)$

p	q	$\neg p$	$\neg q$	$p \to \neg p$	$p \to q$	$(p \to \neg p) \vee (p \to q)$
T	T	F	F	F	T	T
T	F	F	T	F	F	F
F	T	T	F	T	T	T
F	F	T	T	T	T	T

ii. $(p \wedge \neg p) \to q$

p	q	$\neg p$	$p \wedge \neg p$	$(p \wedge \neg p) \to q$
T	T	F	T	T
T	F	F	F	T
F	T	T	T	T
F	F	T	F	T

iii. $(\neg(p \to q) \leftrightarrow (p \wedge \neg q)$

p	q	$\neg p$	$(p \to q$	$\neg(p \to q)$	$p \wedge \neg q$	$(\neg(p \to q)$ $\to \neg(p \wedge q)$	$(\neg(p \to q)$ $\to \neg(p \wedge q)$	$(\neg(p \to q)$ $\leftrightarrow \neg(p \wedge q)$

T	T	F	T	F	F	T	T	T
T	F	F	F	T	T	T	T	T
F	T	T	T	F	F	T	T	T
F	F	T	T	F	F	T	T	T

Q3. Show that each of the following statements is a tautology.

i. $(p \wedge q) \to p$

ii. $p \to (p \vee q)$

iii. $\neg(p \to q) \to p)$

iv. $\neg q \wedge (p \to q) \to \neg p$

`olution:

i. $(p \wedge q) \to p$

P	q	$p \wedge q$	$(p \wedge q) \to p$
T	T	T	T
T	F	F	T
F	T	F	T
F	F	F	T

ii. $p \to (p \vee q)$

P	q	$p \vee q$	$p \to (p \vee q)$
T	T	T	T
T	F	T	T
F	T	T	T
F	F	F	T

iii. $\neg(p \to q) \to p$

P	q	$\neg(p \to q)$	$\neg(p \to q) \to p$
T	T	F	T
.			
F	F	F	T

iv. $\neg q \wedge (p \to q) \to \neg p$

p	q	$\neg p$	$\neg q$	$p \to q$	$\neg q \wedge (p \to q)$	$\neg q \wedge (p \to q) \to \neg p$
T	T	F	F	T	F	T
T	F	F	T	F	F	T
F	T	T	F	T	F	T
F	F	T	T	T	T	T

Q4. Determine whether each of the following is a tautology, a contingency or an absurdity.

i. $p \wedge \neg p$

ii. $p \to (p \to q)$

iii. $q \vee (\neg q \vee p))$

Solution:

i. $p \wedge \neg p$

p	$\neg p$	$p \wedge \neg p$
T	F	F
F	T	F

ii. $p \to (q \to p)$

P	q	$q \to p$	$p \to (q \to p)$
T	T	T	T
T	F	T	T
F	T	F	T
F	F	T	T

iii. $q \vee (\neg q \vee p))$

P	Q	$\neg q$	$\neg q \vee p$	$q \vee (\neg q \vee p)$
T	T	F	T	T
T	F	T	T	T
F	T	F	F	T
F	F	T	T	T

Q5. Prove that $p \vee (\neg p \wedge \neg q) \vee (p \wedge q) = p \vee (\neg p \wedge \neg q)$

Solution:

L.H.S = $p \vee (\neg p \wedge \neg q) \vee (p \wedge q)$

T	F	F	T	F	F	T	T
F	T	T	F	F	F	F	F
F	F	T	T	T	F	T	T

R.H.S = $p \vee (\neg p \wedge \neg q)$

p	q	$\neg p$	$\neg q$	$\neg p \wedge \neg q$	$p \vee (\neg p \wedge \neg q)$
T	T	F	F	F	T
T	F	F	T	F	T
F	T	T	F	F	F

F	F	T	T	T	T

So, L.H.S = R.H.S

Hence Prove.

Exercise 2.5

Q1. Convert the following theorems to logical and prove them by constructing truth tables.

i. **(A∩B)' = A'∪B'** ii. **(A∪B)∩C = (A∪B)∪C**

iii. **(A∩B)∩C = A∩(B∩C)** iv. **A∪(B∩C) = (A∪B) ∩ (A∪C)**

Solution:

i. **(A∩B)' = A'∪B'**

L.H.S $(A \cap B)' \rightarrow \sim(p \wedge q)$

P	Q	$p \wedge q$	$\sim(p \wedge q)$
F	F	F	T

R.H.S $A' \cup B' \rightarrow (\sim p \vee \sim q)$

P	Q	$\sim p$	$\sim q$	$\sim p \vee \sim q$
T	T	F	F	F
T	F	F	T	T
F	T	T	F	T

F	F	T	T	T

ii. **(A∪B)∩C = (A∪B)∪C**

L.H.S $(A \cup B) \cap C \rightarrow (p \vee q) \vee r$

p	Q	r	$p \vee q$	$(p \vee q) \vee r$
T	T	T	T	T
T	T	F	T	T
T	F	T	T	T
F	F	T	T	T
F	F	F	F	F

R.H.S $(A \cup B) \cup C \rightarrow p \vee (q \vee r)$

p	Q	r	$q \vee r$	$p \vee (q \vee r)$
T	T	T	T	T
T	T	F	T	T
T	F	T	T	T

T	F	F	F	T
F	T	T	T	T
F	T	F	T	T
F	F	T	T	T
F	F	F	F	F

iii. **(A∩B)∩C = A∩(B∩C)**

L.H.S $(A \cap B) \cap C \rightarrow (p \wedge q) \wedge r$

T	F	T	F	F
T	F	F	F	F
F	T	T	F	F
F	T	F	F	F
F	F	T	F	F
F	F	F	F	F

R.H.S $A \cap (B \cap C) \rightarrow p \wedge (q \wedge r)$

P	q	r	$q \wedge r$	$p \wedge (q \wedge r)$
T	T	T	T	T
T	T	F	F	F
T	F	T	F	F
T	F	F	F	F
F	T	T	T	F
F	T	F	F	F
F	F	T	F	F
F	F	F	F	F

iv. **A∪(B∩C) = (A∪B) ∩ (A∪C)**

L.H.S $A \cup (B \cap C) \rightarrow p \vee (q \wedge r)$

p	q	r	$q \wedge r$	$p \vee (q \wedge r)$
T	T	T	T	T
T	T	F	F	T
T	F	T	F	T
T	F	F	F	T
F	T	T	T	T
F	T	F	F	F

F	F	T	F	F
F	F	F	F	F

R.H.S $(A \cup B) \cap (A \cup C) \rightarrow (p \vee q) \wedge (p \vee r)$

p	Q	r	$p \vee q$	$p \vee r$	$(p \vee q) \wedge (p \vee r)$
T	T	T	T	T	T
T	T	F	T	T	T
T	F	T	T	T	T
T	F	F	T	T	T
F	F	T	F	F	F
F	F	F	F	F	F

L.H.S = R.H.S

Hence Proved

Exercise 2.6

Q1. For A = {1,2,3,4}, find the following relations in A. State the domain and range of each relation. Also draw the graph of each.

i. {(x,y) | y = x} **ii. {(x,y) | y=x + 1}** **iii. {(x,y) | x+y<5}**

iv. {(x,y) | x+y=10}

Solution:

i. {(x,y) | y = x}


Where, A = {1,2,3,4}

Sc r = {(1,1), (2,2), (3,3),(4,4)

and

Dom (r) = {1,2,3,4}


And range (r) = {1,2,3,4}



ii. {(x,y) | y=x + 1}; Where A = {1,2,3,4}

r = {(1,4), (2,3), (3,2), (3,1)}

So, Dom (r) = {1,2,3,4} = {1,2,3,4}




iii. {(x,y) | x+y<5}, where A = {1,2,3,4}

r ={(1,1),(1,2),(1,3),(2,1), (2,2), (3,1)}

So, Dom(r)= {1,2,3}

& range(r)= {1,2,3}




iv. {(x,y) | x+y=5}

r ={(2,4), (3,3), (3,4), (4,2), (4,3), (4,4)}

So Dom (r)= {2,3,4}

& Range (r)= {4,3,2} or {2,3,4}



Q2. Repeat Q-1 when A = R the set of real number. Which of the real lines are functions?

Solution:

i. A = R

r ={(x,y) | y = x}


r ={(1,1), (2,2), (3,3),....} U {-1, -1}

{-2,-2),.....}

or

= {.. (- 2, 2), (-1, -1), ((0,0), (1,1),


(2,2), (3,3),.....}




ii. A x R

r = {(x+y) | (x+y = 5)}

r = { ....(-1, 6) (0,5), (1,5), (2,3), (3,2), (4,1) (5,0), (6,-1),...}



iii. {(x,y) | x+y=5}



iv. {(x,y) | x+y>5}



Q3. Which of the following diagrams represent functions and of which type?


Fig. (1)

Fig. (2)

Fig. (3)

Fig. (4)

Solution:

i. Fig (1), does not represent a function because element '1' has not unique image.

ii. Fig (2), represent a (1-1) one–to–one function. So , it is on to function.

iii. Fig. (3), represent a (1-1) one-to-one function, so, it is onto (injunction) function.

iv. Fig (4), represents a function which in onto function

Q4. Find the inverse of each of the following relations. Tell whether each relation and its inverse is a function or not.

i. {(2,1), (3,2), (4,3), (5,4), (6,5)}

ii. {(1,3), (2,5), (3,7), (4,9), (5,11)}

iii. {(x,y) | y = 2x+3 x ∈ R }

iv {(x,y) | y² = 4ax, x ≥ 0}

v. {(x,y) x²+y² = 9 , |x, y | x,y ≤ 3}

Solution:

i. r= {(2,1), (3,2), (4,3), (5,4), (6,5)}

r⁻¹={(2,1), (3,2), (4,3), (5,4), (6,5)}

dom (r⁻¹)= {1,2,3,4,5}

So, it is a function

ii. r= {(1,3), (2,5), (3,7), (4,9), (5,11)}

r⁻¹={(3,1), (2,5), (3,7), (4,9), (5,11)}

dom (r⁻¹)= {3,5,7,9,11}

So, it is a function

iii. r= {(x,y) | y = 2x+3, x ∈R}

r⁻¹={(x,y) | y = 2x+3, x ∈R}

It is a function

iv. r=((x,y) | y² = 4ax, x ≥ 0)

r⁻¹ ={(x,y) | x = $\frac{y^2}{4a}$, y ∈ R }

It is not a function

v. r= {(x,y) x²+y² = 9 , | x,y | ≤ |3|}

r⁻¹= {(x,y) x²+y² = 9 , |x, y | ≤ |3|}

It is not a function

Exercise 2.7

Q1. Complete the table, indicating by a tick mark those properties which are satisfied by the specified set of numbers.

property	Set of numbers	Natural	Whole	Integers	Rational	Reals
Closure	⊕					
Closure	⊗					
Associative	⊕					
Associative	⊗					
Identity	⊕					
Identity	⊗					
Inverse	⊕					
Inverse	⊗					
Commutative	⊕					
Commutative	⊗					

Solution:

Property	Set of numbers	Natural	Whole	Integers	Rational	Reals

Closure	⊕	✓	✓	✓	✓	✓
Closure	⊗	✓	✓	✓	✓	✓
Associative	⊕	✓	✓	✓	✓	✓
Associative	⊗	✓	✓	✓	✓	✓
Identity	⊕	X	✓	✓	✓	✓
Identity	⊗	✓	✓	✓	✓	✓
Inverse	⊕	x	x	✓	✓	✓
Inverse	⊗	x	x	x	x	x
Commutative	⊕	✓	✓	✓	✓	✓
Commutative	⊗	✓	✓	✓	✓	✓

Q2. What are the field axioms? In what respect does the field of real numbers differ from that of complex numbers?

Solution:

Field

A non-empty set 'F' is said to be field if for all x,y, $\exists$ $\in$ F, the following axiom are satisfied.

I. A field (F, +) is in abhain group i.e.

a. $x+y \in F;$ $\forall x, y \in F$

b. $(x+y)+z = x+(y+z)$ $\forall x,y,z \in F$

c. $x+0 = 0+x = x;$ $\forall x \in F$

d. $x+-x = -x+x = 0,$ $\forall x \in F$

e. $x+y = y+x;$ $\forall x, y \in F$

II. A field (F,) is an abehain group i.e

a. $x.y \in F;$ $\forall x, y \in F$

b. $x.(y.z) = (x.y).z;$ $\forall x,y,z \in F$

c. $x.1 = 1.x = x;$ $\forall x \in F$

d. $x.\frac{1}{x} = \frac{1}{x}.x = 1,$ $\forall x \in F$

e. $x.y = y.x;$ $\forall x, y \in F$

III. $\forall$ x, y, z ∈ F, the right distributive law holds i.e.

$x.(y+z) = xy+ xz.$

The field of real numbers differs. From the field of complex numbers in the since that an order relation can be defined in the field of real numbers, that is we can say that one real number is smaller than the others real number. So, they form two different fields.

Q3. Show that the adjoining table is that of multiplication of the elements of the set of residue classes modulo.5.

*	0	1	2	3	4
0	0	0	0	0	0
1	0	1	2	3	4
2	0	2	4	1	3
3	0	3	1	4	2
4	0	4	3	2	1

Solution:

As 0 ✱ 0 = 0

1 ✱ 0 = 0

2 ✱ 0 = 0

3 ✱ 0 = 0

4 ✱ 0 = 0

This show it is product table

2 ✱ 3 = 6 gives 1

2✱ 4 = 8 gives 2

and

3 ✱ 2 = 6 gives 1

3✱ 3 = 9 gives 4

3 ✱ 4 = 12 gives 3

4 ✱ 2 = 3 gives 3

4 ✱ 3 = 12 gives 2

4 ✱ 4 = 16 gives 1

This table shows that all elements of table is smaller than 5 So, the table is a multiplication able of the set of residue Classes modulo 5.

Q4. Prepare a table of addition of the elements of the set of residue classes module 4.

Solution:

The table of addition of the element of-the set of residue classes 'H' is given below elements 0,1,2,3.

⊕	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	0	1
3	3	0	1	2

5. Which of the following binary operations shown in tables (a) or (b) is commutative?

✱	a	b	c	d
a	a	b	c	d
b	b	c	b	a
c	c	d	b	c
d	a	a	b	ß

✱	a	b	c	d
a	a	c	b	d
b	c	d	b	a
c	b	b	a	c
d	d	a	c	d

Solution:

As the multiplication table is not symmetric that is when row and columns interchange the

table change.

In table (a)

a✱c = b, c✱a=c

Shows, a✱c ≠ c✱a

In table (b)

a✱c = b; c✱a

Shows, a✱c ≠ c✱a

Both, a and tables have no commutative property.

Q6. Supply the missing elements of the third row of the given table so that the operation ✱ may be associative.

✱	a	b	c	d
A	a	b	c	d
B	b	a	c	d
C	-	-	-	-
D	d	c	c	d

Solution:

Let x,y, z, t be the required element. If they are associative with binary operation ✱,

i.e

(i)✱(z) ✱a = c✱ (b✱a)....(1)

using the table, we have d ✱ b = c, b ✱ a = b, put these values in eq (i)

c ✱ a =z ✱ b;

➔ x = z, [∵a✱a = x, c✱b =z],

ii. (d ✱ b) ✱ b = d ✱ (b ✱ b)

c ✱ b =d ✱ a [∵ c✱ a = c, b ✱ b =b]

➔ y = d [∵ c ✱ b = y, d ✱ a=d]

iii. (d ✱ b) ✱ c = d ✱ (b ✱ c)

c ✱ c =d ✱ c = c [∵ d ✱ b = c , d ✱ c = c]

➔ Z = c [∵ d ✱ b = c , d ✱ c = c]

iv. (d ✱ b) ✱ d = d ✱ (b ✱ d)

c ✱ d =d ✱ d [∵ d ✱ b = c , d ✱ d = d]

➔ t = d [∵ c ✱ d = t , d ✱ d]

Q7. What operation is represented by the adjoining table? Name the identity element of the relevant set. If it exists. Is the operation associative? find the inverse of 0 1 2 3 if they exist.

✱	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	0	1
3	3	0	1	2

Solution:

0 ✱ 0 = 0

0 ✱ 1 =1

0 ✱ 2 = 2

0 ✱ 3 = 3

This shows that the operation is addition.

Identity element = 0

Because if does not change any number when it is added to it

0 ✱ 2 = 2 ✱ 0

Associative property

1 ✱ (2 ✱ 3) = (1 ✱ 2) ✱ 3.

1 ✱ 1 = 3 ✱ 3

2 = 2

Hence proved they have associative property

Inverse of

0 is 0

1 is 3

2 is 2

3 is 1

Exercise 2.8

Q1. Question performed on the two member set $Y = \{0, 1\}$ is shown in the adjoining table. Answer the questions.

i. What is the identity element if exists?

ii. What is the inverse of 1?

iii. Is the set S, under the given operation a group? Justify or Vidit Justify?

$*$	0	1
0	0	1
1	1	0

Solution

i. **Identity element = 0**

$1 * 0 = 1$

$0 * 1 = 1$

ii. **Inverse of $1^{-1} = 1$**

$1 * 1 = 1$

iii. **The set $S = \{0, 1\}$**

i. **Closure property**

$*$	0	1
0	0	1
1	1	0

Here all the elements are 0 and 1.

Hence, set S is closed.

ii. **Associative property**

$(1 * 1) * 1 = 1 * (1 * 1)$

$0 * 1 = 1 * 0$

$1 = 1$

iii. **Identity element exists**

$1 * 0 = 1$

$0 * 1 = 1$

iv. **Inverse for identity.**

$1^{-1} = 1$ (1 is its own inverse)

$0^{-1} = 0$ (0 is its own inverse)

v. **Commutative law**

$1 * 0 = 0 * 1$

$1 = 1$

commutative law holds

Hence, set S is abelian

Q2. The operation $*$ is performed on the set $S = \{1, 2, 3, 4\}$ shown in the adjoining table, show that the set is an abelian group?

$*$	1	2	3	4
1	1	2	3	4
2	2	1	4	3
3	3	4	1	2
4	4	3	2	1

$*$	1	2	3	4
1	1	2	3	4
2	2	1	4	3
3	3	4	1	2
4	4	3	2	1

Solution

i. **Associative property**

$(2 * 3) * 4 = 3 * (2 * 4)$

$1 * 4 = 3 * 3$

$4 = 4$

Associative is verified

ii. **Closure law**

$2 * 3 = 4$ (1 belongs to given table)

iii. **Inverse law**

$1 * 1 = 1$ and $2 * 2 = 2$

$3 * 4 = 4 * 3 = 1$

iv. **Identity exists**

$1 * 1 = 1$ and $2 * 1 = 2$

$3 * 1 = 3$ and $4 * 1 = 4$

v. **Commutative law**

OR

$1 * 2 = 2 * 1$

$2 = 2$

Yes, commutative law holds

Hence, the set is an abelian group

Q3. List each of the following sets. determine whether or not the set forms a group with respect to the indicated operation. Justify.

Set	Operation
i. The set of rational numbers	$\times$
ii. The set of rational numbers	$+$
iii. The set of positive rational numbers	$\times$
iv. The set of integers	$+$
v. The set of integers	$\times$

Solution

i. **The set of rational number under multiplication.**

Set of rational number = $\mathbb{Q}$

Let $a \in \mathbb{Q}$,

$a \neq 0$ then $a^{-1} \in \mathbb{Q}$

Therefore, the set of rational numbers is not a group w.r.t the binary operation multiplication.

ii. **The set of rational numbers under addition.**

iii. **Closure property**

$a + b = c \quad [\because a, b, c \in \mathbb{Q}]$

iv. **Associative property**

$(a + b) + c = a + (b + c) \quad [\because a, b, c \in \mathbb{Q}]$

v. **Identity element**

$a + 0 = a = 0 + a \quad [\because a, 0 \in \mathbb{Q}]$

vi. **Inverse element**

$a + (-a) = 0 = (-a) + a \quad [\because a \in \mathbb{Q}]$

vii. **Commutative property**

$a + b = b + a \quad [\because a, b \in \mathbb{Q}]$

Therefore, the set of rational numbers under addition forms a group.

iii. **The set of positive rational numbers under multiplication.**

Let $\mathbb{Q}$ denotes the set of positive rational numbers, then $\forall\, a \in \mathbb{Q}$

i. **Closure law**

$a \times b \in \mathbb{Q} \quad [\forall\, a, b \in \mathbb{Q}]$

Product of two rational numbers is also a positive rational number.

ii. **Associative law**

$(a \times b) \times c = a \times (b \times c) \quad [\forall\, a, b, c \in \mathbb{Q}]$

Product of these rational numbers is also a positive rational number so, Associative law holds

iii. **Identity element**

$1 \in \mathbb{Q} \quad [\forall\, a \in \mathbb{Q}]$

$a \times 1 = 1 \times a = a$

Hence, '1' is the identity element of $\mathbb{Q}$.

iv. **Inverse element**

$a \times \frac{1}{a} = \frac{1}{a} \times a = 1 \quad [\forall\, a \in \mathbb{Q}]$

Hence, $\frac{1}{a}$ is positive rational number & group under multiplication.

v. **Commutative law**

$a \times b = b \times a \quad [\forall\, a, b \in \mathbb{Q}]$

It is true that group $(\mathbb{Q}, \times)$ is abelian group.

vi. **The set of integers under addition.**

Let $\mathbb{I}$ be the set of integers then

$\mathbb{I} = \{\ldots -3, -2, -1, 0, 1, 2, 3, \ldots\}$

i. **Closure law**

$a + b \in \mathbb{I} \quad [\forall\, a, b \in \mathbb{I}]$

Sum of two integers is also an integer.

ii. **Associative law**

$(a + b) + c = a + (b + c) \quad [\forall\, a, b, c \in \mathbb{I}]$

iii. **Identity element**

$a + 0 = a = 0 + a \quad [\forall\, a \in \mathbb{I}, 0 \in \mathbb{I}]$

Hence, 0 is identity element of $\mathbb{I}$.

iv. **Inverse element**

$a + (-a) = 0 = (-a) + a \quad [\forall\, a \in \mathbb{I}]$

Hence integer under addition is a case.

v. **Commutative property**

$a + b = b + a \quad [\forall\, a, b \in \mathbb{I}]$

Hence, the group is Abelian group.

vii. **The set of integers under multiplication.**

Let $\mathbb{I}$ be set of integers then

$\mathbb{I} = \{\ldots -3, -2, -1, 0, 1, 2, 3, \ldots\}$

i. **Closure law**

$a \times b \in \mathbb{I} \quad [\forall\, a, b \in \mathbb{I}]$

ii. **Associative law**

$(a \times b) \times c = a \times (b \times c) \quad [\forall\, a, b, c \in \mathbb{I}]$

iii. **Identity law**

$a \times 1 = 1 \times a = a \quad [\forall\, a, b, c \in \mathbb{I}]$

iv. **Inverse element**

$a \times \frac{1}{a} = \frac{1}{a} \times a = 1 \quad [\forall\, a \in \mathbb{I}]$

Hence integer under addition is a case.

v. **Commutative property**

$a \times b = b \times a \quad [\forall\, a, b \in \mathbb{I}]$

Hence, the group is Abelian group.

v. The set of integers under multiplication.

Let $\mathbb{I}$ be set of integers then

$\mathbb{I} = \{\ldots -3, -2, -1, 0, 1, 2, 3, \ldots\}$

i. **Closure law**

$a \times b \in \mathbb{I} \quad [\forall\, a, b \in \mathbb{I}]$

ii. **Associative law**

$(a \times b) \times c = a \times (b \times c) \quad [\forall\, a, b, c \in \mathbb{I}]$

iii. **Identity law**

$a \times 1 = 1 \times a = a \quad [\forall\, a, b, c \in \mathbb{I}]$

iv. **Inverse element**

$a \times \frac{1}{a} = \frac{1}{a} \times a = 1 \quad [\forall\, a \in \mathbb{I}]$

Hence integer under addition is a case.

v. **Commutative property**

$a \times b = b \times a \quad [\forall\, a, b \in \mathbb{I}]$

Hence, the group is Abelian group.

vi. The set of integers under multiplication.

Let $\mathbb{I}$ be set of integers then

$\mathbb{I} = \{\ldots -3, -2, -1, 0, 1, 2, 3, \ldots\}$

i. **Closure law**

$a \times b \in \mathbb{I} \quad [\forall\, a, b \in \mathbb{I}]$

ii. **Associative law**

$(a \times b) \times c = a \times (b \times c) \quad [\forall\, a, b, c \in \mathbb{I}]$

iii. **Identity law**

$a \times 1 = 1 \times a = a \quad [\forall\, a, b, c \in \mathbb{I}]$

iv. **Inverse element**

$a \times \frac{1}{a} = \frac{1}{a} \times a = 1 \quad [\forall\, a \in \mathbb{I}]$

Hence integer under addition is a case.

v. **Commutative property**

$a \times b = b \times a \quad [\forall\, a, b \in \mathbb{I}]$

Hence, the group is Abelian group.

vii. The set of integers under addition.

Let $\mathbb{I}$ be set of integers then

$\mathbb{I} = \{\ldots -3, -2, -1, 0, 1, 2, 3, \ldots\}$

i. **Closure law**

$a + b \in \mathbb{I} \quad [\forall\, a, b \in \mathbb{I}]$

Sum of two integers is also an integer.

ii. **Associative law**

$(a + b) + c = a + (b + c) \quad [\forall\, a, b, c \in \mathbb{I}]$

iii. **Identity element**

$a + 0 = a = 0 + a \quad [\forall\, a \in \mathbb{I}, 0 \in \mathbb{I}]$

Hence, 0 is identity element of $\mathbb{I}$.

iv. **Inverse element**

$a + (-a) = 0 = (-a) + a \quad [\forall\, a \in \mathbb{I}]$

Hence integer under addition is a case.

v. **Commutative property**

$a + b = b + a \quad [\forall\, a, b \in \mathbb{I}]$

Hence, the group is Abelian group.

viii. The set of integers under multiplication.

Let $\mathbb{I}$ be set of integers then

$\mathbb{I} = \{\ldots -3, -2, -1, 0, 1, 2, 3, \ldots\}$

i. **Closure law**

$a \times b \in \mathbb{I} \quad [\forall\, a, b \in \mathbb{I}]$

ii. **Associative law**

$(a \times b) \times c = a \times (b \times c) \quad [\forall\, a, b, c \in \mathbb{I}]$

iii. **Identity law**

$a \times 1 = 1 \times a = a \quad [\forall\, a, b, c \in \mathbb{I}]$

iv. **Inverse element**

$a \times \frac{1}{a} = \frac{1}{a} \times a = 1 \quad [\forall\, a \in \mathbb{I}]$

Hence integer under addition is a case.

v. **Commutative property**

$a \times b = b \times a \quad [\forall\, a, b \in \mathbb{I}]$

Hence, the group is Abelian group.

ix. The set of integers under subtraction.

Let $\mathbb{I}$ be set of integers then

$\mathbb{I} = \{\ldots -3, -2, -1, 0, 1, 2, 3, \ldots\}$

i. **Closure law**

$a - b \in \mathbb{I} \quad [\forall\, a, b \in \mathbb{I}]$

Sum of two integers is also an integer.

ii. **Associative law**

$(a - b) - c = a - (b - c) \quad [\forall\, a, b, c \in \mathbb{I}]$

iii. **Identity element**

$a - 0 = a = 0 - a \quad [\forall\, a \in \mathbb{I}, 0 \in \mathbb{I}]$

Hence, 0 is identity element.

iv. **Inverse element**

$a - (-a) = 0 = (-a) - a \quad [\forall\, a \in \mathbb{I}]$

Hence integer under addition is a case.

v. **Commutative property**

$a - b = b - a \quad [\forall\, a, b \in \mathbb{I}]$

Hence, the group is Abelian group.

x. The set of integers under division.

Let $\mathbb{I}$ be set of integers then

$\mathbb{I} = \{\ldots -3, -2, -1, 0, 1, 2, 3, \ldots\}$

i. **Closure law**

$a \div b \in \mathbb{I} \quad [\forall\, a, b \in \mathbb{I}]$

Sum of two integers is also an integer.

ii. **Associative law**

$(a \div b) \div c = a \div (b \div c) \quad [\forall\, a, b, c \in \mathbb{I}]$

iii. **Identity element**

$a \div 0 = a = 0 \div a \quad [\forall\, a \in \mathbb{I}, 0 \in \mathbb{I}]$

Hence, 0 is identity element.

iv. **Inverse element**

$a \div (-a) = 0 = (-a) \div a \quad [\forall\, a \in \mathbb{I}]$

Hence integer under addition is a case.

v. **Commutative property**

$a \div b = b \div a \quad [\forall\, a, b \in \mathbb{I}]$

Hence, the group is Abelian group.

Q4. Let S be the set of all non-singular matrices over the real field from a non-abelian group under multiplication.

Let '1' denotes the set of all non-singular 2 x multiple over the real field.

Then

i. **Closure law**

$\forall\, a, b \in S \Rightarrow a \times b \in S$

ii. **Associative law**

$\forall\, a, b, c \in S, (a \times b) \times c = a \times (b \times c)$

iii. **Identity Element**

$\exists\, e \in S$ such that $a \times e = e \times a = a$

Here, I_2 is a unit matrix. therefore, a unit group as well.

$\forall\, a \in S, e = I_2$

iv. **Inverse law**

$\forall\, a \in S, a^{-1} \in S$ such that $a \times a^{-1} = a^{-1} \times a = I_2$

v. **Commutative law**

$\forall\, a, b \in S, a \times b \neq b \times a$

So, the S the set of all 2 x 2 non-singular matrices over the real field is a non-abelian group under multiplication.