CONTENTS

Unit No.	Units	Page No.
01	Real Numbers	01
02	Logarithms	30
03	Set and Functions	54
04	Factorization and Algebraic Manipulation	93
05	Linear Equations and Inequalities	117
06	Trigonometry	153
07	Coordinate Geometry	201
08	Logic	237
09	Similar Figures	252
10	Graphs of Functions	288
11	Loci and Construction	316
12	Information Handling	345
13	Probability	382

www.ilmkidunya.com


UNIT 01

Real Numbers


Introduction to Real Numbers

The history of numbers comprises thousands of years, from ancient civilization to the modern Arabic system.

Sumerians: (4500–1900BCE) used a sexagesimal (base 60) system for counting. The Sumerians used a small cone, bead, large cone, large perforated cone, sphere and perforated sphere, corresponding to 1, 10, 60 (a large unit), 600.



Egyptians: (3000–2000 BCE) used a decimal (base 10) system for counting. Here are some of the symbols used by the Egyptians, as shown in the figure below: The Egyptians usually wrote numbers left to right, starting with the highest denominator. For example, 2525 would be written with 2000 first, then 500, 20, and 5.



Romans: (500BCE-500CE) used the Roman materials system for counting. Roman numerals represent a number system that was widely used throughout Europe as the standard writing system until the late Middle ages. The ancient Romans explained that when a number reaches 10 it is not easy to count on one's fingers. Therefore, there was a need to create a proper number system that could be used for trade and communications. Roman numerals use 7 letters to represent different numbers. These are I, V X, L,C,D and M which represent the numbers 1,5,10,50,100,500 and 1000 respectively.

Indians: (500-1200)CE developed the concept of zero (0) and made a significant contribution to the decimal (base 10) system.

Ancient Indian mathematicians have contributed immensely to the field of mathematics. The invention of zero is attributed to Indians, and this contribution outweighs all others made by any other nation since it is the basis of the decimal number system, without which no advancement in mathematics would have been possible. The number system used today was invented by Indians, and it is still called Indo-Arabic numerals because Indians invented them and the Arab merchants took them to the Western world.



Arabs: (800-1500CE) introduced Arabic numerals (0-9) to Europe. The Islamic world underwent significant developments in mathematics. Muhammad bin Musa al-



www.ilmkidunya.com

Khwarzimi played a key role in this transformation, introducing algebra as a distinct field in the 9th century. Al-Khwarzimi's approach departing from earlier arithmetical traditions, laid the groundwork for a arithmetization of algebra, influencing mathematical thought for an extended period. Successors like Al-Karaji expanded on his work, contributing to advancements in various mathematical domains. The practicality and broad applicability of these mathematical methods facilitated the dissemination of Arabic mathematics to the West, contributing substantially to the evolution of Western mathematics.

Modern era (1700-present): Developed modern number systems e.g., binary system base -2) and hexadecimal system (base-16).


The Arabic system is the basis for modern decimal system used globally today. Its development and refinement comprise thousands of years from ancient Sumerians to modern mathematicians.

Combination of Rational and

In the modern era, the set [1,2,3,...] was adopted as the counting set. This counting set represents the set of natural numbers was extended to set of real numbers which is used most frequently in everyday life.

Real Numbers:

The set of Real numbers is the union (combination) of the set of rational numbers and irrational numbers i,e., $R = Q \cup Q'$



Rational Numbers:

The set of rational numbers is defined as the set of numbers that contains those elements which can be expressed as quotient of two integers. For example, $\frac{2}{3}, \frac{4}{5}, \frac{6}{7}$....*etc.*

$$Q = \left\{ \frac{p}{q}; p, q \in Z \wedge q \neq 0 \right\}$$

Irrational Numbers:

The set of irrational numbers Q' contains those elements which can not be expressed as quotient of integers.

$$Q' = \left\{ x \neq \frac{p}{q}; p, q \in Z \wedge q \neq 0 \right\}$$

For example, $\pi, e, \sqrt{2}, \sqrt{3}, \sqrt{5}$ and $\sqrt{7}$ *etc.* are irrational numbers.

Decimal Representation of Rational Numbers

(i) Terminating Decimal Numbers:

A decimal number with a finite number of digits after the decimal point is called a terminating decimal number.

For example $\frac{1}{4} = 0.25, \frac{8}{25} = 0.32, \frac{3}{8} = 0.375$,

$\frac{4}{5} = 0.8$

are all terminating decimals.

(ii) Recurring and Non-Terminating Decimal Numbers

The decimal numbers with a repeating pattern of digits after the decimal point are called recurring decimal numbers.

For example,

www.ilmkidunya.com

$\frac{1}{3} = 0.333... = 0.\overline{3}$ (the 3 repeats infinitely)

$\frac{1}{6} = 0.1666... = 0.1\overline{6}$ (the 6 repeats infinitely)

Example 1: Identify the following decimal numbers as rational or irrational numbers:

(i) 0.35 09301001 (ii)0.444... 09301002

(iii) $3.\overline{5}$ 09301003

(iv) 3.36788542... 09301004

(v) 1.709975947... 09301005

Solution:

(i) 0.35 is a terminating decimal number, therefore it is a rational number.

(ii) 0.444... is a recurring decimal number, therefore it is a rational number.

(iii) $3.\overline{5} = 3.5555...$ is a recurring decimal number, therefore it is a rational number.

(iv) 3.3678542... is a non-terminating and non-recurring decimal number. Therefore, it represents an irrational number.

(v) 1.709975947... is a non-terminating and non-recurring decimal number, it is an irrational number.


Representation of Real Numbers on number line
YouTube
Online Lecture
publications.unique.edu.pk UGI.publications uniquenotesofficial @Uniquepublications 0324-6666661-2-3

Representation of rational and irrational numbers on number line

Example 2: Represent $\sqrt{5}$ on a number line. 09301006

Solution:

$\sqrt{5}$ can be located on the real line by geometric construction. As $\sqrt{5} = 2.236...$ which is near to 2. Mark a perpendicular line of $m\overline{AB} = 1$ unit at A, where $m\overline{OA} = 2$ units, and we have a right-angle triangle OAB. By using Pythagoras theorem



$$\left(m\overline{OB}\right)^2 = \left(m\overline{OA}\right)^2 + \left(m\overline{AB}\right)^2$$

$$\Rightarrow m\overline{OB} = \sqrt{(2)^2 + (1)^2} = \sqrt{4+1} = \sqrt{5}$$

$$\Rightarrow m\overline{OB} = \sqrt{5}$$

Draw an arc of radius $m\overline{OB} = \sqrt{5}$ taking O as centre, we got point "P" representing $\sqrt{5}$ on the number line

So, $\left|\overline{OP}\right| = \sqrt{5}$

Example 3: Express the following recurring decimals as the rational number $\frac{p}{q}$, where p and q are integers.

(i) $0.\overline{5}$ 09301007 (ii) $0.\overline{93}$ 09301008

www.ilmkidunya.com

(i) $0.\overline{5}$

Solution:

$0.\overline{5}$

Let $x = 0.\overline{5}$

$x = 0.5555\ldots$ _____(i)

Multiply both sides by 10

$10x = 10(0.5555\ldots)$

$10x = 5.555\ldots$ _____(ii)

Subtracting eq. (i) from eq. (ii)

$10x - x = (5.555\ldots) - (0.5555\ldots)$

$9x = 5$

$\Rightarrow \quad x = \frac{5}{9}$

Which shows the rational number in the form $\frac{p}{q}$.

(ii) $0.\overline{93}$

Solution

$0.\overline{93}$

$x = 0.93939393\ldots$ _____(i)

Multiply both sides by 100, we get

$100x = 100\,(0.93939393\ldots)$

$100x = 93.939393\ldots$ ...(ii)

Subtracting (i) from (ii)

$100x - x = (93.939393\ldots) - (0.93939393\ldots)$

$99x = 93$

$x = \frac{93}{99}$ which is a rational number in the form of $\frac{p}{q}$.

Example 4: Insert two rational numbers between 2 and 3. 09301009

Solution

There are infinite rational numbers between 2 and 3.

We find any two of them

For this, find the average of 2 and 3

$$= \frac{2+3}{2} = \frac{5}{2}$$

So, $\frac{5}{2}$ is a rational number between 2 and 3, to find another rational number between 2 and 3 we will again find average of $\frac{5}{2}$ and 3 i.e,

$$= \left(\frac{5}{2} + 3\right) \div 2$$

$$= \left(\frac{5+6}{2}\right) \div \frac{2}{1}$$

$$= \left(\frac{11}{2}\right) \times \frac{1}{2} = \frac{11}{4}$$

Hence two rational numbers between 2 and 3 are $\frac{5}{2}$ and $\frac{11}{4}$.

Try Yourself

What will be the product of two irrational numbers?

The product of two irrational numbers may or may not be an irrational number.

In most cases, the product of two irrational numbers will be irrational numbers but not always.

Two irrational numbers may multiply to form a rational number.

For example,

$\sqrt{2} \times \sqrt{8} = \sqrt{16} = 4$ (rational) and

$\sqrt{2} \times \sqrt{3} = \sqrt{6}$ (irrational)

Properties of Real Numbers

All calculations involving addition, subtraction, multiplication, and division of real numbers are based on their properties. In this section, we shall discuss these properties.

www.ilmkidunya.com

Additive properties

Name of the property	$\forall a, b, c \in R$	Examples
Closure	$a+b \in R$	$2+3=5 \in R$
Commutative	$a+b=b+a$	$2+5=5+2$ $7=7$
Associative	$a+(b+c)=(a+b)+c$	$2+(3+5)=(2+3)+5$ $2+8=5+5$ $10=10$
Identity	$a+0=a=0+a$	$5+0=5=0+5$
Inverse	$a+(-a)=-a+a=0$	$6+(-6)=(-6)+6=0$

Name of the property	$\forall a, b, c \in R$	Examples
Closure	$ab \in R$	$2 \times 5 = 10 \in R$
Commutative	$ab=ba$	$2 \times 3 = 3 \times 2 = 6$
Associative	$a(bc)=(ab)c$	$2 \times (3 \times 5) = (2 \times 3) \times 5$ $2 \times 15 = 6 \times 5$ $30 = 30$
Identity	$a \times 1 = 1 \times a = a$	$5 \times 1 = 1 \times 5 = 5$
Inverse	$a \times \frac{1}{a} = \frac{1}{a} \times a = 1$	$7 \times \frac{1}{7} = \frac{1}{7} \times 7 = 1$

Do you know?

0 and 1 are the additive and multiplicative identities of real numbers respectively.

Remember!

$0 \in R$ has no multiplicative Inverse.

Distributive Properties

For all real numbers a, b, c

(i) $a(b+c)=ab+ac$ is called left distributive property of multiplication over addition.

(ii) a(b–c) = ab–ac is called left distributive property of multiplication over subtraction.

(iii) $(a+b)c=ac+bc$ is called right distributive property of multiplication over addition.

(iv) (a–b)c = ac–bc is called right distributive property of multiplication over subtraction.

Properties of Equality of Real number

i.	Reflexive property	$\forall a \in R,\ a=a$
ii.	Symmetric property	$\forall a,b \in R,\ a=b \Rightarrow b=a$
iii.	Transitive property	$\forall a,b,c \in R,\ a=b \wedge b=c \Rightarrow a=c$
iv	Additive properly.	$\forall a,b,c \in R,\ a=b \Rightarrow a+c=b+c$
v.	Multiplicative property	$\forall a,b,c \in R,\ a=b \Rightarrow ac=bc$
vi.	Cancellation property w.r.t addition	$\forall a,b,c \in R,\ a+c=b+c \Rightarrow a=b$
vii.	Cancellation property w.r.t multiplication	$\forall a,b,c \in R$, and $c \neq 0$ $ac=bc \Rightarrow a=b$

www.ilmkidunya.com

Order Properties

i.	Trichotomy property	$\forall a,b \in R$, either $a=b$ or $a>b$ or $a<b$
ii.	Transitive Property	$\forall a,b,c \in R$ • $a>b \wedge b>c \Rightarrow a>c$ • $a<b \wedge b<c \Rightarrow a<c$
iii.	Additive property	$\forall a,b,c \in R$ • $a>b \Rightarrow a+c>b+c$ • $a<b \Rightarrow a+c<b+c$
iv	Multiplicative property	$\forall a,b,c \in R$ • $a>b \Rightarrow ac>bc$ if $c>0$ • $a<b \Rightarrow ac<bc$ if $c>0$ • $a>b \Rightarrow ac<bc$ if $c<0$ • $a<b \Rightarrow ac>bc$ if $c<0$ • $a>b \wedge c>d \Rightarrow ac>bd$ • $a<b \wedge c<d \Rightarrow ac<bd$
v	Division property	• $\forall a,b,c \in R$ • $a<b \Rightarrow \frac{a}{c}<\frac{b}{c}$ if $c>0$ • $a<b \Rightarrow \frac{a}{c}>\frac{b}{c}$ if $c<0$ • $a>b \Rightarrow \frac{a}{c}>\frac{b}{c}$ if $c>0$ • $a>b \Rightarrow \frac{a}{c}<\frac{b}{c}$ if $c<0$
vi	Reciprocal property	$\forall a, b \in R$ and have same sign • $a<b \Rightarrow \frac{1}{a}>\frac{1}{b}$ • $a>b \Rightarrow \frac{1}{a}<\frac{1}{b}$

Example 5: If $a=\frac{2}{3}, b=\frac{3}{2}, c=\frac{5}{3}$ then verify the distributive property over addition. 09301010

Solution: (i) Left distributive property

$a(b+c)=ab+ac$

L.H.S = $a(b+c)$	R.H.S = $ab+ac$
$=\frac{2}{3}\left(\frac{3}{2}+\frac{5}{3}\right)$ $=\frac{2}{3}\left(\frac{9+10}{6}\right)$	$=\left(\frac{2}{3}\right)\left(\frac{3}{2}\right)+\left(\frac{2}{3}\right)\left(\frac{5}{3}\right)$ $=1+\frac{10}{9}$

www.ilmkidunya.com

$=\frac{2}{3}\left(\frac{19}{6}\right)$	$=\frac{9+10}{9}$
$=\frac{19}{9}$	$=\frac{19}{9}$

L.H.S = R.H.S
Hence proved.

(ii) Right distributive property

$(a+b)c = ac+bc$

L.H.S = $(a+b)c$	R.H.S = $ac+bc$
$=\left(\frac{2}{3}+\frac{3}{2}\right)\frac{5}{3}$	$=\left(\frac{2}{3}\right)\left(\frac{5}{3}\right)+\left(\frac{3}{2}\right)\left(\frac{5}{3}\right)$
$=\left(\frac{4+9}{6}\right)\frac{5}{3}$	$=\frac{10}{9}+\frac{15}{6}$
$=\left(\frac{13}{6}\right)\left(\frac{5}{3}\right)$	$=\frac{20+45}{18}$
$=\frac{65}{18}$	$=\frac{65}{18}$

L.H.S = R.H.S
Hence Proved

Example 6: Identify the property that justifies the statement

(i) If $a > 13$ then $a+2 > 15$ 09301011
(ii) If $3 < 9$ and $6 < 12$ then $9 < 21$ 09301012
(iii) If $7 > 4$ and $5 > 3$ then $35 > 12$ 09301013
(iv) If $-5 < -4 \Rightarrow 20 > 16$ 09301014

Solution

(i) $a > 13$
Add 2 on both sides
$a+2 > 13+2$
$a+2 > 15$
(order property w.r.t addition)

(ii) As $3 < 9$ and $6 < 12$
$\Rightarrow 3+6 < 9+12$
$9 < 21$
(order property w.r.t addition)

(iii) $7 > 4$ and $5 > 3$
$\Rightarrow 7 \times 5 > 4 \times 3$
$\Rightarrow 35 > 12$
(order property w.r.t multiplication)

(iv) As $-5 < -4$
Mulyplying both sides by -4
$-5 \times -4 > -4 \times -4$
$\Rightarrow 20 > 16$
(order property w.r.t multiplication)

Exercise 1.1

Q.1 Identify each of the following as a rational or irrational numbers:

Solution:

Rational numbers

(i) 2.353535 09301015 (ii) $0.\overline{6}$ 09301016

(ix) $\frac{15}{4}$ 09301017 (x) $\left(2-\sqrt{2}\right)(2+\sqrt{2})$ 09301018

Irrational numbers

(iii) 2.236067... 09301019 (iv) $\sqrt{7}$ 09301020

(v) e 09301021 (vi) π 09301022

(vii) $5+\sqrt{11}$ 09301023 (viii) $\sqrt{3}+\sqrt{13}$ 09301024

Q.2 Represent the following numbers on number line:

(i) $\sqrt{2}$ 09301025

Solution:


$\sqrt{2}$ can be located on the real line by geometric construction. Mark a perpendicular line of $m\overline{AB} = 1$ unit at A, where $m\overline{OA} = 1$ unit, and we have a right-angle triangle OAB. By using Pythagoras theorem

$$\left(m\overline{OB}\right)^2 = \left(m\overline{OA}\right)^2 + \left(m\overline{AB}\right)^2$$

$$\sqrt{\left(m\overline{OB}\right)^2} = \sqrt{\left(m\overline{OA}\right)^2 + \left(m\overline{AB}\right)^2}$$

$$m\overline{OB} = \sqrt{(1)^2+(1)^2} = \sqrt{1+1} = \sqrt{2}$$

www.ilmkidunya.com



Taking O as centre, draw an arc of radius $m\,\overline{OB} = \sqrt{2}$ Which cut the number line at P. W e get point "P" representing $\sqrt{2}$ on the number line

So, $\left|\overline{OP}\right| = \sqrt{2}$

(ii) $\sqrt{3}$ 09301026

Solution:

$\sqrt{3}$ can be located on the real line by geometric method. Mark a line of $m\overline{AB} = 1$ unit at A, With centre at A draw an arc of radius 2 units above the line. From point B draw a perpendicular line segment so that it cuts the arc at C. Join A to C. We have a right-angled ΔABC *in which* $m\overline{AB} = 1$ unit and $m\overline{AC} = 2$ units. By using Pythagoras theorem.

$$\left(\text{m}\overline{\text{AC}}\right)^2 = \left(\text{m}\overline{\text{AB}}\right)^2 + \left(\text{m}\overline{\text{BC}}\right)^2$$

$$(2)^2 = (1)^2 + \left(\text{m}\overline{\text{BC}}\right)^2$$


$$4 = 1 + \left(\text{m}\overline{\text{BC}}\right)^2$$

$$4 - 1 = \left(\text{m}\overline{\text{BC}}\right)^2$$

$$3 = \left(\text{m}\overline{\text{BC}}\right)^2$$

$$\sqrt{\left(\text{m}\overline{\text{BC}}\right)^2} = \sqrt{3}$$

$$\text{m}\overline{\text{BC}} = \sqrt{3}$$




Now consider B is at 0.Taking B as centre, draw an arc of radius $m\,\overline{BC} = \sqrt{3}$, which cut the number line at P. We get point "P" representing $\sqrt{3}$ on the number line

So, $\left|\overline{BP}\right| = \sqrt{3}$

(iii) $4\frac{1}{3}$ 09301027


Solution:



Point P represents $4\frac{1}{3}$ on the number line.

(iv) $-2\frac{1}{7}$ 09301028


Solution:



Point P represents $-2\frac{1}{7}$ on the number line.

(v) $\frac{5}{8}$ 09301029

Solution:




www.ilmkidunya.com

Point P represents $\frac{5}{8}$ on the number line.

(vi) $2\frac{3}{4}$ 09301030

Solution:



Point P represents $2\frac{3}{4}$ on the number line.

Q.3 Express the following as a rational number $\frac{p}{q}$ where p and q are integers and $q \neq 0$.

(i) $0.\overline{4}$ 09301031

Solution:

Let $x = 0.\overline{4}$

$x = 0.4444\ldots$ _____ (i)

Multiplying both sides by "10", we get

$10x = 10\ (0.4444\ldots)$

$10x = 4.444\ldots$ _____ (ii)

Subtracting eq.(i) from (ii)

$10x - x = (4.444\ldots) - (0.4444\ldots)$

$9x = 4$

$x = \frac{4}{9}$

$\Rightarrow 0.\overline{4} = \frac{4}{9}$

(ii) $0.\overline{37}$ 09301032

Solution:

Let $x = 0.\overline{37}$

$x = 0.37373737\ldots$ _____ (i)

Multiplying both sides by "100"

$100x = 100\ (0.37373737\ldots)$

$100x = 37.373737\ldots$ _____ (ii)

Subtracting eq.(i) from (ii)

$100x - x = (37.373737\ldots) - (0.37373737\ldots)$

$99x = 37$

$x = \frac{37}{99}$

$\Rightarrow 0.\overline{37} = \frac{37}{99}$

(iii) $0.\overline{21}$ 09301033

Solution:

Let $x = 0.\overline{21}$

$x = 0.21212121\ldots$ _____ (i)

Multiplying both sides by "100"

$100x = 100\ (0.21212121\ldots)$

$100x = 21.212121\ldots$ _____ (ii)

Subtracting eq.(i) from (ii)

$100x - x = (21.212121\ldots) - (0..21212121\ldots)$

$99x = 21$

$x = \frac{21}{99} = \frac{7}{33}$

$\Rightarrow 0.\overline{21} = \frac{7}{33}$

Q.4 Name the property used in the following. 09301034

Solution:

Sr. No.		Property Name
(i)	$(a + 4) + b = a + (4+b)$	Associative property w.r.t addition
(ii)	$\sqrt{2} + \sqrt{3} = \sqrt{3} + \sqrt{2}$	Commutative property w.r.t addition
(iii)	$x - x = 0$	Additive Inverse
(iv)	$a(b+c) = a\,b + a\,c$	Left distributive property of multiplication over addition.
(v)	$16 + 0 = 16$	Additive Identity
(vi)	$100 \times 1 = 100$	Multiplicative identity
(vii)	$4 \times (5 \times 8) = (4 \times 5) \times 8$	Associative property w.r.t multiplication
(viii)	$ab = ba$	Commutative property w.r.t multiplication.

www.ilmkidunya.com

5. Name the property used in the following:

Solution:

(i) $-3 < -1 \Rightarrow 0 < 2$ 09301035

Additive property of inequality

(ii) If $a < b$ **then** $\frac{1}{a} > \frac{1}{b}$ 09301036

Reciprocal property

(iii) If $a < b$ **then** $a+c < b+c$ 09301037

Additive property of inequality

(iv) If $ac < bc$ **and** $c > 0$ **then** $a < b$

Cancellation property of inequality w.r.t multiplication.

(v) If $ac < bc$ **and** $c < 0$ **then** $a > b$ 09301038

Cancellation property of inequality w.r.t multiplication.

(vi) Either $a > b$ **or** $a = b$ **or** $a < b$

Trichotomy property

6. Insert two rational numbers between

(i) $\frac{1}{3}$ **and** $\frac{1}{4}$ 09301039

Solution:

Two rational numbers between $\frac{1}{3}$ and $\frac{1}{4}$

Average of $\frac{1}{3}$ and $\frac{1}{4} = \left(\frac{1}{3}+\frac{1}{4}\right)\div 2$

$$= \left[\frac{4+3}{12}\right]\times\frac{1}{2}$$

$$= \frac{7}{12}\times\frac{1}{2} = \frac{7}{24}$$

Now we find average of $\frac{1}{3}$ and $\frac{7}{24}$

Average of $\frac{1}{3}$ and $\frac{7}{24} = \left(\frac{1}{3}+\frac{7}{24}\right)\div 2$

$$= \frac{8+7}{24}\times\frac{1}{2}$$

$$= \frac{15}{24}\times\frac{1}{2} = \frac{15}{48} = \frac{5}{16}$$

Thus $\frac{5}{16}$ and $\frac{7}{24}$ are two rational numbers between $\frac{1}{3}$ and $\frac{1}{4}$.

(ii) 3 and 4 09301040

Solution:

Two rational numbers between 3 and 4.

Average of 3 and $4 = \frac{3+4}{2} = \frac{7}{2}$

Average of $\frac{7}{2}$ and $4 = \left(\frac{7}{2}+4\right)\div 2$

$$= \left(\frac{7+8}{2}\right)\div\frac{2}{1}$$

$$= \frac{15}{2}\times\frac{1}{2}$$

$$= \frac{15}{4}$$

Thus $\frac{7}{2}$ and $\frac{15}{4}$ are two rational numbers between 3 and 4.

(iii) $\frac{3}{5}$ **and** $\frac{4}{5}$ 09301041

Solution:

Two rational numbers between $\frac{3}{5}$ and $\frac{4}{5}$

Average of $\frac{3}{5}$ and $\frac{4}{5} = \left(\frac{3}{5}+\frac{4}{5}\right)\div 2$

$$= \left(\frac{3+4}{5}\right)\times\frac{1}{2}$$

$$= \frac{7}{5}\times\frac{1}{2} = \frac{7}{10}$$

Average of $\frac{7}{10}$ and $\frac{4}{5} = \left(\frac{7}{10}+\frac{4}{5}\right)\div 2$

$$= \left(\frac{7+8}{10}\right)\times\frac{1}{2}$$

$$= \frac{15}{10}\times\frac{1}{2}$$

$$= \frac{15}{20} = \frac{3}{4}$$

Thus $\frac{7}{10}$ and $\frac{3}{4}$ are two rational numbers between $\frac{3}{5}$ and $\frac{4}{5}$.

www.ilmkidunya.com

Radical Expressions

If n is a positive integer greater than 1, and a is a real number, then any real number x such that $x = \sqrt[n]{a}$ is called n^{th} root of a.

Here $\sqrt{\ }$ is called radical, and n is the index of radical. A real number under the radical sign is called a radicand. $\sqrt[3]{5}, \sqrt[5]{7}$ are examples of radical form.

Laws of Radicals and Indices

Laws of Radical	Laws of Indices
(i) $\sqrt[n]{ab} = \sqrt[n]{a}.\sqrt[n]{b}$	(i) $a^m.a^n = a^{m+n}$
(ii) $\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}$	(ii) $(a^m)^n = a^{mn}$
(iii) $\sqrt[n]{a^m} = (\sqrt[n]{a})^m$	(iii) $(ab)^n = a^n b^n$
(iv) $\left(\sqrt[n]{a}\right)^n = (a^{\frac{1}{n}})^n = a$	(iv) $\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}$
	(v) $\frac{a^m}{a^n} = a^{m-n}$
	(vi) $a^0 = 1$

Example 7

Simplify the following:

(i) $\sqrt[4]{16x^4y^8}$ 09301042

Solution:

$\sqrt[4]{16x^4y^8}$

$= (16x^4y^8)^{\frac{1}{4}}$ $\quad \sqrt[n]{a} = a^{\frac{1}{n}}$

$= (16)^{\frac{1}{4}}(x^4)^{\frac{1}{4}}(y^8)^{\frac{1}{4}}$ $\quad \because (ab)^m = a^m b^m$

$= 2^{4\cdot\frac{1}{4}} \times x^{4\cdot\frac{1}{4}} \times y^{8\cdot\frac{1}{4}}$ $\quad \because (a^m)^n = a^{mn}$

$= 2xy^2$

(ii) $\sqrt[3]{27x^6y^9z^3}$ 09301043

Solution:

$\sqrt[3]{27x^6y^9z^3}$

$= \left(27x^6y^9z^3\right)^{\frac{1}{3}}$ $\quad \because \sqrt[n]{a} = a^{\frac{1}{n}}$

$= (27)^{\frac{1}{3}}(x^6)^{\frac{1}{3}}(y^9)^{\frac{1}{3}}(z^3)^{\frac{1}{3}}$ $\quad \because (ab)^m = a^m b^m$

$= (3^3)^{\frac{1}{3}}(x^6)^{\frac{1}{3}}(y^9)^{\frac{1}{3}}(z^3)^{\frac{1}{3}}$ $\quad \because (a^m)^n = a^{mn}$

$= 3^{3\times\frac{1}{3}} \cdot x^{6\times\frac{1}{3}} \cdot y^{9\times\frac{1}{3}} \cdot z^{3\times\frac{1}{3}}$

$= 3x^2y^3z$

(iii) $(64)^{-\frac{4}{3}}$ 09301044

Solution:

$(64)^{-\frac{4}{3}}$

$= \frac{1}{(64)^{\frac{4}{3}}}$

$= \frac{1}{(4)^{\frac{4}{3}}} = \frac{1}{4^{3\times\frac{4}{3}}}$

$= \frac{1}{4^4} = \frac{1}{256}$

Surds and their Applications

An irrational radical with rational radicand is called a surd. For example: $\sqrt{7}, \sqrt{2}, \sqrt[3]{11}$ are surds but $\sqrt{\pi}, \sqrt{e}$ are not surds.

The different type of surds are as follow:

(i) A surd that contains a single term is called a monomial e.g., $\sqrt{5}, \sqrt{7}$

(ii) A surd that contains the sum of two monomial surds is called a binomial surd e.g., $\sqrt{3}+\sqrt{5}, \sqrt{2}+\sqrt{7}$ etc.

(iii) $\sqrt{a}+\sqrt{b}$ and $\sqrt{a}-\sqrt{b}$ are called conjugate surds of each other.

Rationalization of denominator

To rationalize a denominator of the form $a + b\sqrt{x}$ or $a - b\sqrt{x}$, we multiply both the numerator and denominator by the conjugate factor.

Example 8: Rationalize the denominator of

(i) $\frac{3}{\sqrt{5}+\sqrt{2}}$ 09301045 (ii) $\frac{3}{\sqrt{5}-\sqrt{3}}$ 09301046

Solution (i):

www.ilmkidunya.com

$$\frac{3}{\sqrt{5}+\sqrt{2}}=\frac{3}{\sqrt{5}+\sqrt{2}}\times\frac{\sqrt{5}-\sqrt{2}}{\sqrt{5}-\sqrt{2}}$$

$$=\frac{3(\sqrt{5}-\sqrt{2})}{(\sqrt{5})^2-(\sqrt{2})^2}$$

$$=\frac{3(\sqrt{5}-\sqrt{2})}{5-2}$$

$$=\frac{3(\sqrt{5}-\sqrt{2})}{3}$$

$$=\sqrt{5}-\sqrt{2}$$

Solution (ii):

$$\frac{3}{\sqrt{5}-\sqrt{3}}=\frac{3}{\sqrt{5}-\sqrt{3}}\times\frac{\sqrt{5}+\sqrt{3}}{\sqrt{5}+\sqrt{3}}$$

$$=\frac{3\left(\sqrt{5}+\sqrt{3}\right)}{\left(\sqrt{5}\right)^2-\left(\sqrt{3}\right)^2}$$

$$=\frac{3\left(\sqrt{5}+\sqrt{3}\right)}{5-3}$$

$$=\frac{3\left(\sqrt{5}+\sqrt{3}\right)}{2}$$

Exercise 1.2

Q.1 Rationalize the denominator of following:

(i) $\frac{13}{4+\sqrt{3}}$ 09301047

Solution:

$$\frac{13}{4+\sqrt{3}}$$

Multiplying and dividing by $4-\sqrt{3}$

$$=\frac{13}{\left(4+\sqrt{3}\right)}\times\frac{\left(4-\sqrt{3}\right)}{\left(4-\sqrt{3}\right)}$$

$$=\frac{13\left(4-\sqrt{3}\right)}{(4)^2-\left(\sqrt{3}\right)^2}$$

$$=\frac{13\left(4-\sqrt{3}\right)}{16-3}$$

$$=\frac{13\left(4-\sqrt{3}\right)}{13}$$

$$=4-\sqrt{3}$$

(ii) $\frac{\sqrt{2}+\sqrt{5}}{\sqrt{3}}$ 09301048

Solution:

$$\frac{\sqrt{2}+\sqrt{5}}{\sqrt{3}}$$

Multiplying and dividing by $\sqrt{3}$

$$=\frac{\left(\sqrt{2}+\sqrt{5}\right)}{\sqrt{3}}\times\frac{\sqrt{3}}{\sqrt{3}}$$

$$=\frac{\sqrt{2}\times\sqrt{3}+\sqrt{5}\times\sqrt{3}}{\left(\sqrt{3}\right)^2}$$

$$=\frac{\sqrt{6}+\sqrt{15}}{3}$$

(iii) $\frac{\sqrt{2}-1}{\sqrt{5}}$ 09301049

Solution:

$$\frac{\sqrt{2}-1}{\sqrt{5}}$$

Multiplying and dividing by "$\sqrt{5}$"

$$=\frac{\left(\sqrt{2}-1\right)}{\sqrt{5}}\times\frac{\sqrt{5}}{\sqrt{5}}$$

$$=\frac{\left(\sqrt{2}-1\right)\sqrt{5}}{\left(\sqrt{5}\right)^2}$$

www.ilmkidunya.com

$= \dfrac{\sqrt{5}\left(\sqrt{2}-1\right)}{5}$

$= \dfrac{\sqrt{10}-\sqrt{5}}{5}$

(iv) $\dfrac{6-4\sqrt{2}}{6+4\sqrt{2}}$ 09301050

Solution:

$\dfrac{6-4\sqrt{2}}{6+4\sqrt{2}}$

Multiplying and dividing by $\left(6-4\sqrt{2}\right)$

$= \dfrac{\left(6-4\sqrt{2}\right)}{\left(6+4\sqrt{2}\right)} \times \dfrac{\left(6-4\sqrt{2}\right)}{6-4\sqrt{2}}$

$= \dfrac{\left(6-4\sqrt{2}\right)^2}{(6)^2-\left(4\sqrt{2}\right)^2}$ $[\because (a-b)^2 = a^2+b^2-2ab]$

$= \dfrac{(6)^2+\left(4\sqrt{2}\right)^2-2(6)\left(4\sqrt{2}\right)}{36-16(2)}$

$= \dfrac{36+16(2)-48\sqrt{2}}{36+32}$

$= \dfrac{36+32-48\sqrt{2}}{4}$

$= \dfrac{68-48\sqrt{2}}{4}$

$= \dfrac{4\left(17-12\sqrt{2}\right)}{4}$

$= 17-12\sqrt{2}$

(v) $\dfrac{\sqrt{3}-\sqrt{2}}{\sqrt{3}+\sqrt{2}}$ 09301051

Solution:

$\dfrac{\sqrt{3}-\sqrt{2}}{\sqrt{3}+\sqrt{2}}$

Multiplying and dividing by $\left(\sqrt{3}-\sqrt{2}\right)$

$= \dfrac{\left(\sqrt{3}-\sqrt{2}\right)}{\sqrt{3}+\sqrt{2}} \times \dfrac{\left(\sqrt{3}-\sqrt{2}\right)}{\sqrt{3}-\sqrt{2}}$

$= \dfrac{\left(\sqrt{3}-\sqrt{2}\right)^2}{\left(\sqrt{3}\right)^2-\left(\sqrt{2}\right)^2}$ $[\because (a-b)^2 = a^2+b^2-2ab]$

$= \dfrac{\left(\sqrt{3}\right)^2+\left(\sqrt{2}\right)^2-2\left(\sqrt{3}\right)\left(\sqrt{2}\right)}{3-2}$

$= \dfrac{3+2-2\sqrt{6}}{1}$

$= \dfrac{5-2\sqrt{6}}{1}$

$= 5-2\sqrt{6}$

vi) $\dfrac{4\sqrt{3}}{\sqrt{7}+\sqrt{5}}$

Solution:

$\dfrac{4\sqrt{3}}{\sqrt{7}+\sqrt{5}} \times \dfrac{\sqrt{7}-\sqrt{5}}{\sqrt{7}-\sqrt{5}}$

$= \dfrac{4\sqrt{3}\left(\sqrt{7}-\sqrt{5}\right)}{\left(\sqrt{7}\right)^2-\left(\sqrt{5}\right)^2}$

$= \dfrac{4\sqrt{3}\left(\sqrt{7}-\sqrt{5}\right)}{7-5}$

$= \dfrac{^{2}\cancel{4}\sqrt{3}\left(\sqrt{7}-\sqrt{5}\right)}{\cancel{2}}$

$= 2\sqrt{3}\left(\sqrt{7}-\sqrt{5}\right)$

Q.2 Simplify the following

(i) $\left(\dfrac{81}{16}\right)^{-\frac{3}{4}}$ 09301052

Solution:

$\left(\dfrac{81}{16}\right)^{-\frac{3}{4}}$

$\because \left(\dfrac{a}{b}\right)^{-m} = \left(\dfrac{b}{a}\right)^{m}$

www.ilmkidunya.com

$= \left(\frac{16}{81}\right)^{\frac{3}{4}}$

$= \left(\frac{2^4}{3^4}\right)^{\frac{3}{4}}$

$= \frac{2^{4\times\frac{3}{4}}}{3^{4\times\frac{3}{4}}}$

$= \frac{2^3}{3^3} = \frac{2\times2\times2}{3\times3\times3}$

$= \frac{8}{27}$

3	81
3	27
3	9
3	3
	1

(ii) $\left(\frac{3}{4}\right)^{-2} \div \left(\frac{4}{9}\right)^{3} \times \frac{16}{27}$ 09301053

Solution:

$\left(\frac{3}{4}\right)^{-2} \div \left(\frac{4}{9}\right)^{3} \times \frac{16}{27}$

$= \left(\frac{4}{3}\right)^{2} \times \left(\frac{9}{4}\right)^{3} \times \frac{2^4}{3^3}$

$= \left(\frac{2^2}{3}\right)^{2} \times \left(\frac{3^2}{2^2}\right)^{3} \times \frac{2^4}{3^3}$

$= \frac{2^4}{3^2} \times \frac{3^6}{2^6} \times \frac{2^4}{3^3}$

$= \frac{2^{4+4}\times3^6}{2^6\times3^{2+3}}$

$= \frac{2^8\times3^6}{2^6\times3^5}$

$= 2^8\times3^6\times2^{-6}\times3^{-5}$

$= 2^{8-6}\times3^{6-5}$

$= 2^2\times3^1$

$= 4\times3$

$= 12$

2	16
2	8
2	4
2	2
	1

3	27
3	9
3	3
	1

(iii) $(0.027)^{-\frac{1}{3}}$ 09301054

Solution:

$(0.027)^{-\frac{1}{3}}$

$= \left(\frac{27}{1000}\right)^{\frac{1}{3}}$

$= \left(\frac{1000}{27}\right)^{\frac{1}{3}} \because \left(\frac{a}{b}\right)^{-m} = \left(\frac{b}{a}\right)^{m}$

$= \left(\frac{10^3}{3^3}\right)^{\frac{1}{3}}$

$= \frac{10^{3\times\frac{1}{3}}}{3^{3\times\frac{1}{3}}}$

$= \frac{10}{3} = 3\frac{1}{3}$

(iv) $\sqrt[7]{\frac{x^{14}\times y^{21}\times z^{35}}{y^{14}z^7}}$ 09301055

Solution:

$\sqrt[7]{\frac{x^{14}\times y^{21}\times z^{35}}{y^{14}z^7}}$

$\sqrt[7]{x^{14}\times y^{21}\times z^{35}\times y^{-14}\times z^{-7}}$

$= \sqrt[7]{x^{14}\times y^{21-14}\times z^{35-7}}$

$= \left(x^{14}\times y^7\times z^{28}\right)^{\frac{1}{7}}$

$= x^{14\times\frac{1}{7}}\times y^{7\times\frac{1}{7}}\times z^{28\times\frac{1}{7}}$

$= x^2yz^4$

(v) $\frac{5\times(25)^{n+1} - 25\times(5)^{2n}}{5\times(5)^{2n+3} - (25)^{n+1}}$ 09301056

Solution:

$\frac{5\times(25)^{n+1} - 25\times(5)^{2n}}{5\times(5)^{2n+3} - (25)^{n+1}}$

$= \frac{5\cdot(5^2)^{n+1} - (5^2)\left(5^{2n}\right)}{5\cdot(5)^{2n+3} - (5^2)^{n+1}}$

www.ilmkidunya.com

$$= \frac{5\times(5^{2n+2})-5^{2n+2}}{5\times(5^{2n+2}\times 5^{1})-5^{2n+2}}$$

$$= \frac{5\times 5^{2n+2}-5^{2n+2}}{5^{2n+2}\times 25-5^{2n+2}}$$

$$= \frac{5^{2n+2}(5-1)}{5^{2n+2}\times 25-5^{2n+2}}$$

$$= \frac{5^{2n+2}(4)}{5^{2n+2}(25-1)}$$

$$= \frac{4}{24}$$

$$= \frac{1}{6}$$

(vi) $\dfrac{(16)^{x+1}+20(4^{2x})}{2^{x-3}\times 8^{x+2}}$ 09301057

Solution:

$$\frac{(16)^{x+1}+20(4^{2x})}{2^{x-3}\times 8^{x+2}}$$

$$= \frac{(2^4)^{x+1}+(5\times 2^2)(2^2)^{2x}}{2^{x-3}\times(2^3)^{x+2}}$$

$$= \frac{2^{4x+4}+(5\times 2^2)(2^{4x})}{2^{x-3}\times 2^{3x+6}}$$

$$= \frac{2^{4x+4}+5\times 2^{4x+2}}{2^{3x+6+x-3}}$$

$$= \frac{2^{4x+4}+5\times 2^{4x+2}}{2^{4x+3}}$$

$$= \frac{2^{4x+2}\times 2^2+5\times 2^{4x+2}}{2^{4x+2}\times 2^1}$$

$$= \frac{2^{4x+2}\times(2^2+5)}{2^{4x+2}\times(2)}$$

$$= \frac{4+5}{2}$$

$$= \frac{9}{2}$$

(vii) $\left(64^{-\frac{2}{3}}\div 9^{\frac{-3}{2}}\right)$ 09301058

Solution:

$$(64)^{-\frac{2}{3}}\div(9)^{-\frac{3}{2}}$$

$$= \frac{(64)^{-\frac{2}{3}}}{(9)^{-\frac{3}{2}}}$$

$$= \frac{9^{\frac{3}{2}}}{64^{\frac{2}{3}}} \qquad \left(\because \frac{a^{-m}}{b^{-n}}=\frac{b^n}{a^m}\right)$$

$$= \frac{(3^2)^{\frac{3}{2}}}{(2^6)^{\frac{2}{3}}}$$

$$= \frac{3^{2\times\frac{3}{2}}}{2^{6\times\frac{2}{3}}}$$

$$= \frac{3^3}{2^4}$$

$$= \frac{27}{2^4}$$

$$= \frac{27}{16}$$

(viii) $\dfrac{3^n\times 9^{n+1}}{3^{n-1}\times 9^{n-1}}$ 09301059

Solution:

$$\frac{3^n\times 9^{n+1}}{3^{n-1}\times 9^{n-1}}$$

$$= \frac{3^n\times(3^2)^{n+1}}{3^{n-1}\times(3^2)^{n-1}}$$

$$= \frac{3^n\times 3^{2n+2}}{3^{n-1}\times 3^{2n-2}}$$

$$= \frac{3^{n+2n+2}}{3^{n-1+2n-2}}$$

$$= \frac{3^{3n+2}}{3^{3n-3}}$$

$$= 3^{3n+2}\times 3^{-3n+3}$$

www.ilmkidunya.com

$= 3^{3n+2-3n+3} \quad \left(\because \frac{a^m}{a^n} = a^{m-n}\right)$

$= 3^5$

$= 3\times3\times3\times3\times3$

$= 243$

(ix) $\dfrac{5^{n+3} - 6.5^{n+1}}{9\times5^n - 4\times5^n}$ 09301060

Solution:

$$\frac{5^{n+3} - 6.5^{n+1}}{9\times5^n - 4\times5^n}$$

$$= \frac{5^n\times5^3 - 6.5^n.5^1}{9\times5^n - 4\times5^n}$$

$$= \frac{5^n\left(5^3 - 6\times5\right)}{5^n\left(9-4\right)}$$

$$= \frac{5^3 - 6\times5}{9-4}$$

$$= \frac{125-30}{9-4}$$

$$= \frac{95}{5}$$

$$= 19$$

Q.3 If $x = 3+\sqrt{8}$ then find the value of

(i) $x+\frac{1}{x}$ (ii) $x-\frac{1}{x}$ (iii) $x^2+\frac{1}{x^2}$

(iv) $x^2-\frac{1}{x^2}$ (v) $x^4+\frac{1}{x^4}$ (vi) $\left(x-\frac{1}{x}\right)^2$

Solution:

$x = 3+\sqrt{8}$ _____(i)

$$\frac{1}{x} = \frac{1}{3+\sqrt{8}}$$

$$\frac{1}{x} = \frac{1}{\left(3+\sqrt{8}\right)}\times\frac{\left(3-\sqrt{8}\right)}{\left(3-\sqrt{8}\right)}$$

$$\frac{1}{x} = \frac{3-\sqrt{8}}{(3)^2-\left(\sqrt{8}\right)^2}$$

$$\frac{1}{x} = \frac{3-\sqrt{8}}{9-8}$$

$$\frac{1}{x} = \frac{3-\sqrt{8}}{1}$$

$\frac{1}{x} = 3-\sqrt{8}$ _____ (ii)

(i) Finding $x+\frac{1}{x}$ 09301061

Adding eq. (i) and (ii)

$$x+\frac{1}{x} = 3+\sqrt{8}+3-\sqrt{8}$$

$x+\frac{1}{x} = 6$ _____ (iii)

(ii) Finding $x-\frac{1}{x}$ 09301062

Subtracting eq.(i) from (ii)

$$x-\frac{1}{x} = \left(3+\sqrt{8}\right)-\left(3-\sqrt{8}\right)$$

$$x-\frac{1}{x} = 3+\sqrt{8}-3+\sqrt{8}$$

$x+\frac{1}{x} = 2\sqrt{8}$ ----------(iv)

(iii) Finding $x^2+\frac{1}{x^2}$ 09301063

Taking square of eq. (iii)

$$\left(x+\frac{1}{x}\right)^2 = (6)^2$$

$$(x)^2+\left(\frac{1}{x}\right)^2+2(x)\left(\frac{1}{x}\right) = 36$$

$$x^2+\frac{1}{x^2} = 36-2$$

$x^2+\frac{1}{x^2} = 34$ ----------(v)

(iv) Finding $x^2-\frac{1}{x^2}$ 09301063a

We know that

$$x^2-\frac{1}{x^2} = (x)^2-\left(\frac{1}{x}\right)^2$$

$$x^2-\frac{1}{x^2} = \left(x+\frac{1}{x}\right)\left(x-\frac{1}{x}\right)$$

www.ilmkidunya.com

Putting the values from (iii) and (iv)

$$x^2-\frac{1}{x^2}=(6)\left(2\sqrt{8}\right)=12\sqrt{8}$$

(v) Finding $x^4+\frac{1}{x^4}$ 09301063b

Taking square of equation (v)

$$\left(x^2+\frac{1}{x^2}\right)^2=(34)^2$$

$$\left(x^2\right)^2+\left(\frac{1}{x^2}\right)^2+2\left(x^2\right)\left(\frac{1}{x^2}\right)=1156$$

$$x^4+\frac{1}{x^4}+2=1156$$

$$x^4+\frac{1}{x^4}=1156-2$$

$$x^4+\frac{1}{x^4}=1154$$

(vi) Finding $\left(x-\frac{1}{x}\right)^2$ 09301063c

Taking square of equation (iv)

$$\left(x-\frac{1}{x}\right)^2=\left(2\sqrt{8}\right)^2$$

$$\left(x-\frac{1}{x}\right)^2=(2)^2(\sqrt{8})^2$$

$$\left(x-\frac{1}{x}\right)^2=4(8)$$

$$\left(x-\frac{1}{x}\right)^2=32$$

Q.4 Find the rational numbers p and q such that $\frac{8-3\sqrt{2}}{4+3\sqrt{2}}=p+q\sqrt{2}$ 09301064

Solution:

$$\mathbf{p+q\sqrt{2}=\frac{8-3\sqrt{2}}{4+3\sqrt{2}}}$$

$$P+q\sqrt{2}=\frac{\left(8-3\sqrt{2}\right)}{\left(4+3\sqrt{2}\right)}\times\frac{\left(4-3\sqrt{2}\right)}{\left(4-3\sqrt{2}\right)}$$

$$=\frac{32-24\sqrt{2}-12\sqrt{2}+\left(3\sqrt{2}\right)^2}{(4)^2-\left(3\sqrt{2}\right)^2}$$

$$=\frac{32-36\sqrt{2}+9(2)}{16-9(2)}$$

$$=\frac{32-36\sqrt{2}+18}{16-18}$$

$$=\frac{50-36\sqrt{2}}{-2}$$

$$=\frac{50}{-2}-\frac{36\sqrt{2}}{-2}$$

$p+q\sqrt{2}=-25+18\sqrt{2}$

By comparing both sides.

$\Rightarrow p=-25, q=18$

Q.5 Simplify the following:

(i) $\frac{(25)^{\frac{3}{2}}\times(243)^{\frac{3}{5}}}{(16)^{\frac{5}{4}}\times(8)^{\frac{4}{3}}}$ 09301065

3	243
3	81
3	27
3	9
3	3
	1

Solution

$$\frac{(25)^{\frac{3}{2}}\times(243)^{\frac{3}{5}}}{(16)^{\frac{5}{4}}\times(8)^{\frac{4}{3}}}$$

$$=\frac{\left(5^2\right)^{\frac{3}{2}}\times\left(3^5\right)^{\frac{3}{5}}}{\left(2^4\right)^{\frac{5}{4}}\times\left(2^3\right)^{\frac{4}{3}}}$$

$$=\frac{5^{2\times\frac{3}{2}}\times3^{5\times\frac{3}{5}}}{2^{4\times\frac{5}{4}}\times2^{3\times\frac{4}{3}}}$$

$$=\frac{5^3\times3^3}{2^5\times2^4}$$

$$=\frac{(5\times3)^3}{2^{5+4}}\qquad\left(\because a^m\times a^n=a^{m+n}\right)$$

$$=\frac{(15)^3}{2^9}$$

$$=\frac{15\times15\times15}{2\times2\times2\times2\times2\times2\times2\times2\times2}$$

$$=\frac{3375}{512}$$

www.ilmkidunya.com

(ii) $\dfrac{54\times\sqrt[3]{(27)^{2x}}}{9^{x+1}+216(3^{2x-1})}$ 09301066

Solution:

$$\frac{54\times\sqrt[3]{(27)^{2x}}}{9^{x+1}+216(3^{2x-1})}$$

2	54
3	27
3	9
3	3
	1

2	216
2	108
2	54
3	27
3	9
3	3
	1

$$= \frac{(2\times3^3)\times\sqrt[3]{(3^3)^{2x}}}{(3^2)^{x+1}+2^3\times3^3(3^{2x-1})}$$

$$= \frac{(2\times3^3)\times\sqrt[3]{(3^{2x})^3}}{3^{2x+2}+2^3\times3^{2x-1+3}}$$

$$= \frac{2\times3^3\times3^{2x}}{3^{2x+2}+2^3\times3^{2x+2}} \qquad \left(\because \sqrt[m]{a^m}=a\right)$$

$$= \frac{2\times3^{2x+3}}{3^{2x+2}(1+2^3)}$$

$$= \frac{2\times3^{2x+3-2x-2}}{(1+8)} \qquad \left(\because \frac{a^m}{a^n}=a^{m-n}\right)$$

$$= \frac{2\times3^1}{9}$$

$$= \frac{6}{9}$$

$$= \frac{2}{3}$$

(iii) $\sqrt{\dfrac{(216)^{\frac{2}{3}}\times(25)^{\frac{1}{2}}}{(0.04)^{\frac{-3}{2}}}}$ 09301067

Solution:

$$\sqrt{\frac{(216)^{\frac{2}{3}}\times(25)^{\frac{1}{2}}}{(0.04)^{\frac{-3}{2}}}}$$

2	216
2	108
2	54
3	27
3	9
3	3
	1

$$= \sqrt{(216)^{\frac{2}{3}}\times(25)^{\frac{1}{2}}\times(0.04)^{\frac{3}{2}}}$$

$$= \sqrt{(2^3\times3^3)^{\frac{2}{3}}\times(5^2)^{\frac{1}{2}}\times\left(\frac{4}{100}\right)^{\frac{3}{2}}}$$

$$= \sqrt{2^{3\times\frac{2}{3}}\times3^{3\times\frac{2}{3}}\times5^{2\times\frac{1}{2}}\times\left(\frac{1}{25}\right)^{\frac{3}{2}}}$$

$$= \sqrt{2^2\times3^2\times5^1\times\frac{1}{(5^2)^{\frac{3}{2}}}}$$

$$= \sqrt{(2\times3)^2\times5^1\times\frac{1}{5^3}}$$

$$= \sqrt{(6)^2\times\frac{1}{5^{3-1}}}$$

$$= \sqrt{\frac{6^2}{5^2}}$$

$$= \sqrt{\left(\frac{6}{5}\right)^2}$$

$$= \frac{6}{5}$$

(iv) $\left(a^{\frac{1}{3}}+b^{\frac{2}{3}}\right)\times\left(a^{\frac{2}{3}}-a^{\frac{1}{3}}b^{\frac{2}{3}}+b^{\frac{4}{3}}\right)$ 09301068

Solution:

$$\left(a^{\frac{1}{3}}+b^{\frac{2}{3}}\right)\times\left(a^{\frac{2}{3}}-a^{\frac{1}{3}}b^{\frac{2}{3}}+b^{\frac{4}{3}}\right)$$

$$=\left(a^{\frac{1}{3}}+b^{\frac{2}{3}}\right)\times\left[\left(a^{\frac{1}{3}}\right)^2-\left(a^{\frac{1}{3}}\right)\left(b^{\frac{2}{3}}\right)+\left(b^{\frac{2}{3}}\right)^2\right]$$

$$\boxed{\because (x+y)(x^2-xy+y^2)=x^3+y^3}$$

$$\therefore \quad = \left(a^{\frac{1}{3}}\right)^3+\left(b^{\frac{2}{3}}\right)^3$$

$$= a^{\frac{1}{3}\times3}+b^{\frac{2}{3}\times3}$$

$$= a+b^2$$

Applications of Real Numbers in Daily Life:

Example 9: The sum of two real numbers is 8, and their difference is 2. Find the numbers. 09301069

www.ilmkidunya.com

Solution:

Let a and b be two real numbers then

$a + b = 8$ ----------(i)

$a - b = 2$ ----------(ii)

Add eq. (i) and eq. (ii)

$2a = 10$

$\Rightarrow a = 5$

Put it in eq. (i)

$\Rightarrow 5 - b = 2$

$\Rightarrow -b = 2 - 5$


$\Rightarrow -b = -3$

$\Rightarrow b = 3$

So, 5 and 3 are required real numbers

Temperature Conversions

In the figure, three types of thermometers are shown.



We can convert three temperature scales Celsius, Fahrenheit and kelvin with each other.

Conversion formulae are given below:

(i) $K = °C + 273$ 09301070

(ii) $°C = \frac{5}{9}(F - 32)°$ 09301071

(iii) $°F = \frac{9°C}{5} + 32$ 09301072

Where K,C and F shows the kelvin, Celsius and Fahrenheit scales respectively.

Example 10: Normal human body temperature is 98.6 F. Convert it into Celsius and kelvin scale. 09301073

Solution:

Given that

$°F = 98.6$

So Convert it into Celsius scale, we use

$$°C = \frac{5}{9}(F - 32)°$$

$$°C = \frac{5}{9}(98.6 - 32)°$$

$$°C = \frac{5}{9}(66.6)°$$

$$°C = (0.55)(66.6)°$$

$$°C = 37°$$

Hence, normal human body temperature at Celsius scale is 37°

Now, we convert it into Kelvin scale

$$K = C + 273°$$

$$K = 37° + 273°$$

$$K = 310 \text{ Kelvin}$$

Profit and Loss:

The profit and loss can be calculated by the following formula.

(i) Profit = Selling price – Cost Price

Profit = SP – CP

$$\text{Profit \%} = \left(\frac{\text{Profit}}{CP} \times 100\right)\%$$

(ii) Loss = Cost Price – Selling Price

Loss = CP – SP

$$\text{Loss \%} = \left(\frac{loss}{CP} \times 100\right)\%$$

Example 11: Hamail purchased a bicycle for Rs. 6590 and sold it for Rs. 6850. Find the profit percentage. 09301074

Solution:

Cost Price = CP = Rs. 6590

Selling Price = SP = Rs. 6850

Profit = SP – CP

= 6850 – 6590

= Rs 260

Now, we find the profit percentage.

$$\text{Profit \%} = \left(\frac{profit}{CP} \times 100\right)\%$$

www.ilmkidunya.com

$$= \left(\frac{260\times100}{6590}\right)\%$$

$$= 3.94\% \approx 4\%$$

Example 12: Umair bought a book for Rs. 850 and sold it for Rs. 720. What was his loss percentage? 09301075

Solution:

Cost price of book = CP = Rs. 850

Selling price of book = SP = Rs. 720

Loss = CP – SP

= 850 – 720

= Rs. 130

$$\text{Loss percentage} = \left(\frac{Loss}{CP}\times100\right)\%$$

$$= \left(\frac{130}{850}\times100\right)\%$$

$$= 15.29\%$$

Example 13: Mr. Saleem, Nadeem, and Tanveer earned a profit of Rs. 450,000 from a business. If their investments in the business are in the ratio 4: 7: 14, find each person's profit. 09301076

Solution:

Profit earned = Rs. 450,000

Given ratios = 4: 7: 14

Sum of ratio = 4 + 7 + 14 = 25

$$\text{Saleem earned profit} = \frac{4}{25}\times450{,}000$$

= Rs. 72,000

$$\text{Nadeem earned profit} = \frac{7}{25}\times450{,}000$$

= Rs. 126,000

$$\text{Tanveer earned profit} = \frac{14}{25}\times450{,}000$$

= Rs. 252,000

Example 14: If the simple profit on Rs. 6,400 for 12 years is Rs. 3,840. Find the rate of profit. 09301077

Solution:

Principal = Rs. 6400

Simple profit = Rs. 3840

Time = 12 years

To find the rate we use the following formula

$$\text{Rate} = \frac{amount\ of\ profit\times100}{time\times principal}$$

$$= \frac{3840\times100}{12\times6400}$$

$$= 5\%$$

Thus, rate of profit is 5%.

Exercise 1.3

Q.1 The sum of three consecutive integers is forty-two, find three integers. 09301078

Solution:

Let x, $x+1$, $x+2$ be three consecutive integers

By condition

$x + (x+1) + (x+2) = 42$

$x + x + 1 + x + 2 = 42$

$3x + 3 = 42$

$3x = 42 - 3$

$3x = 39$

$x = \frac{39}{3}$

$x = 13$

Now the 1st integer = x = 13

2nd integer = $x+1$ = 13+1 = 14

3rd integer = $x+2$ = 13+2 = 15


Thus 13,14 and 15 are required three consecutive integers.

Q.2 The diagram shows right angled ΔABC in which the length of $\overline{AC}$ is $(\sqrt{3}+\sqrt{5})$ cm. The area of Δ ABC is $(1+\sqrt{15})$ cm^2, Find the length $\overline{AB}$ in the

www.ilmkidunya.com

form $(a\sqrt{3}+b\sqrt{5})$ cm where a and b are integers.

09301079



Solution:

Given


$m\overline{AC} = (\sqrt{3}+\sqrt{5})$ cm

Area of $\Delta ABC = (1+\sqrt{15})$ cm^2

To find = $m\overline{AB}$ = ?

We know that, Area of $\Delta = \frac{1}{2}(b \times h)$

Area of $\Delta ABC = \frac{1}{2}(m\overline{AC} \times m\overline{AB})$



$$1+\sqrt{15} = \frac{1}{2}\left[(\sqrt{3}+\sqrt{5}) \times m\overline{AB}\right]$$

$$\Rightarrow \frac{2(1+\sqrt{15})}{\sqrt{3}+\sqrt{5}} = m\overline{AB}$$

$\Rightarrow$ Multiplying and dividing by $(\sqrt{3}-\sqrt{5})$

$$\Rightarrow m\overline{AB} = \frac{(2+2\sqrt{15})}{(\sqrt{3}+\sqrt{5})} \times \frac{(\sqrt{3}-\sqrt{5})}{\sqrt{3}-\sqrt{5}}$$

$$= \frac{2\sqrt{3}-2\sqrt{5}+2\sqrt{15}\times\sqrt{3}-2\sqrt{15}\times\sqrt{5}}{(\sqrt{3})^2-(\sqrt{5})^2}$$

$\because a^2-b^2 = (a+b)(a-b)$

$$= \frac{2\sqrt{3}-2\sqrt{5}+2\sqrt{45}-2\sqrt{75}}{3-5}$$

$$= \frac{2(\sqrt{3}-\sqrt{5}+\sqrt{9\times5}-\sqrt{25\times3})}{-2}$$

$$= \frac{\sqrt{3}-\sqrt{5}+3\sqrt{5}-5\sqrt{3}}{-1}$$

$$= \frac{-4\sqrt{3}+2\sqrt{5}}{-1}$$

$$= 4\sqrt{3}-2\sqrt{5}$$

Thus $m\overline{AB} = (4\sqrt{3}-2\sqrt{5})$ cm.

Q.3 A rectangle has sides of length $(2+\sqrt{18})$ m and $(5-\frac{4}{\sqrt{2}})$ m. Express the area of the rectangle in the form $a+b\sqrt{2}$ where a and b are integers.

Solution:

09301080

Let length of rectangle = L = $(2+\sqrt{18})$ m

Width of rectangle = W = $\left(5-\frac{4}{\sqrt{2}}\right)$ m

We know that

Area of Rectangle = A = L × W

$$A = (2+\sqrt{18})\times\left(5-\frac{4}{\sqrt{2}}\right)$$

$$A = \left(2+\sqrt{9\times2}\right)\times\left(5-\frac{2\times2}{\sqrt{2}}\right)$$

$$A = (2+3\sqrt{2})\times(5-2\sqrt{2}) \qquad \left(\because \frac{a}{\sqrt{a}} = \sqrt{a}\right)$$

$$A = 10 - 4\sqrt{2} + 15\sqrt{2} - (3\sqrt{2})(2\sqrt{2})$$

$$A = 10 + 11\sqrt{2} - 6\sqrt{4}$$

$$A = 10 + 11\sqrt{2} - 6(2)$$

$$A = 10 + 11\sqrt{2} - 12$$

$$A = -2 + 11\sqrt{2}$$

Thus area of rectangle is $(-2+11\sqrt{2})m^2$

www.ilmkidunya.com

Q.4 Find two numbers whose sum is 68 and whose difference is 22. 09301081

Solution:

Sum = 68

Difference = 22

Let x and y be required numbers

By given conditions:

Sum: $x + y = 68$ _____ (i)

Diff. $x - y = 22$ _____ (ii)

Adding eq. (i) and (ii),

$$x + \cancel{y} = 68$$
$$x - \cancel{y} = 22$$
$$2x = 90$$
$$x = \frac{90}{2}$$
$$x = 45$$

Put it in eq. (i),

$x + y = 68$

$45 + y = 68$

$y = 68 - 45$

$y = 23$

Thus required numbers are 45 and 23.

Q.5 The weather in Lahore was unusually warm during the summer of 2024. The TV news reported temperatures as high as 48°C. By using the formula, $\left(°F = \frac{9}{5}°C + 32\right)$ find the temperature as Fahrenheit scale. 09301082

Solution:

Temperature in degree centigrade = 48°C

formula: $°F = \frac{9}{5}°C + 32$

Temperature in Fahrenheit,

$$°F = \frac{9}{5} \times 48 + 32$$
$$°F = 9 \times 9.6 + 32$$
$$°F = 86.4 + 32$$
$$°F = 118.4$$

Q.6 The sum of the ages of the father and son is 72. Six years ago the father's age was 2 times the age of the son. What was Son's age six years ago? 09301083

Solution:

Let father's age = x

Son's age = y

By condition:

$x + y = 72$ _____ (i)

Six year ago,

Father's age = $x - 6$

Son's age = $y - 6$

By condition:

Father's age = 2 times the son's age

$(x - 6) = 2(y - 6)$

$x - 6 = 2y - 12$

$x - 2y = 6 - 12$

$x - 2y = -6$ _____ (ii)

Subtracting eq. (ii) from (i)

$$\cancel{x} + y = 72$$
$$\cancel{x} - 2y = -6$$
$$3y = 78$$
$$y = \frac{78}{3}$$
$$\boxed{y = 26}$$

Put y = 26 in eq.(i)

$x + 26 = 72$

$x = 72 - 26$

$\boxed{x = 46}$

Six years ago,

Son's age = $y - 6 = 26 - 6 = 20$ years

Father's age = $x - 6 = 46 - 6 = 40$ years

Q.7 Mirha bought a toy for Rs.1500 and sold for Rs.1520. What was her profit percentage? 009301084

Solution:

The cost price=CP=Rs.1500

The selling price=SP=Rs.1520

The profit amount=SP – CP

=Rs.1520 – Rs.1500=Rs.20

$$\text{Profit percentage} = \frac{\text{Profit}}{\text{C.P}} \times 100\%$$

www.ilmkidunya.com

Profit percentage $= \frac{20}{1500} \times 100\%$

$= \frac{20}{15 \times \cancel{100}} \times \cancel{100}\% = 1.33\%$

Q.8 The annual income of Tayab is Rs. 960,000, while the exempted amount is Rs. 130,000. How much tax would he have to pay at the rate of 0.75%.

09301085

Solution:

Annual income = Rs.9,60,000/-
Exempted amount = Rs.130,000/-
Tax rate = 0.75%
We know that
Taxable income
= Annual income – exempted amount
Taxable income = 960,000 – 130,000
Taxable income = 830,000
Tax amount = 0.75% of Taxable income

$= \frac{0.75}{100} \times 830,000$

$= \frac{75}{100 \times 100} \times 830,000$

$= 75 \times 83$

= Rs.6,225

Thus Tayab will pay tax of Rs.6,225.

Q.9 Find the compound markup on Rs. 375,000 for one year at the rate of 14% compounded markup annually.

09301086

Solution:

Principal amount = P = Rs.375,000/-
Time, = t = 1year
Rate, R = 14%
We know that
Compound markup = P×T×R

$= 375,000 \times 1 \times 14\%$

$= 375,000 \times \frac{14}{100}$

$= 3,750 \times 14$

= Rs. 52,500/-

Thus compound mark up is Rs.52,500/-

Review Exercise 1

Q.1 Choose the correct option.

i. $\sqrt{7}$ is: 09301087
(a) Integer
(b) Rational number
(c) Irrational number
(d) Natural number

ii. π and e are: 09301088
(a) Natural number
(b) Integers
(c) Rational number
(d) Irrational number

iii. If n is not a perfect square then $\sqrt{n}$ is: 09301089
(a) Rational number
(b) Natural number
(c) Integer
(d) Irrational number

iv. $\sqrt{3}+\sqrt{5}$ is: 09301090
(a) Whole number
(b) Integer
(c) Rational number
(d) Irrational number

v. For all $x \in R$, $x = x$ is called: 09301091
(a) Reflexive property
(b) Transitive number
(c) Symmetric property
(d) Trichotomy property

vi. Let $a, b, c \in R$ then $a > b$ and $b > c$ $\Rightarrow a > c$ is called _____property. 09301092
(a) Trichotomy (b) Transitive
(c) Additive (d) Multiplicative

vii. $2^x \times 8^x = 64$ then $x =$ 09301093
(a) $\frac{3}{2}$ (b) $\frac{3}{4}$
(c) $\frac{5}{6}$ (d) $\frac{2}{3}$

www.ilmkidunya.com

viii. Let $a, b, \in R$ then $a = b$ and $b = a$ is called _____property. 09301094

(a) Reflexive (b) Symmetric
(c) Transitive (d) Additive

ix. $\sqrt{75} + \sqrt{27} =$ 09301095

(a) $\sqrt{102}$ (b) $9\sqrt{3}$
(c) $5\sqrt{3}$ (d) $8\sqrt{3}$

x. The product of $(3+\sqrt{5})(3-\sqrt{5})$ is: 09301096

(a) Prime number
(b) odd number
(c) Irrational number
(d) Rational number

Answer Key

i	c	ii	d	iii	d	iv	d	v	a
vi	b	vii	a	viii	b	ix	d	x	d

Multiple Choice Questions (Additional)

History of Real numbers

1. Which number system was used by the Sumerians? 09301097
(a) Decimal (b) hexadecimal
(c) Sexagesimal (d) Binary

2. The sexagesimal system is a number system with the base: 09301098
(a) 2 (b) 10
(c) 16 (d) 60

3. Which number system was used by the Egyptians? 09301099
(a) Decimal (b) hexadecimal
(c) Sexagesimal (d) Binary

4. How many letters are used in Roman numeral system? 09301100
(a) 3 (b) 5
(c) 7 (d) 10

5. In Roman counting the letter "L" represents the number: . 09301101
(a) 10 (b) 50
(c) 100 (d) 500

6. The invention of zero is attributed to: 09301102
(a) Arabs (b) Egyptians
(c) Sumerians (d) Indians

7. Which number system is known as Indo-Arabic numerals? 09301103
(a) Decimal (b) hexadecimal
(c) Sexagesimal (d) Binary

8. Who did introduce the numerals (0-9) to Europe? 09301104
(a) Arabs (b) Egyptians
(c) Sumerians (d) Indian

9. Which of the following is one of the modern number systems? 09301105
(a) Roman Numerals
(b) Egyptians numerals
(c) Sexagesimal system
(d) hexadecimal system

Real numbers

10. $Q \cup Q' =$ ____ 09301106

(a) Q' (b) Q
(c) R (d) ϕ

11. $Q \cap Q' =$ ____ 09301107
(a) Q' (b) Q
(c) R (d) ϕ

12. Q and Q' are _____sets. 09301108
(a) disjoint (b) over lapping
(c) intersecting (d) supper

13. For each prime number P, $\sqrt{P}$ is an: 0930110
(a) Irrational (b) Rational
(c) Real (d) Whole

www.ilmkidunya.com

Properties of real numbers

14. Name the property of real numbers used in $\pi + (-\pi) = 0$. 09301110
(a) Additive inverse
(b) Multiplicative inverse
(c) Additive identity
(d) Multiplicative identity

15. Name the property of real numbers used in $\frac{1}{2} \times 1 = \frac{1}{2}$. 09301111
(a) Additive identity
(b) Additive Inverse
(c) Multiplicative identity
(d) Multiplicative Inverse

16. If $x < y$ and $z < 0$ then: 09301112
(a) $xz < yz$ (b) $xz > yz$
(c) $xz = yz$ (d) $\ddot{x} > y$

17. If a, b∈ R then only one of a = b or a < b or a > b holds is called --- property. 093011
(a) Trichotomy (b) Transitive
(c) Additive (d) Multiplicative

Radical Expressions

18. In $\sqrt[n]{a}$, the symbol $\sqrt{}$ is called: 09301114
(a) radical sign (b) index
(c) exponent (d) base

19. In $\sqrt[n]{a^m}$ 'n' is called: 09301115
(a) base (b) radical sign
(c) index (d) radical

20. $(27x)^{\frac{2}{3}} =$ _____ 09301116
(a) $\frac{\sqrt[3]{x^2}}{9}$ (b) $\frac{\sqrt{x^3}}{9}$
(c) $\frac{\sqrt[3]{x^2}}{8}$ (d) $9\sqrt[3]{x^2}$

21. Write $\sqrt[5]{x}$ in exponential form 09301117
(a) x (b) x^5
(c) $x^{\frac{1}{5}}$ (d) $x^{\frac{5}{2}}$

22. Writing $m^{\frac{2}{3}}$ with radical sign we get: 09301118
(a) $\sqrt[3]{m^2}$ (b) $\sqrt{m^3}$
(c) $\sqrt[2]{m^3}$ (d) $\sqrt{m^6}$

23. In $\sqrt[3]{5}$, the radicand is: 09301119
(a) 3 (b) $\frac{1}{3}$
(c) 5 (d) 35

Surds

24. Which of the following is a Surd? 09301120
(a) $\sqrt{7}$ (b) $\sqrt{9}$
(c) $\sqrt{\pi}$ (d) $\sqrt{e}$

25. Which of the following is a binomial Surd? 09301121
(a) $4\sqrt{3}$ (b) $\sqrt{16}$
(c) $7+\sqrt{\pi}$ (d) $2-\sqrt{3}$

26. A surd which contains a single term is called surd. 09301122
(a) Monomial (b) Binomial
(c) Trinomial (d) None

27. Conjugate factor of the Surd $a+b\sqrt{x}$ is: 09301123
(a) $a+b\sqrt{x}$ (b) $a-b\sqrt{x}$
(c) $-a-b\sqrt{x}$ (d) $-a+b\sqrt{x}$

28. $(4+\sqrt{2})(4-\sqrt{2})$ is equal to: 09301124
(a) 14 (b) −14
(c) 12 (d) 8

29. $(\sqrt{x}+\sqrt{y})(\sqrt{x}-\sqrt{y})(x+y) = \ldots$ 09301125
(a) $(x+y)$ (b) $(x-y)$
(c) (x^2+y^2) (d) (x^2-y^2)

30. $\frac{1}{2-\sqrt{3}} =$ ____ 09301126
(a) $2+\sqrt{3}$ (b) $2-\sqrt{3}$
(c) $-2+\sqrt{3}$ (d) $-2-\sqrt{3}$

Answer Key

1	c	2	d	3	a	4	c	5	b	6	d	7	a	8	a	9	d	10	c
11	d	12	a	13	a	14	a	15	c	16	b	17	a	18	a	19	c	20	d
21	c	22	a	23	c	24	a	25	d	26	a	27	b	28	a	29	d	30	a

www.ilmkidunya.com

Q.2 If $a=\frac{3}{2}, b=\frac{5}{3}$ and $c=\frac{7}{5}$ then verify that

(i) $a(b+c)=ab+ac$ 09301127

Solution:

$a(b+c)=ab+ac$

L.H.S $= a(b+c)$

$$=\frac{3}{2}\left(\frac{5}{3}+\frac{7}{5}\right)$$

$$=\frac{3}{2}\left(\frac{25+21}{15}\right)$$

$$=\frac{3}{2}\left(\frac{46}{15}\right)$$

$$=\frac{23}{5} \quad \text{(i)}$$

Now, R.H.S $= ab+ac$

$$=\frac{3}{2}\times\frac{5}{3}+\frac{3}{2}\times\frac{7}{5}$$

$$=\frac{5}{2}+\frac{21}{10}$$

$$=\frac{25+21}{10}$$

$$=\frac{46}{10}$$

$$=\frac{23}{5} \quad \text{(ii)}$$

From (i) and (ii), L.H.S = R.H.S

Hence $a(b+c)=ab+ac$

(ii) $(a+b)c=ac+bc$ 09301128

Solution

$(a+b)c=ac+bc$

L.H.S $= (a+b)c$

$$=\left(\frac{3}{2}+\frac{5}{3}\right)\times\frac{7}{5}$$

$$=\left(\frac{9+10}{6}\right)\times\frac{7}{5}$$

$$=\frac{19}{6}\times\frac{7}{5}$$

$$=\frac{133}{30} \quad \text{(i)}$$

Now, R.H.S $= ac+bc$

$$=\frac{3}{2}\times\frac{7}{5}+\frac{5}{3}\times\frac{7}{5}$$

$$=\frac{21}{10}+\frac{7}{3}$$

$$=\frac{63+70}{30}$$

$$=\frac{133}{30} \quad \text{(ii)}$$

From (i) and (ii), L.H.S = R.H.S

Hence $(a+b)c=ac+bc$ is proved

Q.3 If $a=\frac{4}{3}, b=\frac{5}{2}, c=\frac{7}{4}$, then verify the associative property of real numbers w.r.t addition and multiplication. 09301129

Solution:

(i) Associative property w.r.t addition.

$(a+b)+c=a+(b+c)$

L.H.S $= (a+b)+c$

$$=\left(\frac{4}{3}+\frac{5}{2}\right)+\frac{7}{4}$$

$$=\left(\frac{8+15}{6}\right)+\frac{7}{4}$$

$$=\frac{23}{6}+\frac{7}{4}$$

$$=\frac{46+21}{12}$$

$$=\frac{67}{12} \quad \text{(i)}$$

2	6 –4
2	3–2
3	3–1
	1–1

R.H.S. $= a+(b+c)$

Now, $=\frac{4}{3}+\left(\frac{5}{2}+\frac{7}{4}\right)$

$$=\frac{4}{3}+\left(\frac{10+7}{4}\right)$$

$$=\frac{4}{3}+\frac{17}{4}$$

$$=\frac{16+51}{12}$$

$$=\frac{67}{12} \quad \text{(ii)}$$

From(i) and (ii) L.H.S = R.H.S.

Hence $a+(b+c)=(a+b)+c$

www.ilmkidunya.com

(ii) Associative property w.r.t multiplication. $(a\times b)\times c = a\times(b\times c)$ 09301130

Solution:

L.H.S $= (a\times b)\times c$

$$= \left(\frac{4}{3}\times\frac{5}{2}\right)\times\frac{7}{4}$$

$$= \frac{20}{6}\times\frac{7}{4}$$

$$= \frac{10}{3}\times\frac{7}{4}$$

$$= \frac{70}{12}$$

$$= \frac{35}{6} \quad \text{(i)}$$

Now, R.H.S $= a\times(b\times c)$

$$= \frac{4}{3}\times\left(\frac{5}{2}\times\frac{7}{4}\right)$$

$$= \frac{4}{3}\times\left(\frac{35}{8}\right)$$

$$= \frac{\cancel{4}^{1}}{3}\times\left(\frac{35}{\cancel{8}_{2}}\right)$$

$$= \frac{35}{6} \quad \text{(ii)}$$

From (i) and (ii) L.H.S = R.H.S

Hence, $(a\times b)\times c = a\times(b\times c)$

Q.4 Is 0 a rational number? Explain.

09301131

Solution:

Yes, 0 is a rational number.

Explanation

A number in the form $\frac{p}{q}$, p where $p, q \in z$ and $q \neq 0$ is a rational number. The number 0 can be written as $\frac{0}{1}, \frac{0}{2}, \frac{0}{3}, \ldots\ldots$ Here $0 \in z$ and $1,2,3, \in z$ so we can say that 0 is a rational number.

Q.5 State trichotomy property of real numbers. 09301132

Solution:

For all values of $a, b \in R$

Either $a > b$ or $a = b$ or $a < b$

This property is called trichotomy property.

Q.6 Find two rational numbers between 4 and 5. 09301133

Solution:

Average of 4 and 5 $= \frac{4+5}{2} = \frac{9}{2}$

Now we find,

Average of $\frac{9}{2}$ and 5 $= \left(\frac{9}{2}+5\right)\div 2$

$$= \left(\frac{9+10}{2}\right)\times\frac{1}{2}$$

$$= \frac{19}{2}\times\frac{1}{2}$$

$$= \frac{19}{4}$$

Thus two rational number between 4 and 5 are $\frac{9}{2}$ and $\frac{19}{4}$.

Q.7 Simplify the following:

(i) $\sqrt[5]{\frac{x^{15}y^{35}}{z^{20}}}$ 09301134

Solution:

$$\sqrt[5]{\frac{x^{15}y^{35}}{z^{20}}} = \left(\frac{x^{15}y^{35}}{z^{20}}\right)^{\frac{1}{5}}$$

$$= \frac{x^{15\times\frac{1}{5}}y^{35\times\frac{1}{5}}}{z^{20\times\frac{1}{5}}}$$

$$= \frac{x^3.y^7}{z^4}$$

(ii) $\sqrt[3]{(27)^{2x}}$ 09301135

Solution:

$$\sqrt[3]{(27)^{2x}}$$

$$= \sqrt[3]{(3^3)^{2x}}$$

www.ilmkidunya.com

$= \sqrt[3]{(3^{2x})^3}$

$= 3^{2x} \quad \because \sqrt[3]{a^3} = a$

(iii) $\dfrac{6(3)^{n+2}}{3^{n+1}-3^n}$ 09301136

Solution:

$$\frac{6(3)^{n+2}}{3^{n+1}-3^n}$$

$$= \frac{6\left(3^n \times 3^2\right)}{3^n 3^1 - 3^n \times 1} \qquad \left(\because a^{m+n} = a^m \times a^n\right)$$

$$= \frac{3^n \times 6 \times 3^2}{3^n\left[3^1 - 1\right]}$$

$$= \frac{3^n \times 3^{-n} \times 6 \times 9}{2}$$

$$= \frac{3^{n-n} \times 6 \times 9}{2}$$

$$= \frac{3^0 \times 54}{2} \qquad (\because 3^0 = 1)$$

$= 1 \times 27$

$= 27$

Q.8 The sum of three consecutive odd integers is 51. Find the three integers. 09301137

Solution:

Sum = 51

Let x, $x+2$, $x+4$ be three consecutive odd numbers.

By condition:

$(x) + (x+2) + (x+4) = 51$

$x + x + 2 + x + 4 = 51$

$3x + 6 = 51$

$3x = 51 - 6$

$3x = 45$

$x = \dfrac{45}{3}$

$x = 15$

1st odd number $= x = 15$

2nd odd number $= x + 2$

$= 15 + 2 = 17$

3rd odd number $= x + 4$

$= 15 + 4 = 19$

Thus, 15, 17 and 19 are required three consecutive odd numbers.

Q.9 Abdullah picked up 96 balls and placed them into two buckets. One bucket has twenty-eight more balls than the other bucket. How many balls were in each bucket? 09301138

Solution

Total balls = 96

Let balls in 1st and 2nd bucket be x and y respectively.

By 1st condition:

$x + y = 96$ ---------- (i)

By 2nd condition:

$x = 28 + y$

$x - y = 28$ ..... (ii)

Adding eq.(i) and (ii)

$$x + y = 96$$
$$x - y = 28$$
$$2x = 124$$

$$x = \frac{124}{2} = 62$$

$$\boxed{x = 62}$$

Put it in eq. (i)

$62 + y = 96$

$y = 96 - 62$

$$\boxed{y = 34}$$

Thus 1st bucket has 62 balls and 2nd bucket has 34 balls.

Q.10 Salma invested Rs. 350,000 in a bank, which paid simple profit at a rate

www.ilmkidunya.com

$7\frac{1}{4}\%$ per annum. After 2 years, the rate was increased to 8% per annum. Find the amount she had at the end of 7 years.

09301139

Solution:

Time period = 7 years.

We divide the period of 7 years into two parts 2 years and 5 years.

Finding profit for 2 years

Principal amount = P = Rs.350,000/-

Profit rate = R = $7\frac{1}{4}\%$ or 7.25%

Time = t = 2 years

We know that

Profit = P×T×R

= 350,000×2×7.25%

$= 700{,}000 \times \frac{7.25}{100}$

$= 700{,}000 \times \frac{725}{100 \times 100}$

= 70 × 725

= Rs.50,750

Finding the profit for 5 years

Principal amount = P = Rs.350,000/-

Profit rate = R = 8%

Time period = T = 5 years

We know that

Profit = P×T×R

= 350,000×5×8%

$= 1{,}750{,}0\cancel{0}\cancel{0} \frac{8}{1\cancel{00}}$

= Rs. 140,000/-

Finding the total profit:

Total profit = Rs. (50,750+140,000)

= Rs. 190,750

Finding the total amount:

Total amount at the end of 7 years

= Rs. (350,000 + 190,750)

= Rs. 540,750/-

www.ilmkidunya.com