Exercise 1.1

1. Write the following quadratic equations in the standard form and point out pure quadratic equations.

i. $(x+7)(x-3) = -7$

Solution:

$(x+7)(x-3)=-7$
$x(x-3)+7(x-3)=-7$
$x^2-3x+7x-21=-7$
$x^2+4x-21+7=0$
$x^2+4x-14=0$

The above equation is a quadratic equation.

ii. $\frac{x^2+4}{3}-\frac{x}{7}=1$

Solution:

$\frac{x^2+4}{3}-\frac{x}{7}=1$

Multiply both sides by 21, we get

$21\times\frac{x^2+4}{3}-21\times\frac{x}{7}=1\times21$
$7(x^2+4)-3x=21$
$7x^2+28-3x=21$
$7x^2-3x+28-21=0$
$7x^2-3x+7=0$

The above equation is a quadratic equation.

iii. $\frac{x}{x+1}+\frac{x+1}{x}=6$

Solution:

$\frac{x}{x+1}+\frac{x+1}{x}=6$
$\frac{x^2+(x+1)^2}{x(x+1)}=6$
$x^2+x^2+2x+1=6x(x+1)$
$2x^2+2x+1=6x^2+6x$
$2x^2-6x^2+2x-6x+1=0$
$-4x^2-4x+1=0$
$-(4x^2+4x-1)=0$
$\Rightarrow 4x^2+4x-1=0$

The above equation is a quadratic equation.

iv $\frac{x+4}{x-2}-\frac{x-2}{x}+4=0$

Solution:

$\frac{x+4}{x-2}-\frac{x-2}{x}+4=0$
$\frac{x(x+4)-(x-2)^2+4x(x-2)}{x(x-2)}=0$
$(x^2+4x)-(x^2-4x+4)+4(x^2-2x)=0$
$x^2+4x-x^2+4x-4+4x^2-8x=0$
$x^2-x^2+4x^2+4x+4x-8x-4=0$
$4x^2+8x-8x-4=0$
$4x^2-4=0$
$4(x^2-4)=0$
$x^2-1=0$

The above equation is a pure quadratic equation.

v. $\frac{x+3}{x+4}-\frac{x-5}{x}=1$

Solution:

$\frac{x+3}{x+4}-\frac{x-5}{x}=1$
$\frac{x(x+3)-(x+4)(x-5)}{x(x+4)}=1$
$(x^2+3x)-x(x-5)-4(x-5)=x(x+4)$
$x^2+3x-x^2+5x-4x+20=x^2+4x$
$x^2-x^2+3x+5x-4x+20=x^2+4x$
$4x+20=x^2+4x$
$-x^2+4x-4x+20=0$
$-x^2-20=0$
$-(x^2-20)=0$
$x^2-20=0$

The above equation is a pure quadratic equation.

vi. $\frac{x+1}{x+2}+\frac{x+2}{x+3}=\frac{25}{12}$

Solution:

$\frac{x+1}{x+2}+\frac{x+2}{x+3}=\frac{25}{12}$
$\frac{(x+1)(x+3)+(x+2)^2}{(x+2)(x+3)}=\frac{25}{12}$
$\frac{x(x+3)+1(x+3)+(x^2+4x+4)}{x(x+3)+2(x+3)}=\frac{25}{12}$
$\frac{x^2+3x+x+3+x^2+4x+4}{x^2+3x+2x+6}=\frac{25}{12}$

$\frac{2x^2+8x+7}{x^2+5x+6}=\frac{25}{12}$
$25(x^2+5x+6)=12(2x^2+8x+7)$
$25x^2+125x+150=24x^2+96x+84$
$x^2+29x+66=0$

The above equation is a quadratic equation.

Q2. Solve by factorization:

i. $x^2-x-20=0$

Solution:

$x^2-x-20=0$
$x^2-5x+4x-20=0$
$x(x-5)+4(x-5)=0$
$(x+4)(x-5)=0$

Either $x+4=0$ or $x-5=0$
$x=-4$ $x=5$

Thus, solution set $=\{-4,5\}$

ii. $3y^2=y(y-5)$

Solution:

$3y^2=y(y-5)$
$3y^2=y^2-5y$
$3y^2-y^2+5y=0$
$2y^2+5y=0$
$y(2y+5)=0$
Either $y=0$ or $2y+5=0$

$2y=-5$
$y=\frac{-5}{2}$

Thus, solution set $=\left\{0,-\frac{5}{2}\right\}$

iii. $4-32x=17x^2$

Solution:

$4-32x=17x^2$
or $17x^2+32x-4=0$
$17x^2+34x-2x-4=0$
$17x(x+2)-2(x+2)=0$
$(17x-2)(x+2)=0$

Either $17x-2=0$ or $x+2=0$
$17x=2$ $x=-2$
$x=\frac{2}{17}$

Thus, solution set $=\left\{\frac{2}{17},-2\right\}$

iv. $x^2-11x=152$

Solution:

$x^2-11x=152$
$x^2-11x-152=0$
$x^2-19x+8x-152=0$
$x(x-19)+8(x-19)=0$
$(x+8)(x-19)=0$

Either $(x+8)=0$ or $(x-19)=0$
$x=-8$ $x=19$

Thus, solution set = {-8,19}

v. $\frac{x+1}{x}+\frac{x}{x+1}=\frac{25}{12}$

Solution:

$\frac{x+1}{x}+\frac{x}{x+1}=\frac{25}{12}$
$\frac{(x+1)^2+x^2}{x(x+1)}=\frac{25}{12}$
$\frac{x^2+2x+1+x^2}{x^2+x}=\frac{25}{12}$
$\frac{2x^2+2x+1}{x^2+x}=\frac{25}{12}$
$25(x^2+x)=12(2x^2+2x+1)$
$25x^2+25x=24x^2+24x+12$
$25x^2-24x^2+25x-24x-12=0$
$x^2+x-12=0$
$x^2+4x-3x-12=0$
$x(x+4)-3(x+4)=0$
$(x-3)(x+4)=0$

Either $x-3=0$ or $x+4=0$
$x=3$ $x=-4$

Thus, solution set $=\{3,-4\}$

ClassNotes, Inc. © 2020

vi. $\frac{2}{x-9}=\frac{1}{x-3}-\frac{1}{x-4}$

Solution:

$\frac{2}{x-9}=\frac{1}{x-3}-\frac{1}{x-4}$
$\frac{2}{x-9}=\frac{(x-4)-(x-3)}{(x-3)(x-4)}$
$\frac{2}{x-9}=\frac{x-4-x+3}{x^2-7x+12}$
$\frac{2}{x-9}=\frac{-1}{x^2-7x+12}$
$2(x^2-7x+12)=-1(x-9)$
$2x^2-14x+24=-x+9$
$2x^2-14x+x+24-9=0$
$2x^2-13x+15=0$
$2x^2-10x-3x+15=0$
$2x(x-5)-3(x-5)=0$
$(2x-3)(x-5)=0$

Either $2x-3=0$ or $x-5=0$
$2x=3$ $x=5$
$x=\frac{3}{2}$ $x=5$

Thus solution set $=\{5,\frac{3}{2}\}$

Q3. Solve the following equations by completing square:

i. $7x^2+2x-1=0$

Solution:

$7x^2+2x-1=0$
$7x^2+2x=1$
$\frac{7x^2}{7}+\frac{2x}{7}=\frac{1}{7}$
$x^2+\frac{2x}{7}=\frac{1}{7}$
$(x)^2+2(x)\left(\frac{1}{7}\right)+\left(\frac{1}{7}\right)^2=\frac{1}{7}+\left(\frac{1}{7}\right)^2$
$\left(x+\frac{1}{7}\right)^2=\frac{1}{7}+\frac{1}{49}$
$\left(x+\frac{1}{7}\right)^2=\frac{8}{49}$

Takin square root on both sides, we get

$x+\frac{1}{7}=\pm\sqrt{\frac{8}{49}}$
$x+\frac{1}{7}=\pm\frac{2\sqrt{2}}{7}$
$x=-\frac{1}{7}\pm\frac{2\sqrt{2}}{7}$
$x=\frac{-1\pm2\sqrt{2}}{7}$

Thus, solution set $=\left\{\frac{-1\pm2\sqrt{2}}{7}\right\}$

ii. $ax^2+4x-a=0$, $a\neq0$

Solution:

$ax^2+4x=a$

$\frac{ax^2}{a}+\frac{4x}{a}=\frac{a}{a}$
$x^2+\frac{4x}{a}=1$
$(x)^2+2(x)\left(\frac{2}{a}\right)+\left(\frac{2}{a}\right)^2=1+\left(\frac{2}{a}\right)^2$
$\left(x+\frac{2}{a}\right)^2=1+\frac{4}{a^2}$
$\left(x+\frac{2}{a}\right)^2=\frac{a^2+4}{a^2}$

Taking square root on both sides, we get

$x+\frac{2}{a}=\pm\sqrt{\frac{a^2+4}{a^2}}$
$x=-\frac{2}{a}\pm\sqrt{\frac{a^2+4}{a^2}}$
$x=\frac{-2\pm\sqrt{a^2+4}}{a}$

Thus, solution set $=\left\{\frac{-2\pm\sqrt{a^2+4}}{a}\right\}$

iii. $11x^2-34x+3=0$

Solution:

$11x^2-34x+3=0$
$11x^2-34x=-3$
$\frac{11x^2}{11}-\frac{34x}{11}=-\frac{3}{11}$
$x^2-\frac{34}{11}x=-\frac{3}{11}$
$(x)^2-2(x)\left(\frac{34}{22}\right)+\left(\frac{34}{22}\right)^2=-\frac{3}{11}+\left(\frac{34}{22}\right)^2$
$\left(x-\frac{34}{22}\right)^2=-\frac{3}{11}+\frac{1156}{484}$

$\left(x-\frac{34}{22}\right)^2=\frac{-132+1156}{484}$
$\left(x-\frac{34}{22}\right)^2=\frac{1024}{484}$

Taking square root on both sides we get

$\left(x-\frac{34}{22}\right)=\pm\sqrt{\frac{1024}{484}}$
$x-\frac{34}{22}=\pm\frac{32}{22}$
$x=\frac{34}{22}\pm\frac{32}{22}$
$x=\frac{34\pm32}{22}$
$x=\frac{34+32}{22}$, $x=\frac{34-32}{22}$
$=\frac{66}{22}$ $=\frac{2}{22}$
$=3$ $=\frac{1}{11}$

Thus solution set $\left\{3,\frac{1}{11}\right\}$

iv. $lx^2+mx+n=0$, $l\neq0$

Solution:

$lx^2+mx+n=0$
$lx^2+mx=-n$
$\frac{lx^2}{l}+\frac{mx}{l}=-\frac{n}{l}$
$x^2+\frac{mx}{l}=-\frac{n}{l}$
$(x)^2+2(x)\left(\frac{m}{2l}\right)+\left(\frac{m}{2l}\right)^2=-\frac{n}{l}+\left(\frac{m}{2l}\right)^2$

$\left(x+\frac{m}{2l}\right)^2=-\frac{n}{l}+\frac{m^2}{4l^2}$
$\left(x+\frac{m}{2l}\right)^2=\frac{-4ln+m^2}{4l^2}$
$\left(x+\frac{m}{2l}\right)^2=\frac{m^2-4ln}{4l^2}$

Taking square root on both sides, we get

$\sqrt{\left(x+\frac{m}{2l}\right)^2}=\pm\sqrt{\frac{m^2-4ln}{4l^2}}$
$x+\frac{m}{2l}=\pm\sqrt{\frac{m^2-4ln}{4l^2}}$
$x=-\frac{m}{2l}\pm\sqrt{\frac{m^2-4ln}{4l^2}}$
$x=\frac{-m\pm\sqrt{m^2-4ln}}{2l}$

Thus solution set $=\left\{\frac{-m\pm\sqrt{m^2-4ln}}{2l}\right\}$

v. $3x^2+7x=0$

Solution:

$3x^2+7x=0$
$\frac{3x^2}{3}+\frac{7x}{3}=\frac{0}{3}$
$x^2+\frac{7}{3}x=0$
$(x)^2+2(x)\left(\frac{7}{6}\right)+\left(\frac{7}{6}\right)^2=0+\left(\frac{7}{6}\right)^2$
$\left(x+\frac{7}{6}\right)^2=\left(\frac{7}{6}\right)^2$

Taking square root on both sides, we get

$\sqrt{\left(x+\frac{7}{6}\right)^2}=\pm\sqrt{\left(\frac{7}{6}\right)^2}$
$x+\frac{7}{6}=\pm\frac{7}{6}$
$x=-\frac{7}{6}\pm\frac{7}{6}$
$x=-\frac{7}{6}+\frac{7}{6}$ or $x=-\frac{7}{6}-\frac{7}{6}$
$x=0$ $x=-\frac{14}{6}$
$x=-\frac{7}{3}$

Thus solution set $=\left\{0,-\frac{7}{3}\right\}$

vi. $x^2-2x-195=0$

Solution:

$x^2-2x-195=0$
$x^2-2x=195$
$(x)^2-2(x)(1)+(1)^2=195+(1)^2$
$(x-1)^2=195+1$
$(x-1)^2=196$

Taking square root on both sides

$\sqrt{(x-1)^2}=\pm\sqrt{196}$
$x-1=\pm14$
$x=1\pm14$
$x=1+14$ or $x=1-14$
$=15$ $=-13$

Thus, solution set $=\{-13,15\}$

vii. $-x^2+\frac{15}{2}=\frac{7}{2}x$

Solution:

$-x^2-\frac{7}{2}x=-\frac{15}{2}$
$-\left(x^2+\frac{7}{2}x\right)=-\frac{15}{2}$
$x^2+\frac{7}{2}x=\frac{15}{2}$
$(x)^2+2(x)\left(\frac{7}{4}\right)+\left(\frac{7}{4}\right)^2=\frac{15}{2}+\left(\frac{7}{4}\right)^2$
$\left(x+\frac{7}{4}\right)^2=\frac{15}{2}+\frac{49}{16}$
$\left(x+\frac{7}{4}\right)^2=\frac{120+49}{16}$
$\left(x+\frac{7}{4}\right)^2=\frac{169}{16}$

Taking square root on both sides, we get

$\sqrt{\left(x+\frac{7}{4}\right)^2}=\pm\sqrt{\frac{169}{16}}$
$x+\frac{7}{4}=\pm\frac{13}{4}$
$x=-\frac{7}{4}\pm\frac{13}{4}$
$x=-\frac{7}{4}+\frac{13}{4}$ or $x=-\frac{7}{4}-\frac{13}{4}$
$x=\frac{3}{2}$ $x=-5$

Thus, solution set $=\left\{\frac{3}{2},-5\right\}$

viii. $x^2+17x+\frac{33}{4}=0$

Solution:

$x^2+17x=-\frac{33}{4}$
$(x)^2+2(x)\left(\frac{17}{2}\right)+\left(\frac{17}{2}\right)^2=-\frac{33}{4}+\left(\frac{17}{2}\right)^2$
$\left(x+\frac{17}{2}\right)^2=-\frac{33}{4}+\frac{289}{4}$
$\left(x+\frac{17}{2}\right)^2=\frac{256}{4}$

Taking square root on both sides,

$\sqrt{\left(x+\frac{17}{2}\right)^2}=\pm\sqrt{\frac{256}{4}}$
$x+\frac{17}{2}=\pm\frac{16}{2}$
$x=-\frac{17}{2}\pm\frac{16}{2}$
$x=-\frac{17}{2}+\frac{16}{2}$ or $x=-\frac{17}{2}-\frac{16}{2}$
$x=-\frac{1}{2}$ $x=-\frac{33}{2}$

Thus solution set $=\left\{-\frac{1}{2},-\frac{33}{2}\right\}$

ix. $4-\frac{8}{3x+1}=\frac{3x^2+5}{3x+1}$

Solution:

$4-\frac{8}{3x+1}=\frac{3x^2+5}{3x+1}$
$\frac{4(3x+1)-8}{3x+1}=\frac{3x^2+5}{3x+1}$
$\frac{12x+4-8}{3x+1}=\frac{3x^2+5}{3x+1}$
$\frac{12x-4}{3x+1}=\frac{3x^2+5}{3x+1}$

Multiplying both sides by $(3x+1)$, we get

$12x-4=3x^2+5$
$3x^2+5-12x+4=0$
$3x^2-12x+9=0$
$3(x^2-4x+3)=0$
$\Rightarrow x^2-4x+3=0$
$x^2-4x=-3$
$(x)^2-2(x)(2)+(2)^2=-3+(2)^2$
$(x-2)^2=1$

Taking square root on both sides,

$\sqrt{(x-2)^2}=\pm\sqrt{1}$
$x-2=\pm1$
$x=2\pm1$
$x=2+1$ or $x=2-1$
$x=3$ $x=1$

Thus solution set $=\{1,3\}$

x. $7(x+2a)^2+3a^2=5a(7x+23a)$

Solution:

$7(x+2a)^2+3a^2=5a(7x+23a)$
$7(x^2+4ax+4a^2)+3a^2=35ax+115a^2$
$7x^2+28ax+28a^2+3a^2=35ax+115a^2$
$7x^2+28ax-35ax+28a^2+3a^2-115a^2=0$
$7x^2-7ax-84a^2=0$
$7(x^2-ax-12a^2)=0$
$x^2-ax-12a^2=0$
$x^2-ax=12a^2$
$(x)^2-2(x)\left(\frac{a}{2}\right)+\left(\frac{a}{2}\right)^2=12a^2+\left(\frac{a}{2}\right)^2$
$\left(x-\frac{a}{2}\right)^2=12a^2+\frac{a^2}{4}$
$\left(x-\frac{a}{2}\right)^2=\frac{48a^2+a^2}{4}$
$\left(x-\frac{a}{2}\right)^2=\frac{49a^2}{4}$

Taking square root on both sides, we get

$\sqrt{\left(x-\frac{a}{2}\right)^2}=\pm\sqrt{\frac{49a^2}{4}}$
$x-\frac{a}{2}=\pm\frac{7a}{2}$
$x=\frac{a}{2}\pm\frac{7a}{2}$
$x=\frac{a}{2}+\frac{7a}{2}$ or $x=\frac{a}{2}-\frac{7a}{2}$
$=\frac{8a}{2}$ $=-\frac{6a}{2}$
$=4a$ $=-3a$

Thus solution set $=\{-3a,4a\}$

Quadratic Formula:

Derivation of quadratic formula by using completing square method.

The quadratic equation in standard form is

$ax^2+bx+c=0, a\neq 0$

Dividing each term of the equation by a, we get

$x^2+\frac{b}{a}x+\frac{c}{a}=0$

Shifting constant term $\frac{c}{a}$ to the right, we have

$x^2+\frac{b}{a}x=-\frac{c}{a}$

adding $\left(\frac{1}{2}\right)^2$ of both sides, we obtain

$x^2+\frac{b}{a}x+\left(\frac{1}{2}\right)^2\left(\frac{b}{a}\right)^2=-\frac{c}{a}+\left(\frac{1}{2}\right)^2\left(\frac{b}{a}\right)^2$

$\left(x+\frac{1}{2}\frac{b}{a}\right)^2=-\frac{c}{a}+\frac{1}{4}\frac{b^2}{a^2}$

$\left(x+\frac{b}{2a}\right)^2=\frac{b^2-4ac}{4a^2}$

Taking square root on both sides, we get

$\sqrt{\left(x+\frac{b}{2a}\right)^2}=\sqrt{\frac{b^2-4ac}{4a^2}}$

or $x+\frac{b}{2a}=\frac{\pm\sqrt{b^2-4ac}}{2a}$ or $x=-\frac{b}{2a}\pm\frac{\sqrt{b^2-4ac}}{2a}$ or $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

Thus, $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$ is known as "quadratic formula"

Exercise 1.2

Q1. Solve the following equations using quadratic formula:

(i) $\frac{1}{x}-x^2=7x$

Solution

$\frac{1}{x}-x^2-7x=0$

$-x^2-7x+1=0$

$-\left(x^2+7x-1\right)=0$

$x=x^2+7x-1=0$

Comparing it with standard quadratic equation, we have

$ax^2+bx+c=0$

Here $a=1, b=7, c=-1$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-7\pm\sqrt{7^2-4(1)(-1)}}{2(1)}$

$x=\frac{-7\pm\sqrt{49+4}}{2}$

$x=\frac{-7\pm\sqrt{53}}{2}$

Thus, solution are $=\left\{\frac{-7\pm\sqrt{53}}{2}\right\}$

(ii) $5x^2+8x+3=0$

Solution

Comparing it with standard quadratic equation, we have

$ax^2+bx+c=0$

Here $a=5, b=8, c=3$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-8\pm\sqrt{8^2-4(5)(3)}}{2(5)}$

$x=\frac{-8\pm\sqrt{64-60}}{10}$

$x=\frac{-8\pm\sqrt{4}}{10}$

$x=\frac{-8\pm 2}{10}$

Then solution are $=\left\{\frac{-8\pm 2}{10}\right\}$

Here $a=5, b=8, c=3$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-8\pm\sqrt{8^2-4(5)(3)}}{2(5)}$

$x=\frac{-8\pm\sqrt{64-60}}{10}$

$x=\frac{-8\pm\sqrt{4}}{10}$

$x=\frac{-8\pm 2}{10}$

$x=\frac{-8+2}{10}, \frac{-8-2}{10}$

Then solution are $=\left\{\frac{-8+2}{10}, \frac{-8-2}{10}\right\}$

(iii) $\sqrt{2}x^2+x-4\sqrt{2}$

Solution

$\sqrt{2}x^2+x+4\sqrt{2}=0$

$\sqrt{2}x^2+x+4\sqrt{2}=0$

Comparing it with standard quadratic equation, we have

$ax^2+bx+c=0$

Here $a=\sqrt{2}, b=1, c=4\sqrt{2}$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-1\pm\sqrt{1^2-4\left(\sqrt{2}\right)\left(4\sqrt{2}\right)}}{2\left(\sqrt{2}\right)}$

$x=\frac{-1\pm\sqrt{1-32}}{2\sqrt{2}}$

$x=\frac{-1\pm\sqrt{-31}}{2\sqrt{2}}$

$x=\frac{-1\pm\sqrt{31}i}{2\sqrt{2}}$

$x=\frac{-1}{2\sqrt{2}}\pm\frac{\sqrt{31}i}{2\sqrt{2}}$ or $x=\frac{-1}{2\sqrt{2}}\pm\frac{\sqrt{31}i}{2\sqrt{2}}$

$x=\frac{-1}{2\sqrt{2}}\pm\frac{\sqrt{31}i}{2\sqrt{2}}$

$x=\frac{-1}{2\sqrt{2}}\pm\frac{\sqrt{31}i}{2\sqrt{2}}$

$x=\frac{-1\pm\sqrt{31}i}{2\sqrt{2}}$

Thus solution are $=\left\{\frac{-1\pm\sqrt{31}i}{2\sqrt{2}}\right\}$

(iv) $4x^2-10x+3=0$

Solution

$4x^2-10x+3=0$

Comparing it with standard quadratic equation, we have

$ax^2+bx+c=0$

Here $a=4, b=-7, c=3$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-(-7)\pm\sqrt{(-7)^2-4(4)(3)}}{2(4)}$

$x=\frac{7\pm\sqrt{49-48}}{8}$

$x=\frac{7\pm\sqrt{1}}{8}$

$x=\frac{7\pm 1}{8}$ or $x=\frac{7+1}{8}$ or $x=\frac{7-1}{8}$

$x=\frac{8}{8}$ or $x=\frac{6}{8}$

$x=1$ or $x=\frac{3}{4}$

Thus solution are $=\left\{1, \frac{3}{4}\right\}$

Here $a=4, b=-7, c=3$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-(-7)\pm\sqrt{(-7)^2-4(4)(3)}}{2(4)}$

$x=\frac{7\pm\sqrt{49-48}}{8}$

$x=\frac{7\pm\sqrt{1}}{8}$

$x=\frac{7\pm 1}{8}$ or $x=\frac{7+1}{8}$ or $x=\frac{7-1}{8}$

$x=\frac{8}{8}$ or $x=\frac{6}{8}$

$x=1$ or $x=\frac{3}{4}$

Thus solution are $=\left\{1, \frac{3}{4}\right\}$

Thus solution are $=\left\{-\frac{1}{2}, 1\right\}$

(vi) $3x^2+8x+5=0$

Solution

$3x^2+8x+5=0$

Comparing it with standard quadratic equation, we have

$ax^2+bx+c=0$

Here $a=3, b=8, c=5$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-8\pm\sqrt{8^2-4(3)(5)}}{2(3)}$

$x=\frac{-8\pm\sqrt{64-60}}{6}$

$x=\frac{-8\pm\sqrt{4}}{6}$

$x=\frac{-8\pm 2}{6}$

$x=\frac{-8+2}{6}$ or $x=\frac{-8-2}{6}$

$x=\frac{-6}{6}$ or $x=\frac{-10}{6}$

$x=-1$ or $x=-\frac{5}{3}$

Thus solution are $=\left\{-1, -\frac{5}{3}\right\}$

(vii) $\frac{1}{x}-\frac{1}{x-2}=\frac{3}{4}, x\neq 0$

Solution

$\frac{1}{x}-\frac{1}{x-2}=\frac{3}{4}, x\neq 0$

$\frac{(x-2)-(x)}{x(x-2)}=\frac{3}{4}, x\neq 0$

$\frac{x-2-x}{x^2-2x}=\frac{3}{4}, x\neq 0$

$\frac{-2}{x^2-2x}=\frac{3}{4}, x\neq 0$

$-8=3x^2-6x, x\neq 0$

$3x^2-6x+8=0, x\neq 0$

Comparing it with standard quadratic equation, we have

$ax^2+bx+c=0$

Here $a=3, b=-6, c=8$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-(-6)\pm\sqrt{(-6)^2-4(3)(8)}}{2(3)}$

$x=\frac{6\pm\sqrt{36-96}}{6}$

$x=\frac{6\pm\sqrt{-60}}{6}$

$x=\frac{6\pm 2\sqrt{15}i}{6}$

$x=\frac{6}{6}\pm\frac{2\sqrt{15}i}{6}$ or $x=\frac{6}{6}\pm\frac{2\sqrt{15}i}{6}$

$x=1\pm\frac{\sqrt{15}i}{3}$ $x\neq 0$

Thus, solution are $=\left\{1\pm\frac{\sqrt{15}i}{3}\right\}$

$\frac{\left(\sqrt{2}x^2+4\sqrt{2}\right)\left(\sqrt{2}x^2+4\sqrt{2}\right)}{\left(\sqrt{2}x^2+4\sqrt{2}\right)}=\frac{x}{3}$

$\frac{\left(\sqrt{2}x^2+4\sqrt{2}\right)\left(\sqrt{2}x^2+4\sqrt{2}\right)}{\left(\sqrt{2}x^2+4\sqrt{2}\right)}=\frac{x}{3}$

$\frac{2x^2+8\sqrt{2}x^2+8\sqrt{2}}{\sqrt{2}x^2+4\sqrt{2}}=\frac{x}{3}$

$\frac{3\left(2x^2+8\sqrt{2}x^2+8\sqrt{2}\right)}{\sqrt{2}x^2+4\sqrt{2}}=x$

$\frac{6x^2+24\sqrt{2}x^2+24\sqrt{2}}{\sqrt{2}x^2+4\sqrt{2}}=x$

$6x^2+24\sqrt{2}x^2+24\sqrt{2}=\sqrt{2}x^3+4\sqrt{2}x$

$\sqrt{2}x^3-6x^2-24\sqrt{2}x^2+4\sqrt{2}x-24\sqrt{2}=0$

Comparing it with standard quadratic equation, we have

$ax^3+bx^2+cx+d=0$

Here $a=\sqrt{2}, b=-6, c=4\sqrt{2}-24\sqrt{2}, d=-24\sqrt{2}$

$\sqrt{2}x^3-6x^2-44\sqrt{2}x-24\sqrt{2}=0$

$\sqrt{2}x^3-6x^2-44\sqrt{2}x-24\sqrt{2}=0$

$\sqrt{2}\left(x^3-3\sqrt{2}x^2-44x-24\sqrt{2}\right)=0$

$x^3-3\sqrt{2}x^2-44x-24\sqrt{2}=0$

Comparing it with standard quadratic equation, we have

$ax^3+bx^2+cx+d=0$

Here $a=1, b=-3\sqrt{2}, c=-44, d=-24\sqrt{2}$

$x^3-3\sqrt{2}x^2-44x-24\sqrt{2}=0$

$x^3-3\sqrt{2}x^2-44x-24\sqrt{2}=0$

$x^3-3\sqrt{2}x^2-44x-24\sqrt{2}=0$

$x^3-3\sqrt{2}x^2-44x-24\sqrt{2}=0$

(ix) $\frac{x}{x-1}+\frac{x-1}{x}=2$

Solution

$\frac{x}{x-1}+\frac{x-1}{x}=2$

$\frac{x(x)+(x-1)(x-1)}{(x-1)(x)}=2$

$\frac{x^2+x^2-2x+1}{x^2-x}=2$

$2x^2-2x+1=2\left(x^2-x\right)$

$2x^2-2x+1=2x^2-2x$

$2x^2-2x^2-2x+2x+1=0$

$1=0$

Comparing it with standard quadratic equation, we have

$ax^2+bx+c=0$

Here $a=1, b=-1, c=-12$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-(-1)\pm\sqrt{(-1)^2-4(1)(-12)}}{2(1)}$

$x=\frac{1\pm\sqrt{1+48}}{2}$

$x=\frac{1\pm\sqrt{49}}{2}$

$x=\frac{1\pm 7}{2}$ or $x=\frac{1-7}{2}$ or $x=\frac{1+7}{2}$

$x=\frac{8}{2}$ or $x=\frac{-6}{2}$

$x=4$ or $x=-3$

Thus, solution are $=\{-3, 4\}$

(x) $\frac{x}{x-1}+\frac{x-1}{x}=2$

Solution

$\frac{x}{x-1}+\frac{x-1}{x}=2$

$\frac{x(x)+(x-1)(x-1)}{(x-1)(x)}=2$

$\frac{x^2+x^2-2x+1}{x^2-x}=2$

$2x^2-2x+1=2\left(x^2-x\right)$

$2x^2-2x+1=2x^2-2x$

$2x^2-2x^2-2x+2x+1=0$

$1=0$

Comparing it with standard quadratic equation, we have

$ax^2+bx+c=0$

Here $a=1, b=-1, c=-12$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-(-1)\pm\sqrt{(-1)^2-4(1)(-12)}}{2(1)}$

$x=\frac{1\pm\sqrt{1+48}}{2}$

$x=\frac{1\pm\sqrt{49}}{2}$

$x=\frac{1\pm 7}{2}$ or $x=\frac{1-7}{2}$ or $x=\frac{1+7}{2}$

$x=\frac{8}{2}$ or $x=\frac{-6}{2}$

$x=4$ or $x=-3$

Thus, solution are $=\{-3, 4\}$

$x=\frac{8\pm\sqrt{96}}{4}$

$x=\frac{8\pm 4\sqrt{6}}{4}$

$x=\frac{8}{4}\pm\frac{4\sqrt{6}}{4}$

$x=2\pm\sqrt{6}$

Thus, solution are $=\left\{2\pm\sqrt{6}\right\}$

(xi) $x^2+2x+1=0$

Solution

$x^2+2x+1=0$

Comparing it with standard quadratic equation, we have

$ax^2+bx+c=0$

Here $a=1, b=2, c=1$

Now $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-2\pm\sqrt{2^2-4(1)(1)}}{2(1)}$

$x=\frac{-2\pm\sqrt{4-4}}{2}$

$x=\frac{-2\pm\sqrt{0}}{2}$

$x=\frac{-2\pm 0}{2}$ or $x=\frac{-2+0}{2}$ or $x=\frac{-2-0}{2}$

$x=\frac{-2}{2}$ or $x=\frac{-2}{2}$

$x=-1$ or $x=-1$

Thus solution are $=\{-1, -1\}$

Exercise 3.3

Q1 - Solve the following equations:

(i) $\frac{3x}{2} + 5 = \frac{2x}{3} - 13 = 0$

Solution:

$\frac{3x}{2} + 5 = \frac{2x}{3} - 13$ ..... (1)

To find Solution:

$\frac{3x}{2} - \frac{2x}{3} = -13 - 5$

$\frac{9x - 4x}{6} = -18$

$\frac{5x}{6} = -18$

$5x = -18 \times 6$

$x = \frac{-108}{5}$

Thus, solution set $= \left\{\frac{-108}{5}\right\}$

(ii) $\frac{2x}{3} + \frac{x}{2} - 4 = 0$

Solution:

$\frac{2x}{3} + \frac{x}{2} - 4 = 0$ ..... (1)

To find Solution:

$\frac{2x}{3} + \frac{x}{2} = 4$

$\frac{4x + 3x}{6} = 4$

$\frac{7x}{6} = 4$

$7x = 4 \times 6$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(iii) $\frac{3x}{4} + \frac{2x}{3} - 3 = 0$

Solution:

$\frac{3x}{4} + \frac{2x}{3} - 3 = 0$ ..... (1)

To find Solution:

$\frac{3x}{4} + \frac{2x}{3} = 3$

$\frac{9x + 8x}{12} = 3$

$\frac{17x}{12} = 3$

$17x = 3 \times 12$

$x = \frac{36}{17}$

Thus, solution set $= \left\{\frac{36}{17}\right\}$

(iv) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(v) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(vi) $\frac{x}{2} + \frac{2x}{3} = \frac{3x}{4} + 1$

Solution:

$\frac{x}{2} + \frac{2x}{3} = \frac{3x}{4} + 1$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{2x}{3} - \frac{3x}{4} = 1$

$\frac{6x + 8x - 9x}{12} = 1$

$\frac{5x}{12} = 1$

$5x = 1 \times 12$

$x = \frac{12}{5}$

Thus, solution set $= \left\{\frac{12}{5}\right\}$

(vii) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(viii) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(ix) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(x) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xi) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xii) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xiii) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xiv) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xv) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xvi) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xvii) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xviii) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xix) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xx) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xxi) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xxii) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xxiii) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xxiv) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xxv) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xxvi) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xxvii) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xxviii) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xxix) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

(xxx) $\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$

Solution:

$\frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 2$ ..... (1)

To find Solution:

$\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = 2$

$\frac{6x + 4x - 3x}{12} = 2$

$\frac{7x}{12} = 2$

$7x = 2 \times 12$

$x = \frac{24}{7}$

Thus, solution set $= \left\{\frac{24}{7}\right\}$

Radical Equations
To solve radical equations, square / power the radical sign to undo or isolate a radical.

$$\sqrt{x}=\sqrt{x+1} \quad \text{or} \quad \sqrt[3]{x}=\sqrt[3]{x+1}$$

Exercise 1.4

Solve the following equations:

(1) $3x+5=\sqrt{9x+16}$

Solution

Squaring both sides, we get

$$(3x+5)^2=\left(\sqrt{9x+16}\right)^2$$

$$9x^2+30x+25=9x+16$$

$$9x^2+30x-9x+25-16=0$$

$$9x^2+21x+9=0$$

$$3x^2+7x+3=0$$

$$a=3,\ b=7,\ c=3$$

$$x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$$

$$x=\frac{-7\pm\sqrt{7^2-4(3)(3)}}{2(3)}$$

$$x=\frac{-7\pm\sqrt{49-36}}{6}$$

$$x=\frac{-7\pm\sqrt{13}}{6}$$

Thus, solution set $=\left\{\frac{-7+\sqrt{13}}{6},\frac{-7-\sqrt{13}}{6}\right\}$ Ans

(2) $\sqrt{x+1}=\sqrt{3x-1}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{3x-1}\right)^2$$

$$x+1=3x-1$$

$$x-3x=-1-1$$

$$-2x=-2$$

$$x=1$$

Thus, solution set $=\{1\}$ Ans

(3) $\sqrt{x+1}=\sqrt{2x-3}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{2x-3}\right)^2$$

$$x+1=2x-3$$

$$x-2x=-3-1$$

$$-x=-4$$

$$x=4$$

Thus, solution set $=\{4\}$ Ans

(4) $\sqrt{2x+1}=\sqrt{3x-2}$

Solution

Squaring both sides, we get

$$\left(\sqrt{2x+1}\right)^2=\left(\sqrt{3x-2}\right)^2$$

$$2x+1=3x-2$$

$$2x-3x=-2-1$$

$$-x=-3$$

$$x=3$$

Thus, solution set $=\{3\}$ Ans

(5) $\sqrt{x+1}=\sqrt{x}+1$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+1\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}(1)+1^2$$

$$x+1=x+2\sqrt{x}+1$$

$$x-x-2\sqrt{x}=1-1$$

$$-2\sqrt{x}=0$$

$$\sqrt{x}=0$$

$$x=0$$

Thus, solution set $=\{0\}$ Ans

(6) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-1}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-1}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-1}+\left(\sqrt{x-1}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-1}+x-1$$

$$x-x-x=-1-1$$

$$-x=-2$$

$$x=2$$

Thus, solution set $=\{2\}$ Ans

(7) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-2}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-2}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-2}+\left(\sqrt{x-2}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-2}+x-2$$

$$x-x-x=-2-1$$

$$-x=-3$$

$$x=3$$

Thus, solution set $=\{3\}$ Ans

(8) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-3}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-3}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-3}+\left(\sqrt{x-3}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-3}+x-3$$

$$x-x-x=-3-1$$

$$-x=-4$$

$$x=4$$

Thus, solution set $=\{4\}$ Ans

(9) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-4}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-4}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-4}+\left(\sqrt{x-4}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-4}+x-4$$

$$x-x-x=-4-1$$

$$-x=-5$$

$$x=5$$

Thus, solution set $=\{5\}$ Ans

(10) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-5}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-5}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-5}+\left(\sqrt{x-5}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-5}+x-5$$

$$x-x-x=-5-1$$

$$-x=-6$$

$$x=6$$

Thus, solution set $=\{6\}$ Ans

(11) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-6}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-6}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-6}+\left(\sqrt{x-6}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-6}+x-6$$

$$x-x-x=-6-1$$

$$-x=-7$$

$$x=7$$

Thus, solution set $=\{7\}$ Ans

(12) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-7}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-7}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-7}+\left(\sqrt{x-7}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-7}+x-7$$

$$x-x-x=-7-1$$

$$-x=-8$$

$$x=8$$

Thus, solution set $=\{8\}$ Ans

(13) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-8}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-8}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-8}+\left(\sqrt{x-8}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-8}+x-8$$

$$x-x-x=-8-1$$

$$-x=-9$$

$$x=9$$

Thus, solution set $=\{9\}$ Ans

(14) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-9}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-9}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-9}+\left(\sqrt{x-9}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-9}+x-9$$

$$x-x-x=-9-1$$

$$-x=-10$$

$$x=10$$

Thus, solution set $=\{10\}$ Ans

(15) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-10}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-10}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-10}+\left(\sqrt{x-10}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-10}+x-10$$

$$x-x-x=-10-1$$

$$-x=-11$$

$$x=11$$

Thus, solution set $=\{11\}$ Ans

(16) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-11}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-11}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-11}+\left(\sqrt{x-11}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-11}+x-11$$

$$x-x-x=-11-1$$

$$-x=-12$$

$$x=12$$

Thus, solution set $=\{12\}$ Ans

(17) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-12}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-12}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-12}+\left(\sqrt{x-12}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-12}+x-12$$

$$x-x-x=-12-1$$

$$-x=-13$$

$$x=13$$

Thus, solution set $=\{13\}$ Ans

(18) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-13}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-13}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-13}+\left(\sqrt{x-13}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-13}+x-13$$

$$x-x-x=-13-1$$

$$-x=-14$$

$$x=14$$

Thus, solution set $=\{14\}$ Ans

(19) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-14}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-14}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-14}+\left(\sqrt{x-14}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-14}+x-14$$

$$x-x-x=-14-1$$

$$-x=-15$$

$$x=15$$

Thus, solution set $=\{15\}$ Ans

(20) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-15}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-15}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-15}+\left(\sqrt{x-15}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-15}+x-15$$

$$x-x-x=-15-1$$

$$-x=-16$$

$$x=16$$

Thus, solution set $=\{16\}$ Ans

(21) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-16}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-16}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-16}+\left(\sqrt{x-16}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-16}+x-16$$

$$x-x-x=-16-1$$

$$-x=-17$$

$$x=17$$

Thus, solution set $=\{17\}$ Ans

(22) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-17}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-17}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-17}+\left(\sqrt{x-17}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-17}+x-17$$

$$x-x-x=-17-1$$

$$-x=-18$$

$$x=18$$

Thus, solution set $=\{18\}$ Ans

(23) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-18}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-18}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-18}+\left(\sqrt{x-18}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-18}+x-18$$

$$x-x-x=-18-1$$

$$-x=-19$$

$$x=19$$

Thus, solution set $=\{19\}$ Ans

(24) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-19}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-19}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-19}+\left(\sqrt{x-19}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-19}+x-19$$

$$x-x-x=-19-1$$

$$-x=-20$$

$$x=20$$

Thus, solution set $=\{20\}$ Ans

(25) $\sqrt{x+1}=\sqrt{x}+\sqrt{x-20}$

Solution

Squaring both sides, we get

$$\left(\sqrt{x+1}\right)^2=\left(\sqrt{x}+\sqrt{x-20}\right)^2$$

$$x+1=\left(\sqrt{x}\right)^2+2\sqrt{x}\sqrt{x-20}+\left(\sqrt{x-20}\right)^2$$

$$x+1=x+2\sqrt{x}\sqrt{x-20}+x-20$$

$$x-x-x=-20-1$$

$$-x=-21$$

$$x=21$$

Thus, solution set $=\{21\}$ Ans

Solved Miscellaneous Exercise. 1

Q1. Multiple Choice Questions:

Four possible answers are given for the following questions. Tick (✓) the correct answer.

(i) Standard form of quadratic equation is:

(a) $ax+c=0, a\neq 0$ **(b)** $ax^2+bx+c=0, a\neq 0$

(c) $ax^2=0, a\neq 0$ **(d)** $ax^2=0, a\neq 0$

(ii) The number of terms in a standard quadratic equation $ax^2+bx+c=0$ **is**

(a) 1 **(b)** 2 **(c)** 3 **(d)** 4

(iii) The number of methods to solve a quadratic equation is:

(a) 1 **(b)** 2 **(c)** 3 **(d)** 4

(iv) The quadratic formula is:

(a) $\frac{-b\pm\sqrt{b^2-4ac}}{2a}$ **(b)** $\frac{b\pm\sqrt{b^2-4ac}}{2a}$

(c) $\frac{-b\pm\sqrt{b^2+4ac}}{2a}$ **(d)** $\frac{b\pm\sqrt{b^2-4ac}}{2a}$

(v) Two linear factors of $x^2-15x+56$ **are:**

(a) $(x-7)$ and $(x+8)$ **(b)** $(x+7)$ and $(x-8)$

(c) $(x-7)$ and $(x-8)$ **(d)** $(x+7)$ and $(x+8)$

(vi) An equation which remains unchanged when x is replaced by $\frac{1}{x}$ **is called a/an**

(a) Exponential equation **(b)** Reciprocal equation

(c) Radical Equation **(d)** None of these

(vii) An equation of the type 3x+3=+4x+0 is a/an:

(a) Exponential equation **(b)** Radical Equation

(c) Reciprocal equation **(d)** None of these

(viii) The solution set of equation $4x^2-16=0$ **is**

(a) $\{14\}$ **(b)** $\{4\}$

(c) $\{\pm 2\}$ **(d)** ± 2

(ix) An equation of the form $2x^4-3x^3+7x^2-3x+2=0$ **is called a/an**

(a) Reciprocal equation **(b)** Radical Equation

(c) Exponential equation **(d)** None of these

Answers:

(i)	b	(ii)	c	(iii)	c	(iv)	a	(v)	c

(vi)	b	(vii)	b	(viii)	c	(ix)	a		

Q2. Write short answers of the following question.

i. $x^2+2x-2=0$

Answer:

$x^2+2x-2=0$

Compare it with

$ax^2+bx+c=0$

Here $a=1, b=2, c=-2$

By using quadratic equation, we get

$x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$

$x=\frac{-(2)\pm\sqrt{(2)^2-4(1)(-2)}}{2(1)}$

$x=\frac{-2\pm\sqrt{4+8}}{2}$

$x=\frac{-2\pm\sqrt{12}}{2}$

$x=\frac{-2\pm 2\sqrt{3}}{2}$

$x=\frac{2(-1\pm\sqrt{3})}{2}=-1\pm\sqrt{3}$

Thus, the solution set $=\{-1\pm\sqrt{3}\}$

ii. Solve by factorization $5x^2=15x$

Answer:

$5x^2=15x$

$5x^2-15x=0$

$5x(x-3)=0$

Either $5x=0$ *or* $x-3=0$

$x=0$ $x=3$

Thus, the solution set $=\{0,3\}$

iii. Write in standard form $\frac{1}{x+4}-\frac{1}{x-4}=\frac{3}{2}$

Solution:

$\frac{1}{x+4}-\frac{1}{x-4}=3$ $\frac{x-4-x-4}{(x+4)(x-4)}=3$

$\frac{2x}{x^2-16}=3$ $3(x^2-16)=2x$

$3x^2-48-2x=0$

$3x^2-2x-48=0$

iv. Write the names of the method for solving a quadratic equation.

Answer: Solution of quadratic equations:

To find solution of a quadratic equation, following methods are used:

(i) Factorization **(ii)** Completing square

(iii) Quadratic formula

v. $Solve\left(2x-\frac{1}{2}\right)^2=\frac{9}{4}$

Answer:

$\left(2x-\frac{1}{2}\right)^2=\frac{9}{4}$

Taking square root on both sides, we get

$2x-\frac{1}{2}=\pm\frac{3}{2}$

$2x=\frac{1}{2}\pm\frac{3}{2}$

Either $2x=\frac{1}{2}+\frac{3}{2}$ *or* $2x=\frac{1}{2}-\frac{3}{2}$

$2x=2$ $2x=-1$

$x=1$ $x=-\frac{1}{2}$

Thus, the solution set $=\left\{-\frac{1}{2},1\right\}$

vi. $Solve\sqrt{3x+18}=x$

Answer:

$\sqrt{3x+18}=x$

Taking square root on both sides, we get

$3x+18=x^2$

$x^2-3x-18=0$

$x^2-6x+3x-18=0$

$x(x-6)+3(x-6)=0$

$(x-6)(x+3)=0$

Either $x-6=0$ *or* $x+3=0$

$x=6$ $x=-3$

Thus, the solution set $=\{-3,6\}$

vii. Define quadratic equation.

Ans: Quadratic Equation:

An equation, which contains the square of the unknown (variable) quantity, but no higher power, is called a **quadratic equation** or an equation of the second degree.

viii. Define reciprocal equation.

Ans: Reciprocal equations of the type:

$ax\left(x^2+\frac{1}{x^2}\right)+b\left(x+\frac{1}{x}\right)+c=0$ or $ax^3+bx^2+cx^2+bx+ax=0$

An equation is said to be a reciprocal equation, if it remains unchanged, when x is replaced by $\frac{1}{x}$.

Replacing x by $\frac{1}{x}$ as $ax^3-bx^2+cx^2-bx+a=0$

$a\left(\frac{1}{x}\right)^3-b\left(\frac{1}{x}\right)^2+c\left(\frac{1}{x}\right)^2-b\left(\frac{1}{x}\right)+a=0$ which is simplified as

$a-bx+cx^2-bx^2+ax^3=0$ we get the same equation.

Thus $ax^3-bx^2+cx^2-bx+a=0$ is a reciprocal equation.

ix. Define exponential equation.

Ans: Exponential equation:

In exponential equations, variable occurs in exponent.

For example. $5^{x+1}+5x^2=26$.

x. Define radical equation.

Ans: Radical equations:

An equation involving expression under the radical sign is called a **radical equation.**

e.g., $\sqrt{x+3}=x+1$ and $\sqrt{x-1}-\sqrt{x-2}=1$

Q3. Fill in the blanks:

(i) The standard form of quadratic equation is _____.

(ii) The number of methods to solve a quadratic equation are _____.

(iii) The name of the method to derive a quadratic formula is _____.

(iv) The solution of the equation $ax^2+bx+c=0, a\neq 0$ is _____.

(v) The solution set of $25x^2-4=0$ is _____.

(vi) An equation of the form $2^x-3.2^{-x}+5=0$ is called a/an _____ equation.

(vii) The solution set of the equation $x^2-9=0$ is _____.

(viii) An equation of the type $x^4+x^3+x^2+x+1=0$ called a/an _____.

(ix) A root of an equation, which do not satisfy the equation is called _____root.

(x) An equation involving impression of the variable under _____is called radical.

Answer

(i)	$ax^2+bx+c=0$	(ii)	3	(iii)	Completing Square
(iv)	$\frac{-b\pm\sqrt{b^2-4ac}}{2a}$	(v)	$\left\{\pm\frac{1}{5}\right\}$	(vi)	Exponential

(vii)	$\{\pm 3\}$	(viii)	Reciprocal	(ix)	Extraneous
(x)	Radical Sign				

Summary

- ✓ An equation, which contains the square of the unknown (variable) quantity, but no higher power, is called a quadratic equation or an equation of the second degree.
- ✓ A second-degree equation in one variable is $ax^2+bx+c=0$
- ✓ Where $a\neq 0$ and a, b, care real numbers, is called the general or standard form of a quadratic equation.
- ✓ An equation is said to be a reciprocal equation, if it remains unchanged, when x is replaced by $\frac{1}{x}$.
- ✓ In exponential equations, variable occur in exponents.
- ✓ An equation involving expression under the radical sign is called a radical equation.
- ✓ Quadratic formula for $ax^2+bx+c=0, a\neq 0$ is $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$
- ✓ Any quadratic equation is solved by the following three methods.
- ✓ **(i)** Factorization **(ii)** Completing square **(iii)**Quadratic formula