Solved Exercise 2.1

1. Find the discriminant of the following given quadratic equation

(i) $2x^2 + 3x + 1 = 0$

Solution:
$a = 2, b = 3, c = 0$
Compare it with
$ax^2 + bx + c = 0$
Discriminant $= b^2 - 4ac = 0$
Also, $\Delta = b^2 - 4ac$
$= (3)^2 - 4(2)(1)$
$= 9 - 8$
$= 1$

(ii) $4x^2 - 4x + 3 = 0$

Solution:
$a = 4, b = -4, c = 3$
Compare it with
$ax^2 + bx + c = 0$
Discriminant $= b^2 - 4ac = 0$
Also, $\Delta = b^2 - 4ac$
$= (-4)^2 - 4(4)(3)$
$= 16 - 48$
$= -32$

(iii) $3x^2 - 30x + 25 = 0$

Solution:
$a = 3, b = -30, c = 25$
Compare it with
$ax^2 + bx + c = 0$
Discriminant $= b^2 - 4ac = 0$
Also, $\Delta = b^2 - 4ac$
$= (-30)^2 - 4(3)(25)$
$= 900 - 300$
$= 600$

(iv) $4x^2 - 7x - 2 = 0$

Solution:
$a = 4, b = -7, c = -2$
Compare it with
$ax^2 + bx + c = 0$
Discriminant $= b^2 - 4ac = 0$
Also, $\Delta = b^2 - 4ac$
$= (-7)^2 - 4(4)(-2)$
$= 49 + 32$
$= 81$

2. Find the nature of the roots of the following given quadratic equations and verify the result by solving the equations

(i) $x^2 - 22x + 120 = 0$

Solution:
$x^2 - 22x + 120 = 0$
Compare it with
$ax^2 + bx + c = 0$
Also $a = 1, b = -22, c = 120$
Also, $\Delta = b^2 - 4ac$

$= (-22)^2 - 4(1)(120)$
$= 484 - 480$
$= 4$
$= (2)^2 > 0$

As the discr. is positive and is a power square. Therefore the roots are rational (real) and unequal. Verification by solving the equation.

$x^2 - 22x + 120 = 0$
$x^2 - 12x - 10x + 120 = 0$
$x(x - 12) - 10(x - 12) = 0$
$(x - 12)(x - 10) = 0$
Either $x - 0 = 0$ or $x - 10 = 0$
$x = 12$ $x = 10$

Thus, the roots are rational (real) and unequal.

(ii) $2x^2 + 3x + 7 = 0$

Solution:
$2x^2 + 3x + 7 = 0$
Compare it with
$ax^2 + bx + c = 0$
Also $a = 2, b = 3, c = 7$
Also, $\Delta = b^2 - 4ac$
$= (3)^2 - 4(2)(7)$
$= 9 - 56$
$= -47$

As the discr. is negative.
Therefore the roots are imaginary and unequal.
Verification by solving the equation.
Let's use formula
Using quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-(3) \pm \sqrt{(3)^2 - 4(2)(7)}}{2(2)}$$

$$= \frac{-3 \pm \sqrt{9 - 56}}{4}$$

$$= \frac{-3 \pm \sqrt{-47}}{4}$$

Thus, the roots are imaginary and unequal.

(iii) $36x^2 - 24x + 9 = 0$

Solution:
$36x^2 - 24x + 9 = 0$
Compare it with
$ax^2 + bx + c = 0$
Also $a = 36, b = -24, c = 9$
Also, $\Delta = b^2 - 4ac$
$= (-24)^2 - 4(36)(9)$
$= 576 - 576$
$= 0$

As the discr. is zero.
Therefore the roots are rational (real) and equal.
Verification by solving the equation.
Let's use formula
Using quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-(-24) \pm \sqrt{(-24)^2 - 4(36)(9)}}{2(36)}$$

$$= \frac{24 \pm \sqrt{576 - 576}}{72}$$

$$= \frac{24 \pm \sqrt{0}}{72}$$

$$= \frac{24}{72}$$

$$= \frac{24}{72}$$
$$= \frac{1}{3}$$

Thus, the roots are real and equal.

(iv) $3x^2 + 5x - 12 = 0$

Solution:
$3x^2 + 5x - 12 = 0$
Compare it with
$ax^2 + bx + c = 0$
Also $a = 3, b = 5, c = -12$
Also, $\Delta = b^2 - 4ac$
$= (5)^2 - 4(3)(-12)$
$= 25 + 144$
$= 169$
$= (13)^2 > 0$

As the discr. is positive and not a perfect square.
Therefore the roots are irrational (real) and unequal.
Verification by solving the equation.
$3x^2 + 5x - 12 = 0$
Using quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-(5) \pm \sqrt{(5)^2 - 4(3)(-12)}}{2(3)}$$

$$= \frac{-5 \pm \sqrt{25 + 144}}{6}$$

$$= \frac{-5 \pm \sqrt{169}}{6}$$

Thus, the roots are irrational (real) and unequal.

3. For what value of k, the expression $3x^2 + 5(k+1)x + 4$ **is perfect square.**

Solution:
$3^2 + 5(k+1)(x) + 4 = 0$
Compare it with
$ax^2 + bx + c = 0$
Also $a = 3, b = 5(k+1), c = 4$
Also, $\Delta = b^2 - 4ac$
$= [5(k+1)]^2 - 4(3)(4)$
$= (5k+5)^2 - 48$
$= (5k)^2 + (5)^2 + 2(5k)(5) - 48$
$= 25k^2 + 25 + 50k - 48$
$= 25k^2 + 50k - 23$

As the discr. is perfect square.
Therefore $\Delta = 0$
$25k^2 + 50k - 23 = 0$
$25k^2 + 50k = 23$
$25(k^2 + 2k) = 23$
$k^2 + 2k = \frac{23}{25}$
$k^2 + 2k + (1)^2 = \frac{23}{25} + (1)^2$
$(k+1)^2 = \frac{23}{25} + 1$
$(k+1)^2 = \frac{23+25}{25}$
$(k+1)^2 = \frac{48}{25}$
$k+1 = \sqrt{\frac{48}{25}}$
$k+1 = \frac{\sqrt{48}}{\sqrt{25}}$
$k+1 = \frac{\sqrt{16 \times 3}}{5}$
$k+1 = \frac{4\sqrt{3}}{5}$
$k = \frac{4\sqrt{3}}{5} - 1$
$k = \frac{4\sqrt{3} - 5}{5}$

$ax^2 + bx + c = 0$
Here $a = 2k + 1, b = 6k, c = 3$
As the roots are equal, As
$\Delta = 0$
$b^2 - 4ac = 0$
$(6)^2 - 4(2k+1)(3) = 0$
$36 - 12(2k+1) = 0$
$36 - 24k - 12 = 0$
$24 - 24k = 0$
$24 = 24k$
$\frac{24}{24} = k$
$1 = k$
$k = 1$

(ii) $x^2 + 2(k+2)x + (k+4) = 0$

Solution:
$x^2 + 2(k+2)x + (k+4) = 0$
Compare it with
$ax^2 + bx + c = 0$
Here $a = 1, b = 2(k+2), c = (k+4)$
As the roots are equal, As
$\Delta = 0$
$b^2 - 4ac = 0$
$[2(k+2)]^2 - 4(1)(k+4) = 0$
$4(k+2)^2 - 4(k+4) = 0$
$4[(k)^2 + (2)^2 + 2(k)(2)] - 4k - 16 = 0$
$4(k^2 + 4 + 4k) - 4k - 16 = 0$
$4k^2 + 16 + 16k - 4k - 16 = 0$
$4k^2 + 12k = 0$
$4k(k + 3) = 0$
$k = 0$

4. Show that the equation $x^2 + (ax + c)^2 = d^2$ **has equal roots, if** $c^2 = d^2(1 + a^2)$

Solution:
$x^2 + (ax + c)^2 = d^2$
$x^2 + a^2x^2 + 2(ax)(c) = d^2$
$(1 + a^2)x^2 + 2(ax)(c) = d^2$

$(1 + a^2)x^2 + 2(ax)(c) - d^2 = 0$
Here $a = (1 + a^2), b = 2(ax), c = -d^2$
As the roots are equal, As
$\Delta = 0$
$b^2 - 4ac = 0$
$[2(ax)]^2 - 4(1 + a^2)(-d^2) = 0$
$4a^2x^2 + 4(1 + a^2)d^2 = 0$
$4a^2x^2 + 4d^2 + 4a^2d^2 = 0$
$4a^2x^2 + 4a^2d^2 + 4d^2 = 0$
$4a^2x^2 + 4a^2d^2 = -4d^2$
$4a^2(x^2 + d^2) = -4d^2$
$x^2 + d^2 = \frac{-4d^2}{4a^2}$
$x^2 + d^2 = -\frac{d^2}{a^2}$
$x^2 = -\frac{d^2}{a^2} - d^2$
Hence proved

5. Find the condition that the roots of the equation $(ax+c)^2 - 4ax = 0$ **are equal.**

Solution:
$(ax+c)^2 - 4ax = 0$
$a^2x^2 + 2(ax)(c) - 4ax = 0$
$a^2x^2 + 2(ax)(c) - 4ax = 0$
$a^2x^2 + 2ax(c - 2) = 0$
As the roots are equal, As
$\Delta = 0$
$[2a(c-2)]^2 - 4(a^2)(0) = 0$
$4(a^2)(c-2)^2 - 0 = 0$
$4(a^2)(c-2)^2 = 0$
$(c-2)^2 = \frac{0}{4a^2}$
$(c-2)^2 = 0$
$c - 2 = \sqrt{0}$
$c - 2 = 0$
$c = 2$

6. Show that the roots of the following equations are rational.

(i) $(k-3)x^2 + 5kx + 1 = 0$

Solution:
$(k-3)x^2 + 5kx + 1 = 0$
Compare it with

$ax^2 + bx + c = 0$
Here $a = (k-3), b = 5k, c = 1$
As the roots are equal, As
$\Delta = 0$
$b^2 - 4ac = 0$
$(5k)^2 - 4(k-3)(1) = 0$
$25k^2 - 4(k-3) = 0$
$25k^2 - 4k + 12 = 0$
$25k^2 - 4k + 12 = 0$
$25k^2 - 4k = -12$
$25(k^2 - \frac{4}{25}k) = -12$
$k^2 - \frac{4}{25}k + (\frac{2}{25})^2 = -\frac{12}{25} + (\frac{2}{25})^2$
$(k - \frac{2}{25})^2 = -\frac{12}{25} + \frac{4}{625}$
$(k - \frac{2}{25})^2 = \frac{-300+4}{625}$
$(k - \frac{2}{25})^2 = -\frac{296}{625}$
$k - \frac{2}{25} = \sqrt{-\frac{296}{625}}$
$k - \frac{2}{25} = \frac{\sqrt{-296}}{\sqrt{625}}$
$k - \frac{2}{25} = \frac{\sqrt{-4 \times 74}}{25}$
$k - \frac{2}{25} = \frac{2\sqrt{-74}}{25}$
$k = \frac{2\sqrt{-74}}{25} + \frac{2}{25}$
$k = \frac{2\sqrt{-74} + 2}{25}$

7. Find the value of k, if the roots of the following equations are equal.

(i) $(2k+3)x^2 + 6kx + 3 = 0$

Solution:
$(2k+3)x^2 + 6kx + 3 = 0$
Compare it with

$ax^2 + bx + c = 0$
Here $a = (2k+3), b = 6k, c = 3$
As the roots are equal, As
$\Delta = 0$
$b^2 - 4ac = 0$
$(6k)^2 - 4(2k+3)(3) = 0$
$36k^2 - 12(2k+3) = 0$
$36k^2 - 24k - 36 = 0$
$36k^2 - 24k = 36$
$36(k^2 - \frac{2}{3}k) = 36$
$k^2 - \frac{2}{3}k + (\frac{1}{3})^2 = 1 + (\frac{1}{3})^2$
$(k - \frac{1}{3})^2 = 1 + \frac{1}{9}$
$(k - \frac{1}{3})^2 = \frac{9+1}{9}$
$(k - \frac{1}{3})^2 = \frac{10}{9}$
$k - \frac{1}{3} = \sqrt{\frac{10}{9}}$
$k - \frac{1}{3} = \frac{\sqrt{10}}{\sqrt{9}}$
$k - \frac{1}{3} = \frac{\sqrt{10}}{3}$
$k = \frac{\sqrt{10}}{3} + \frac{1}{3}$
$k = \frac{\sqrt{10} + 1}{3}$

(ii) $(k+2)x^2 - 5(k+1)x + (2k+5) = 0$

Solution:
$(k+2)x^2 - 5(k+1)x + (2k+5) = 0$
Compare it with
$ax^2 + bx + c = 0$
Here $a = (k+2), b = -5(k+1), c = (2k+5)$
As the roots are equal, As
$\Delta = 0$
$b^2 - 4ac = 0$
$[-5(k+1)]^2 - 4(k+2)(2k+5) = 0$
$25(k+1)^2 - 4(k+2)(2k+5) = 0$
$25[(k)^2 + (1)^2 + 2(k)(1)] - 4[2k^2 + 5k + 4k + 10] = 0$
$25(k^2 + 1 + 2k) - 4[2k^2 + 9k + 10] = 0$
$25k^2 + 25 + 50k - 8k^2 - 36k - 40 = 0$
$17k^2 + 14k - 15 = 0$
$17k^2 + 14k = 15$
$17(k^2 + \frac{14}{17}k) = 15$
$k^2 + \frac{14}{17}k + (\frac{7}{17})^2 = \frac{15}{17} + (\frac{7}{17})^2$
$(k + \frac{7}{17})^2 = \frac{15}{17} + \frac{49}{289}$
$(k + \frac{7}{17})^2 = \frac{255+49}{289}$
$(k + \frac{7}{17})^2 = \frac{304}{289}$
$k + \frac{7}{17} = \sqrt{\frac{304}{289}}$
$k + \frac{7}{17} = \frac{\sqrt{304}}{\sqrt{289}}$
$k + \frac{7}{17} = \frac{\sqrt{16 \times 19}}{17}$
$k + \frac{7}{17} = \frac{4\sqrt{19}}{17}$
$k = \frac{4\sqrt{19}}{17} - \frac{7}{17}$
$k = \frac{4\sqrt{19} - 7}{17}$

8. $x^2 - \left(\frac{k}{2}\right)x + 1 = 0, k > 0$ **are real.**

Solution:
$x^2 - \left(\frac{k}{2}\right)x + 1 = 0$
Here $a = 1, b = -\left(\frac{k}{2}\right), c = 1$

Also,
$\Delta = b^2 - 4ac$
$= \left(-\frac{k}{2}\right)^2 - 4(1)(1)$
$= \frac{k^2}{4} - 4$
$= \frac{k^2 - 16}{4}$
$= \frac{1}{4}(k^2 - 16)$
$= \frac{1}{4}(k^2 - 4^2)$
$= \frac{1}{4}(k+4)(k-4)$

As the roots are real.
Therefore $\Delta \geq 0$
$\frac{1}{4}(k+4)(k-4) \geq 0$
$(k+4)(k-4) \geq 0$
Hence, $k \geq 4$ or $k \leq -4$

9. Show that the roots of the equation $(a^2+b^2)x^2 - 2(ac+bd)x + (c^2+d^2) = 0$ **are equal.**

Solution:
$(a^2+b^2)x^2 - 2(ac+bd)x + (c^2+d^2) = 0$
Here $a = (a^2+b^2), b = -2(ac+bd), c = (c^2+d^2)$
Also,
$\Delta = b^2 - 4ac$
$= [-2(ac+bd)]^2 - 4(a^2+b^2)(c^2+d^2)$
$= 4(ac+bd)^2 - 4(a^2+b^2)(c^2+d^2)$
$= 4[(ac)^2 + (bd)^2 + 2(ac)(bd)] - 4[a^2c^2 + a^2d^2 + b^2c^2 + b^2d^2]$
$= 4[a^2c^2 + b^2d^2 + 2abcd] - 4[a^2c^2 + a^2d^2 + b^2c^2 + b^2d^2]$
$= 4a^2c^2 + 4b^2d^2 + 8abcd - 4a^2c^2 - 4a^2d^2 - 4b^2c^2 - 4b^2d^2$
$= 8abcd - 4a^2d^2 - 4b^2c^2$
$= -4a^2d^2 + 8abcd - 4b^2c^2$
$= -4[a^2d^2 - 2abcd + b^2c^2]$
$= -4[ad - bc]^2$

As the roots are equal.
Therefore $\Delta = 0$
$-4[ad - bc]^2 = 0$
$[ad - bc]^2 = \frac{0}{-4}$
$[ad - bc]^2 = 0$
$ad - bc = \sqrt{0}$
$ad - bc = 0$
$ad = bc$
Hence the roots are rational.

10. Show that the roots of the equation
$(b-c)x^2 + (c-a)x + (a-b) = 0$ **are real.**

Solution:
$(b-c)x^2 + (c-a)x + (a-b) = 0$
Here $a = (b-c), b = (c-a), c = (a-b)$
Also,
$\Delta = b^2 - 4ac$
$= (c-a)^2 - 4(b-c)(a-b)$
$= c^2 + a^2 - 2ac - 4[ab - b^2 - ac + bc]$
$= c^2 + a^2 - 2ac - 4ab + 4b^2 + 4ac - 4bc$

$= a^2 + 4b^2 + c^2 - 4ab - 4bc + 2ac$
$= (a)^2 + (-4b)^2 + (c)^2 + 2(a)(-4b) + 2(-4b)(c) + 2(a)(c)$
$= [a - 4b + c]^2 \geq 0$

Hence the roots of equation are real.

Cube roots of unity and their properties
The cube roots of unity
Let's consider now the cube root of unity.

$$x = \sqrt[3]{1}$$

Solution:

$x = \sqrt[3]{1}$

$x^3 = 1^3$

$x^3 - 1^3 = 0$

$(x-1)(x^2+x+1) = 0$ [using $a^3 - b^3 = (a-b)(a^2+ab+b^2)$]

Either $x-1=0$ or $x^2+x+1=0$

or $x = 1$ or $x = \frac{-1 \pm \sqrt{1-4}}{2}$

$x = \frac{-1 \pm \sqrt{-3}}{2} = \frac{-1 \pm \sqrt{3}i}{2}$

Or we take note of what we:

$x = \left(\frac{-1+\sqrt{3}i}{2}\right)$ and $x = \left(\frac{-1-\sqrt{3}i}{2}\right)$

Recognize complete cube roots of unity as 1^{st} and 2^{nd}.

The first complete cube root of unity are $\frac{-1+\sqrt{3}i}{2}$ and $\frac{-1-\sqrt{3}i}{2}$.

If we name anyone of these as ω (pronounced as omega), then the others is ω^2

The second complete cube root of unity are

(a) Prove that each of the complex cube roots of unity is the square of the other.
Proof:

The complex cube roots of unity are $\frac{-1+\sqrt{3}i}{2}$ and $\frac{-1-\sqrt{3}i}{2}$.

We prove that:

Solution:

$\left(\frac{-1+\sqrt{3}i}{2}\right)^2 = \frac{-1-\sqrt{3}i}{2}$ and $\left(\frac{-1-\sqrt{3}i}{2}\right)^2 = \frac{-1+\sqrt{3}i}{2}$

$\left(\frac{-1+\sqrt{3}i}{2}\right)\left(\frac{-1+\sqrt{3}i}{2}\right) = \frac{-1-\sqrt{3}i}{2}$ $\left(\frac{-1-\sqrt{3}i}{2}\right)\left(\frac{-1-\sqrt{3}i}{2}\right) = \frac{-1+\sqrt{3}i}{2}$

$\frac{1-\sqrt{3}i-\sqrt{3}i+3i^2}{4}$ $\frac{1+\sqrt{3}i+\sqrt{3}i+3i^2}{4}$

$\frac{1-2\sqrt{3}i-3}{4}$ $\frac{1+2\sqrt{3}i-3}{4}$

$\frac{-2-2\sqrt{3}i}{4}$ $\frac{-2+2\sqrt{3}i}{4}$

$\frac{-1-\sqrt{3}i}{2}$ $\frac{-1+\sqrt{3}i}{2}$

Thus, each of the complex cube root of unity is the square of the other, that's,

$\omega = \left(\frac{-1+\sqrt{3}i}{2}\right)$ then $\omega^2 = \frac{-1-\sqrt{3}i}{2}$ Also

$\omega = \frac{-1-\sqrt{3}i}{2}$

(b) Prove that the product of three cube roots of unity is one.
Proof:

Solution:

The cube roots of unity are

$\left(\frac{-1+\sqrt{3}i}{2}\right)$, 1, $\left(\frac{-1-\sqrt{3}i}{2}\right)$

The product of the roots of unity $= \left(\frac{-1+\sqrt{3}i}{2}\right)\left(\frac{-1-\sqrt{3}i}{2}\right)(1)$

$= \frac{1-\sqrt{3}i-\sqrt{3}i+3i^2}{4}(1) = \frac{1-2\sqrt{3}i-3}{4} = \frac{-2-2\sqrt{3}i}{4}$

$= \frac{-1-\sqrt{3}i}{2}(1) = \omega \cdot \omega^2 = 1$

(c) Prove that each complex cube root of unity is reciprocal of the other.
Proof:

We know that $\omega^3 = 1$ $\omega \cdot \omega^2 = 1, \omega$

$\omega = \frac{1}{\omega^2}$ or $\omega^2 = \frac{1}{\omega}$

Thus, each complex cube root of unity is reciprocal of the other.

(d) Prove that the sum of all the cube roots of unity is zero.

$\omega, 1, \omega, \omega^2 = 0$

Proof:

The cube roots of unity are

$\left(\frac{-1+\sqrt{3}i}{2}\right)$, 1, $\left(\frac{-1-\sqrt{3}i}{2}\right)$

If $\omega = \left(\frac{-1+\sqrt{3}i}{2}\right)$ then $\omega^2 = \left(\frac{-1-\sqrt{3}i}{2}\right)$

The sum of the roots $1 + \omega + \omega^2$

$= 1 + \left(\frac{-1+\sqrt{3}i}{2}\right) + \left(\frac{-1-\sqrt{3}i}{2}\right)$

$= 1 + \frac{-1+\sqrt{3}i-1-\sqrt{3}i}{2} = 1 + \frac{-2}{2} = 1 - 1 = 0$

Thus $1 + \omega + \omega^2 = 0$

We can apply index as following below, that is

$\left(\frac{-1+\sqrt{3}i}{2}\right)^3 = \left(\frac{-1-\sqrt{3}i}{2}\right)^3 = 1$ and $\omega^3 = 1$

Exercise 2.2

1. Find the cube roots of -8. - 07 JA

(i) The three cube roots of -1

Solution:

Let $x^3 = -1$

$x^3 + 1 = 0$

$(x+1)(x^2 - x + 1) = 0$

Either $x + 1 = 0$ or $x^2 - x + 1 = 0$

$x = -1$ Where $a = 1, b = -1, c = 1$

Using quadratic formula

$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)(1)}}{2(1)}$

$x = \frac{1 \pm \sqrt{1-4}}{2}$

$x = \frac{1 \pm \sqrt{-3}}{2}$

$x = \frac{1 \pm \sqrt{3}i}{2}$ or $x = \frac{1-\sqrt{3}i}{2}$

$x = \left(\frac{1+\sqrt{3}i}{2}\right)$ or $x = \left(\frac{1-\sqrt{3}i}{2}\right)$

$x = \omega$ $x = \omega^2$

These cube roots of -1 are $-1, \omega, \omega^2$

(ii) The three cube roots of 8

Solution:

Let $x^3 = 8$

$x^3 - 8 = 0$

$(x-2)(x^2 + 2x + 4) = 0$

Either $x - 2 = 0$ or $x^2 + 2x + 4 = 0$

$x = 2$ Where $a = 1, b = 2, c = 4$

Using quadratic formula

$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

$x = \frac{-(2) \pm \sqrt{(2)^2 - 4(1)(4)}}{2(1)}$

$x = \frac{-2 \pm \sqrt{4-16}}{2}$

$x = \frac{-2 \pm \sqrt{-12}}{2}$

$x = \frac{-2 \pm \sqrt{12}i}{2}$

$x = \frac{-2 \pm 2\sqrt{3}i}{2}$

$x = \frac{2(-1 \pm \sqrt{3}i)}{2}$

$x = \left(-1 \pm \sqrt{3}i\right)$ or $x = \left(-1 - \sqrt{3}i\right)$

$x = -\omega$ $x = -\omega^2$

These cube roots of 8 are $2, 2\omega, 2\omega^2$

(iii) The three cube roots of -27

Solution:

Let $x^3 = -27$

$x^3 + 27 = 0$

$(x+3)(x^2 - 3x + 9) = 0$

$(x+3)(x^2 - 3x + 9) = 0$

Either $x + 3 = 0$ or $x^2 - 3x + 9 = 0$

$x = -3$ Where $a = 1, b = -3, c = 9$

Using quadratic formula

$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

$x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(9)}}{2(1)}$

$x = \frac{3 \pm \sqrt{9-36}}{2}$

$x = \frac{3 \pm \sqrt{-27}}{2}$

$x = \frac{3 \pm \sqrt{27}i}{2}$

$x = \frac{3 \pm 3\sqrt{3}i}{2}$

$x = \left(\frac{3+3\sqrt{3}i}{2}\right)$ or $x = \left(\frac{3-3\sqrt{3}i}{2}\right)$

$x = -3\omega$ $x = -3\omega^2$

These cube roots of -27 are $-3, -3\omega, -3\omega^2$

(iv) The three cube roots of 64

Solution:

Let $x^3 = 64$

$x^3 - 64 = 0$

$(x-4)(x^2 + 4x + 16) = 0$

Either $x - 4 = 0$ or $x^2 + 4x + 16 = 0$

$x = 4$ Where $a = 1, b = 4, c = 16$

Using quadratic formula

$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

$x = \frac{-(4) \pm \sqrt{(4)^2 - 4(1)(16)}}{2(1)}$

$x = \frac{-4 \pm \sqrt{16-64}}{2}$

$x = \frac{-4 \pm \sqrt{-48}}{2}$

$x = \frac{-4 \pm \sqrt{48}i}{2}$

$x = \frac{-4 \pm 4\sqrt{3}i}{2}$

$x = \frac{4(-1 \pm \sqrt{3}i)}{2}$

$x = \left(-1 \pm \sqrt{3}i\right)$ or $x = \left(-1 - \sqrt{3}i\right)$

$x = -\omega$ $x = -\omega^2$

These cube roots of 64 are $4, 4\omega, 4\omega^2$

2. If evaluate

$\left(1 - \omega - \omega^2\right)^3$

Solution:

$\left(1 - \omega - \omega^2\right)^3$, since $\omega^2 = \frac{1}{\omega}$

$= \left(1 - \omega - \frac{1}{\omega}\right)^3$ $= \omega \cdot \omega^2 = 1$

$= \left(1 - \frac{\omega^2 - 1}{\omega}\right)^3$

$= \left(1 - \frac{\omega^2 - 1}{\omega}\right)^3$

$= \left(\frac{\omega - \omega^2 + 1}{\omega}\right)^3$

$= \frac{(\omega - \omega^2 + 1)^3}{\omega^3}$ $\omega^3 = 1$

$= \frac{(\omega - \omega^2 + 1)^3}{1}$

$= (\omega - \omega^2 + 1)^3$

(b) $\left(3 - 3\omega - 3\omega^2\right)^3$

Solution

$\left(3 - 3\omega - 3\omega^2\right)^3 = 3^3\left(1 - \omega - \omega^2\right)^3$

$= 27\left(1 - \omega - \omega^2\right)^3$ $\omega \cdot \omega^2 = 1$

$= 27(1)^3$

$= 27 \times 1$

$= 27$

(c) $\left(3 + 4\omega + 4\omega^2\right)^3$

Solution

$\left(3 + 4\omega + 4\omega^2\right)^3 = 3^3\left(1 + \frac{4}{3}\omega + \frac{4}{3}\omega^2\right)^3$

$= 27\left(1 + \frac{4}{3}\omega + \frac{4}{3}\omega^2\right)^3$ $\omega \cdot \omega^2 = 1$

$= 27\left(\frac{3}{3} + \frac{4}{3}\omega + \frac{4}{3}\omega^2\right)^3$ $3\omega = 3\omega \cdot \omega^2 = 3\omega^3$

$= 27\left(\frac{3}{3} + \frac{4}{3}\omega + \frac{4}{3}\omega^2\right)^3$ $\omega^3 = 1$

$= 27\left(\frac{3}{3}\right)^3$

$= 27(1)^3$

$= 27 \times 1$

$= 27$

(d) $\left(2 + 2\omega - 2\omega^2\right)\left(3 - 3\omega + 3\omega^2\right)$

Solution:

$\left(2 + 2\omega - 2\omega^2\right)\left(3 - 3\omega + 3\omega^2\right)$

$= 2\left(1 + \omega - \omega^2\right)3\left(1 - \omega + \omega^2\right)$

$= 6\left(1 + \omega - \omega^2\right)\left(1 - \omega + \omega^2\right)$ $(1 + \omega = -\omega^2$

$= 6\left(-\omega^2 - \omega^2\right)\left(1 - \omega + \omega^2\right)$

$= 6\left(-2\omega^2\right)\left(1 - \omega + \omega^2\right)$

$= -12\omega^2\left(1 - \omega + \omega^2\right)$

$= -12\omega^2 + 12\omega^3 - 12\omega^4$

$= -12\omega^2 + 12(1) - 12\omega^3 \cdot \omega$

$= -12\omega^2 + 12 - 12(1)\omega$

$= -12\omega^2 + 12 - 12\omega$

$= 12 - 12\omega - 12\omega^2$

$= 12 - 12\left(\omega + \omega^2\right)$

$= 12 - 12(-1)$

$= 12 + 12$

$= 24$

(e) $\left(1 + \sqrt{2}\omega\right)^3 + \left(1 - \sqrt{2}\omega\right)^3$

Solution:

$\left(1 + \sqrt{2}\omega\right)^3 + \left(1 - \sqrt{2}\omega\right)^3$

$= \left[(a+b)^3 + (a-b)^3\right]$ $= 2\left(a^3 + 3ab^2\right)$ $\omega^3 = 1, \frac{1}{\omega} = \omega^2$

$= 2\left[a^3 + 3ab^2\right]$ $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ $(a-b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$

$= 2\left[a^3 + 3a^2b^2\right]$

$= 2\left[1 + 3(1)^2\left(\sqrt{2}\right)^2\right]$ $\omega^3 = 1$

$= 2[1 + 6]$

$= 2[7]$

$= 14$

(e)(ii) $\left(1 + \sqrt{2}\omega^2\right)^3 + \left(1 - \sqrt{2}\omega^2\right)^3$

Solution:

$\left(1 + \sqrt{2}\omega^2\right)^3 + \left(1 - \sqrt{2}\omega^2\right)^3$

$= \left[(a+b)^3 + (a-b)^3\right]$ $= 2\left(a^3 + 3ab^2\right)$ $\omega^3 = 1, \frac{1}{\omega} = \omega^2$

$= 2\left[a^3 + 3ab^2\right]$ $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ $(a-b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$

$= 2\left[a^3 + 3a^2b^2\right]$

$= 2\left[1 + 3(1)^2\left(\sqrt{2}\right)^2\right]$ $\omega^3 = 1$

$= 2[1 + 6]$

$= 2[7]$

$= 14$

(e)(iii) $\left(1 + \sqrt{2}\omega\right)^3 + \omega^{-3}$

Solution:

$= 1 + \sqrt{2}\omega^3$

$= \omega^{-3} = \omega^{-3}$

$= 1 + \sqrt{2}(1)$

$= \omega^{-3} = \omega^3 \cdot \omega^{-3}$

$= 1 + \sqrt{2}$

$= \omega^0$

$= 1$

(e)(iv) $\left(1 + \sqrt{2}\omega^2\right)^3 + \omega^{-3}$

Solution:

$= 1 + \sqrt{2}\omega^3$

$= \omega^{-3} = \omega^{-3}$

$= 1 + \sqrt{2}(1)$

$= \omega^{-3} = \omega^3 \cdot \omega^{-3}$

$= 1 + \sqrt{2}$

$= \omega^0$

$= 1$

$= \left(\omega^2\right)^3 + \omega^{-3}$ $\omega^{-3} = \frac{1}{\omega^3}$

$= \omega^6 + \omega^{-3}$

$= \omega^3 \cdot \omega^3 + \omega^{-3}$

$= 1 \cdot 1 + \omega^{-3}$

$= 1 + \omega^{-3}$

$= 1 + \omega^3 \cdot \omega^{-3}$

$= 1 + \omega^0$

$= 1 + 1$

$= 2$

3. Prove that $x^2 + y^2 - 2xy = (x + y\omega)(x + y\omega^2)$, if $\omega^3 = 1$.

Solution:

$x^2 + y^2 - 2xy = (x + y\omega)(x + y\omega^2)$

R.H.S $= (x + y\omega)(x + y\omega^2)$

$= x(x + y\omega^2) + y\omega(x + y\omega^2)$ $\omega^3 = 1, \omega \cdot \omega^2 = 1$

$= x^2 + xy\omega^2 + xy\omega + y^2\omega^3$

$= x^2 + xy\omega^2 + xy\omega + y^2(1)$ $\omega^3 = 1$

$= x^2 + xy\omega^2 + xy\omega + y^2$ $\omega^2 + \omega = -1$

$= x^2 + xy(-1) + y^2$

$= x^2 - xy + y^2$

$= x^2 - 2xy + xy + y^2$

$= x^2 - 2xy + y^2$

$= L.H.S$

Hence Proved.

4. Prove that $x^2 + y^2 + z^2 - 2yz - 2zx - 2xy = (x + y + z)(x + y\omega + z\omega^2)(x + y\omega^2 + z\omega)$,

Solution:

$x^2 + y^2 + z^2 - 2yz - 2zx - 2xy = (x + y + z)(x + y\omega + z\omega^2)(x + y\omega^2 + z\omega)$

R.H.S $= (x + y + z)(x + y\omega + z\omega^2)(x + y\omega^2 + z\omega)$

$= (x + y + z)\left[x(x + y\omega^2 + z\omega) + y\omega(x + y\omega^2 + z\omega) + z\omega^2(x + y\omega^2 + z\omega)\right]$

$= (x + y + z)\left[x^2 + xy\omega^2 + xz\omega + xy\omega + y^2\omega^3 + yz\omega^2 + xz\omega^2 + yz\omega^3 + z^2\omega^4\right]$

$= (x + y + z)\left[x^2 + xy\omega^2 + xz\omega + xy\omega + y^2(1) + yz\omega^2 + xz\omega^2 + yz(1) + z^2\omega\right]$ $\omega^3 = 1$

$= (x + y + z)\left[x^2 + xy\omega^2 + xz\omega + xy\omega + y^2 + yz\omega^2 + xz\omega^2 + yz + z^2\omega\right]$ $\omega^4 = \omega \cdot \omega^3 = \omega$

$= (x + y + z)\left[x^2 + y^2 + xy\omega + xy\omega^2 + xz\omega + xz\omega^2 + yz\omega^2 + yz + z^2\omega\right]$

$= (x + y + z)\left[x^2 + y^2 + xy(\omega + \omega^2) + xz(\omega + \omega^2) + yz(\omega^2 + 1) + z^2\omega\right]$ $\omega^2 + \omega = -1, \omega^2 + 1 = -\omega$

$= (x + y + z)\left[x^2 + y^2 + xy(-1) + xz(-1) + yz(-\omega) + z^2\omega\right]$

$= (x + y + z)\left[x^2 + y^2 - xy - xz - yz\omega + z^2\omega\right]$

$= x^3 + xy^2 + xz^2 - x^2y - x^2z - xyz\omega + xz^2\omega + xy^2 + y^3 + yz^2 - x^2y - xyz\omega - y^2z\omega + yz^2\omega + x^2z + y^2z + z^3\omega - xyz - xz^2\omega - yz^2\omega$

$= x^3 + y^3 + z^3 - 3xyz$

$= L.H.S$

Hence Proved.

5. Prove that $(1 + \omega)^3 + (1 + \omega^2)^3 = (x + \omega^2)^3 + (x + \omega)^3$ **... 2x factors=2,**

Solution:

L.H.S $= (1 + \omega)^3 + (1 + \omega^2)^3 = (-\omega^2)^3 + (-\omega)^3$ ... 2x factors

$= (1 + \omega)(1 + \omega)^2 + (1 + \omega^2)^3 = (-\omega^2)^3 + (-\omega)^3$ ... 2x factors

$= (1 + \omega)(1 + \omega)^2 + (1 + \omega^2)^3 = -\omega^6 - \omega^3$ ... 2x factors

$= (1 + \omega)(1 + \omega^2 + 2\omega) + (1 + \omega^2)^3 = -\omega^3 \cdot \omega^3 - \omega^3$ ... 2x factors $\omega^3 = 1$

$= (1 + \omega)(\omega^2 + 2\omega) + (1 + \omega^2)^3 = -(1)(1) - (1)$ ... 2x factors

$= (1 + \omega)(\omega^2 + 2\omega) + (1 + \omega^2)^3 = -1 - 1$ ... 2x factors

$= (1 + \omega)(\omega^2 + 2\omega) + (1 + \omega^2)^3 = -2$ ... 2x factors

$= \left[(-\omega^2)^3 + (-\omega)^3\right] = -2$ ... 2x factors $\omega^2 + \omega = -1$

$= \left[-\omega^6 - \omega^3\right] = -2$ ... 2x factors $\omega^3 = 1$

$= \left[-(1)^2 - (1)\right] = -2$ ... 2x factors $1 + \omega = -\omega^2$

$= -1 - 1 = -2$ ... 2x factors $1 + \omega^2 = -\omega$

$= -2$ ... 2x factors

$= -2$

$= R.H.S$

Hence Proved.

Roots and co-efficient of a quadratic equation

We know that $\frac{-b+\sqrt{b^2-4ac}}{2a}$ and $\frac{-b-\sqrt{b^2-4ac}}{2a}$ are roots of the equation $ax^2+bx+c=0$ where a,b are coefficients of x^2 and x respectively. While c is the constant term.

Relation between roots and co-efficient of a quadratic equation

If $\alpha = \frac{-b+\sqrt{b^2-4ac}}{2a}$ and $\beta = \frac{-b-\sqrt{b^2-4ac}}{2a}$

Show in turn with the roots and the position of the roots as follows;

Sum of the roots $\alpha + \beta$

$$= \frac{-b+\sqrt{b^2-4ac}}{2a} + \frac{-b-\sqrt{b^2-4ac}}{2a}$$
$$= \frac{-b+\sqrt{b^2-4ac}-b-\sqrt{b^2-4ac}}{2a}$$
$$= \frac{-2b}{2a} = -\frac{b}{a}$$

Product of the roots $\alpha \times \beta$

$$= \frac{-b+\sqrt{b^2-4ac}}{2a} \times \frac{-b-\sqrt{b^2-4ac}}{2a}$$
$$= \frac{(-b+\sqrt{b^2-4ac})(-b-\sqrt{b^2-4ac})}{2a \times 2a}$$
$$= \frac{b^2+b\sqrt{b^2-4ac}-b\sqrt{b^2-4ac}-(b^2-4ac)}{4a^2}$$
$$= \frac{b^2-b^2+4ac}{4a^2} = \frac{4ac}{4a^2} = \frac{c}{a}$$

If we divide the sum of roots and product of roots by A and P respectively we get;

$\frac{\alpha+\beta}{A} = \frac{-\frac{b}{a}}{\frac{-b}{a}} = \frac{-b}{a} \times \frac{a}{-b} = \frac{-1}{-1}$ and $\frac{\alpha \times \beta}{P} = \frac{\frac{c}{a}}{\frac{c}{a}} = \frac{c}{a} \times \frac{a}{c} = \frac{-1}{-1}$

Exercise 2.3

1. Without solving, find the sum and the product of the following quadratic equations.

(i) $x^2 - 7x + 3 = 0$

Solution:

$x^2 - 7x + 3 = 0$

Here $a=1, b=-7, c=3$

Let α and β be the roots of the given equation.

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{(-7)}{1} = 7$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{3}{1}$

(ii) $2x^2 + 7x - 11 = 0$

Solution:

$2x^2 + 7x - 11 = 0$

Here $a=2, b=7, c=-11$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{7}{2}$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{-11}{2}$

(iii) $px^2 - cpx + r = 0$

Solution:

$px^2 - cpx + r = 0$

Here $a=p, b=-cp, c=r$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{(-cp)}{p} = c$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{r}{p}$

(iv) $\frac{7}{2}x^2 - 2ax + 9a = 0$

Solution:

$\frac{7}{2}x^2 - 2ax + 9a = 0$

Here $a=\frac{7}{2}, b=-2a, c=9a$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{(-2a)}{\frac{7}{2}} = \frac{4a}{7}$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{9a}{\frac{7}{2}}$

(v) $(x+a)(x-a) + (x-a)(x+a) - c^2 = 0$

Solution:

$(x+a)(x-a) + (x-a)(x+a) - c^2 = 0$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{(0)}{2} = 0$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{-2a^2-c^2}{2}$

(vi) $\frac{x}{a} + \frac{a}{x} = \frac{b}{c}, x \neq 0$

Solution:

$\frac{x}{a} + \frac{a}{x} = \frac{b}{c}, x \neq 0$

Here $a=\frac{1}{a}, b=-\frac{b}{c}, c=1$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{\left(-\frac{b}{c}\right)}{\frac{1}{a}} = \frac{ab}{c}$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{1}{\frac{1}{a}} = a$

2. Find the value of k, if

(i) Sum of the roots of the equation $2kx^2 - 5x + 8 = 0$ is twice the product of the roots.

Solution:

$2kx^2 - 5x + 8 = 0$

Here $a=2k, b=-5, c=8$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{(-5)}{2k} = \frac{5}{2k}$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{8}{2k}$

As one of the roots is twice the product of the roots, so

$\alpha = \beta + 2\alpha\beta$

$\frac{5}{2k} = \frac{8}{2k}$

$5 = 8$

$\alpha \neq \beta$

(ii) Sum of the roots of the equation $x^2 + (k-7)x - 7k = 0$ is three times the product of the roots.

Solution:

$x^2 + (k-7)x - 7k = 0$

Here $a=1, b=(k-7), c=-7k$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{(k-7)}{1} = (-k+7)$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{-7k}{1} = -7k$

As one of the roots α is the product of the roots, so

$\alpha = \alpha\beta$

$-k+7 = -7k$

$-k+7k = -7$

$6k = -7$

$k = \frac{-7}{6}$

(iii) Sum of the squares of the roots of the equation $4kx^2 + 12kx - 9 = 0$ is 2.

Solution:

$4kx^2 + 12kx - 9 = 0$

Here $a=4k, b=12k, c=-9$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{12k}{4k} = -3$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{-9}{4k}$

As sum of the squares of roots is 2, so

$\alpha^2 + \beta^2 = 2$

$k = \frac{-9}{8}$

3. Find k, if

(i) Sum of the squares of the roots of the equation $4kx^2 + 12kx - 9 = 0$ is 2.

Solution:

$4kx^2 + 12kx - 9 = 0$

Here $a=4k, b=12k, c=-9$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{12k}{4k} = -3$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{-9}{4k}$

As sum of the squares of roots is two the product of the roots, so

$(\alpha+\beta)^2 - 2\alpha\beta = 2$ $\quad$ $(-3)^2 - 2\left(\frac{-9}{4k}\right) = 2$

$\left(\frac{-3}{1}\right)^2 + \left(\frac{9}{2k}\right) = 2$ $\quad$ $\alpha^2 + \beta^2 = (\alpha+\beta)^2 - 2\alpha\beta$

$\frac{9}{1} + \frac{9}{2k} = 2$

$\frac{18k+9}{2k} = \frac{2}{1}$ $\quad$ $\frac{18k+9}{2k} = \frac{2}{1}$

$18k + 9 = 4k$ $\quad$ $18k - 4k = -9$

$k = \frac{-9}{14}$ $\quad$ $14k = -9$

$k = \frac{-9}{14}$

(ii) Sum of the squares of the roots of the equation $x^2 - 10x + 16 = 0$ is 4.

Solution:

$x^2 - 10x + 16 = 0$

Here $a=1, b=-10, c=16$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{(-10)}{1} = 10$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{16}{1} = 16$

As sum of the squares of roots is four the product of the roots, so

$(\alpha+\beta)^2 - 2\alpha\beta = 4$ $\quad$ $(\alpha+\beta)^2 = \alpha^2 + \beta^2 + 2\alpha\beta$

$(10)^2 - 2(16) = 4$ $\quad$ $\alpha^2 + \beta^2 = (\alpha+\beta)^2 - 2\alpha\beta$

$100 - 32 = 4$

$\alpha^2 + \beta^2 = 4$

$\alpha^2 + \beta^2 = 68$ $\neq 4$

$\alpha^2 + \beta^2 = 4$ is not possible.

(iii) Sum of the squares of the roots of the equation $x^2 - 2(k-1)x + k = 0$ is 6.

Solution:

$x^2 - 2(k-1)x + k = 0$

Here $a=1, b=-2(k-1), c=k$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{-2(k-1)}{1} = 2(k-1)$

And product of roots $= \alpha\beta = \frac{c}{a} = \frac{k}{1} = k$

As sum of the squares of roots is six the product of the roots, so

$\alpha^2 + \beta^2 = 6$

$(\alpha+\beta)^2 - 2\alpha\beta = 6$ $\quad$ $(\alpha+\beta)^2 = \alpha^2 + \beta^2 + 2\alpha\beta$

$[2(k-1)]^2 - 2(k) = 6$ $\quad$ $\alpha^2 + \beta^2 = (\alpha+\beta)^2 - 2\alpha\beta$

$4(k^2 - 2k + 1) - 2k = 6$

$4k^2 - 8k + 4 - 2k = 6$

$4k^2 - 10k + 4 = 6$

$4k^2 - 10k + 4 - 6 = 0$

$4k^2 - 10k - 2 = 0$

$2k^2 - 5k - 1 = 0$

$k = \frac{-(-5) \pm \sqrt{(-5)^2 - 4(2)(-1)}}{2(2)}$ $\quad$ or $\quad$ $k = \frac{5 \pm \sqrt{33}}{4}$

$k = \frac{5+\sqrt{33}}{4}$ $\quad$ $k = \frac{5-\sqrt{33}}{4}$

4. Find a, if

(i) The roots of the equation $x^2 - ax + p = 0$ differ by unity.

Solution:

$x^2 - ax + p = 0$

Here $a=1, b=-a, c=p$

Let α and β be the roots of the given equation

Then $\alpha + \beta = \frac{a}{1}$ and $\alpha\beta = \frac{p}{1}$

$\alpha - \beta = 1$ $\quad$ $\alpha\beta = p$

$(\alpha - \beta)^2 = 1^2$ $\quad$ $\alpha = \beta + 1$ (where α is greater)

$(\alpha+\beta)^2 - 4\alpha\beta = 1$ $\quad$ $(\beta+1)\beta = p$

$a^2 - 4p = 1$ $\quad$ $\beta^2 + \beta = p$

$a^2 = 1 + 4p$ $\quad$ $\beta^2 + \beta - p = 0$

(ii) The roots of the equation $x^2 + 4x + 3(1+k) = 0$ differ by 2.

Solution:

$x^2 + 4x + 3(1+k) = 0$

Here $a=1, b=4, c=3(1+k)$

Let α and β be the roots of the given equation

Then $\alpha + \beta = -\frac{b}{a} = -\frac{4}{1}$ and $\alpha\beta = \frac{c}{a} = \frac{3(1+k)}{1}$

$\alpha - \beta = 2$ $\quad$ $\alpha\beta = \beta(\beta+2)$

$(\alpha - \beta)^2 = 2^2$ $\quad$ $\alpha\beta = \beta^2 + 2\beta$ (where α is greater)

$(\alpha+\beta)^2 - 4\alpha\beta = 4$ $\quad$ $\beta(\beta+2) = 3(1+k)$

$(-4)^2 - 4(3)(1+k) = 4$ $\quad$ $\beta^2 + 2\beta = 3 + 3k$

$16 - 12(1+k) = 4$

$16 - 12 - 12k = 4$

$4 - 12k = 4$

$-12k = 4 - 4$

$-12k = 0$

$k = 0$

$\beta + \beta + 2 = 3 + 3k$

$2\beta + 2 = 3 + 3k$

$2\beta = 3 + 3k - 2$

$2\beta = 1 + 3k$

$\beta = \frac{1+3k}{2}$

$\beta(\beta+2) = 3(1+k)$

$\left(\frac{1+3k}{2}\right)\left(\frac{1+3k}{2}+2\right) = 3(1+k)$

$\left(\frac{1+3k}{2}\right)\left(\frac{1+3k+4}{2}\right) = 3(1+k)$

$\frac{(1+3k)(5+3k)}{4} = 3(1+k)$

$(1+3k)(5+3k) = 12(1+k)$

$5 + 3k + 15k + 9k^2 = 12 + 12k$

$9k^2 + 18k + 5 - 12 - 12k = 0$

$9k^2 + 6k - 7 = 0$

For $\alpha = \beta + 2$ we get;

$\alpha = \frac{1+3k}{2} + 2$

$\alpha = \frac{1+3k+4}{2}$

$\alpha = \frac{5+3k}{2}$

For $\alpha = \beta$ we get;

$\frac{5+3k}{2} = \frac{1+3k}{2}$

$5 + 3k = 1 + 3k$

$5 = 1$

$\alpha \neq \beta$

(iii) The roots of the equation $3x^2 - 2x + 7(1+k) = 0$ differ by 3.

Solution:

$3x^2 - 2x + 7(1+k) = 0$

Here $a=3, b=-2, c=7(1+k)$

Let α and β be the roots of the given equation

Then $\alpha + \beta = -\frac{b}{a} = -\frac{-2}{3} = \frac{2}{3}$ and $\alpha\beta = \frac{c}{a} = \frac{7(1+k)}{3}$

$\alpha - \beta = 3$ $\quad$ $\alpha\beta = \beta(\beta+3)$

$(\alpha - \beta)^2 = 3^2$ $\quad$ $\alpha = \beta + 3$ (where α is greater)

$(\alpha+\beta)^2 - 4\alpha\beta = 9$ $\quad$ $\beta(\beta+3) = \frac{7(1+k)}{3}$

$\left(\frac{2}{3}\right)^2 - 4\left(\frac{7(1+k)}{3}\right) = 9$ $\quad$ $\beta^2 + 3\beta = \frac{7+7k}{3}$

$\frac{4}{9} - \frac{28(1+k)}{3} = 9$

$\frac{4 - 84(1+k)}{9} = \frac{9}{1}$

$4 - 84 - 84k = 81$

$-80 - 84k = 81$

$-84k = 81 + 80$

$-84k = 161$

$k = -\frac{161}{84}$

$\beta + \beta + 3 = \frac{2}{3}$

$2\beta + 3 = \frac{2}{3}$

$2\beta = \frac{2}{3} - 3$

$2\beta = \frac{2-9}{3}$

$2\beta = -\frac{7}{3}$

$\beta = -\frac{7}{6}$

$\beta(\beta+3) = \frac{7(1+k)}{3}$

$\left(-\frac{7}{6}\right)\left(-\frac{7}{6}+3\right) = \frac{7(1+k)}{3}$

$\left(-\frac{7}{6}\right)\left(\frac{-7+18}{6}\right) = \frac{7(1+k)}{3}$

$\left(-\frac{7}{6}\right)\left(\frac{11}{6}\right) = \frac{7(1+k)}{3}$

$-\frac{77}{36} = \frac{7(1+k)}{3}$

$\frac{-77}{36} = \frac{7+7k}{3}$

$-77 = 84 + 84k$

$-77 - 84 = 84k$

$-161 = 84k$

$k = -\frac{161}{84}$

For $\alpha = \beta + 3$ we get;

$\alpha = -\frac{7}{6} + 3$

$\alpha = \frac{-7+18}{6}$

$\alpha = \frac{11}{6}$

For $\alpha = \beta$ we get;

$\beta = \beta + 3$

$0 = 3$

$\alpha \neq \beta$

(iv) The roots of the equation $3x^2 - 2x + 7(1+k) = 0$ differ by their squares.

Solution:

$3x^2 - 2x + 7(1+k) = 0$

Here $a=3, b=-2, c=7(1+k)$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{-2}{3} = \frac{2}{3}$ and $\alpha\beta = \frac{c}{a} = \frac{7(1+k)}{3}$

Now $\alpha - \beta = \alpha^2 - \beta^2$ $\quad$ and $\alpha\beta = \beta\alpha$ $\quad$ ..... (i)

Now $\alpha - \beta = (\alpha+\beta)(\alpha-\beta)$ $\quad$ ..... (ii)

Eq. (i) = Eq. (ii)

$(\alpha+\beta)(\alpha-\beta) = \alpha - \beta$

$(\alpha+\beta)(\alpha-\beta) - (\alpha-\beta) = 0$

$(\alpha-\beta)(\alpha+\beta-1) = 0$

$\alpha - \beta = 0$ $\quad$ or $\quad$ $\alpha + \beta - 1 = 0$

$\alpha = \beta$ $\quad$ $\alpha + \beta = 1$

$\alpha + \beta = \frac{2}{3}$

$\frac{2}{3} \neq 1$

$\alpha \neq \beta$

For $\alpha = \beta$ we get;

$\beta = 1 - \beta$

$\beta + \beta = 1$

$2\beta = 1$

$\beta = \frac{1}{2}$

For $\alpha = \beta$ and $\beta = 1 - \alpha$ we get;

$\alpha = 1 - \alpha$

$\alpha + \alpha = 1$

$2\alpha = 1$

$\alpha = \frac{1}{2}$

$\beta = 1 - \alpha$

$\beta = 1 - \frac{1}{2}$

$\beta = \frac{2-1}{2}$

$\beta = \frac{1}{2}$

$\alpha = \beta$

(v) The roots of the equation $3x^2 - 2x + 7(1+k) = 0$ are reciprocal to each other.

Solution:

$3x^2 - 2x + 7(1+k) = 0$

Here $a=3, b=-2, c=7(1+k)$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{-2}{3} = \frac{2}{3}$ and $\alpha\beta = \frac{c}{a} = \frac{7(1+k)}{3}$

$\alpha = \frac{1}{\beta}$ $\quad$ and $\quad$ $\alpha\beta = 1$ $\quad$ ..... (i)

$\alpha\beta = \beta\alpha$ $\quad$ ..... (ii)

Eq. (i) = Eq. (ii)

$\alpha\beta = 1$

$\frac{7(1+k)}{3} = 1$

$7 + 7k = 3$

$7k = 3 - 7$

$7k = -4$

$k = -\frac{4}{7}$

(vi) The roots of the equation $3x^2 - 2x + 7(1+k) = 0$ are negative reciprocal to each other.

Solution:

$3x^2 - 2x + 7(1+k) = 0$

Here $a=3, b=-2, c=7(1+k)$

Let α and β be the roots of the given equation

Then Sum of roots $= \alpha + \beta = -\frac{b}{a} = -\frac{-2}{3} = \frac{2}{3}$ and $\alpha\beta = \frac{c}{a} = \frac{7(1+k)}{3}$

$\alpha = -\frac{1}{\beta}$ $\quad$ and $\quad$ $\alpha\beta = -1$ $\quad$ ..... (i)

$\alpha\beta = \beta\alpha$ $\quad$ ..... (ii)

Eq. (i) = Eq. (ii)

$\alpha\beta = -1$

$\frac{7(1+k)}{3} = -1$

$7 + 7k = -3$

$7k = -3 - 7$

$7k = -10$

$k = -\frac{10}{7}$

5. Find a, if one of the roots of the following equations is equal to a given constant.

(i) $(2a-3)x^2 + (7a-5)x + (2a-4) = 0$

Solution:

$(2a-3)x^2 + (7a-5)x + (2a-4) = 0$

Here $a=(2a-3), b=(7a-5), c=(2a-4)$

Let α and β be the roots of the given equation

Then $\alpha + \beta = -\frac{b}{a} = -\frac{(7a-5)}{(2a-3)}$ $\quad$ and $\quad$ $\alpha\beta = \frac{c}{a} = \frac{(2a-4)}{(2a-3)}$

$\alpha + \beta = \frac{-7a+5}{2a-3}$ $\quad$ ..... (i)

$\alpha\beta = \frac{2a-4}{2a-3}$ $\quad$ ..... (ii)

Comparing eq.(i) with eq.(ii) we get

$\frac{-7a+5}{2a-3} = \frac{2a-4}{2a-3}$

$-7a + 5 = 2a - 4$

$-7a - 2a = -4 - 5$

$-9a = -9$

$a = \frac{-9}{-9}$

$a = 1$

(ii) $4x^2 - (2+5a)x + (5a-7) = 0$

Solution:

$4x^2 - (2+5a)x + (5a-7) = 0$

Here $a=4, b=-(2+5a), c=(5a-7)$

Let α and β be the roots of the given equation

Then $\alpha + \beta = -\frac{b}{a} = -\frac{-(2+5a)}{4} = \frac{(2+5a)}{4}$ $\quad$ and $\quad$ $\alpha\beta = \frac{c}{a} = \frac{(5a-7)}{4}$

$\alpha + \beta = \frac{2+5a}{4}$ $\quad$ ..... (i)

$\alpha\beta = \frac{5a-7}{4}$ $\quad$ ..... (ii)

Comparing eq.(i) with eq.(ii) we get

$\frac{2+5a}{4} = \frac{5a-7}{4}$

$2 + 5a = 5a - 7$

$5a - 5a = -7 - 2$

$0 = -9$

$a \neq \beta$

(iii) $4x^2 - (2+5a)x + (5a-7) = 0$

Solution:

$4x^2 - (2+5a)x + (5a-7) = 0$

Here $a=4, b=-(2+5a), c=(5a-7)$

Let α and β be the roots of the given equation

Then $\alpha + \beta = -\frac{b}{a} = -\frac{-(2+5a)}{4} = \frac{(2+5a)}{4}$ $\quad$ and $\quad$ $\alpha\beta = \frac{c}{a} = \frac{(5a-7)}{4}$

$\alpha + \beta = \frac{2+5a}{4}$ $\quad$ ..... (i)

$\alpha\beta = \frac{5a-7}{4}$ $\quad$ ..... (ii)

Comparing eq.(i) with eq.(ii) we get

$\frac{2+5a}{4} = \frac{5a-7}{4}$

$2 + 5a = 5a - 7$

$5a - 5a = -7 - 2$

$0 = -9$

$\alpha \neq \beta$

Symmetric functions of the roots of a quadratic equation:
Define symmetric functions of the roots of a quadratic equation:
Definition:
Symmetric functions are those functions in which the roots involved are such that the value of the expressions involving them remain unaltered, when roots are interchanged.

For example, If

$$f(\alpha,\beta)=\alpha^2+\beta^2, \text{ then}$$
$$f(\beta,\alpha)=\beta^2+\alpha^2=\alpha^2+\beta^2 \qquad (\because \beta^2+\alpha^2=\alpha^2+\beta^2)$$
$$=f(\alpha,\beta)$$

Exercise 2.4

1. If α , β are the roots of the equation x²+px+q=0, then evaluate

(i) $\alpha^2+\beta^2$

Solution:
Solution:
$x^2+px+q=0$

Here a=1, b=p, c=q

Let α , β be the roots of the given equation

Then $\alpha+\beta=-\frac{b}{a}$ *and* $\alpha\beta=\frac{c}{a}$

$=-\frac{p}{1}$ $\alpha\beta=\frac{q}{1}$

$=-p$ $=q$

Now $\alpha^2+\beta^2=(\alpha+\beta)^2-2\alpha\beta$

$=(-p)^2-2(q)$

$=p^2-2q$

(ii) $\alpha^3\beta+\alpha\beta^3$

Solution:
$\alpha^3\beta+\alpha\beta^3$
$x^2+px+q=0$
Here a=1, b=p, c=q

Let α , β be the roots of the given equation

Then $\alpha+\beta=-\frac{b}{a}$ *and* $\alpha\beta=\frac{c}{a}$

$=-\frac{p}{1}$ $\alpha\beta=\frac{q}{1}$

$=-p$ $=q$

Now $\alpha^3\beta+\alpha\beta^3=\alpha\beta(\alpha^2+\beta^2)$

$=\alpha\beta\left[(\alpha+\beta)^2-2\alpha\beta\right]$

$=q\left[(-p)^2-2q\right]$

$=q\left(p^2-2q\right)$

(iii) $\frac{\alpha}{\beta}+\frac{\beta}{\alpha}$

Solution:
$\frac{\alpha}{\beta}+\frac{\beta}{\alpha}$
$x^2+px+q=0$

Here a=1, b=p, c=q

Let α , β be the roots of the given equation

Then $\alpha+\beta=-\frac{b}{a}$ *and* $\alpha\beta=\frac{c}{a}$

$=-\frac{p}{1}$ $\alpha\beta=\frac{q}{1}$

$=-p$ $=q$

Now $\frac{\alpha}{\beta}+\frac{\beta}{\alpha}=\frac{\alpha^2+\beta^2}{\alpha\beta}$

$=\frac{(\alpha+\beta)^2-2\alpha\beta}{\alpha\beta}$

$=\frac{(-p)^2-2q}{q}=\frac{1}{q}\left(p^2-2q\right)$

2. If α , β are the roots of the equation 4x²-5x+6=0, then find the values of

(i) $\frac{1}{\alpha}+\frac{1}{\beta}$

Solution:
$\frac{1}{\alpha}+\frac{1}{\beta}$
$4x^2-5x+6=0$
Here a=4, b=-5, c=6

Let α , β be the roots of the given equation

Then $\alpha+\beta=-\frac{b}{a}$ *and* $\alpha\beta=\frac{c}{a}$

$=-\frac{(-5)}{4}$ $\alpha\beta=\frac{6}{4}$

$=\frac{5}{4}$ $=\frac{3}{2}$

Now $\frac{1}{\alpha}+\frac{1}{\beta}=\frac{\alpha+\beta}{\alpha\beta}$

$=\frac{5}{\frac{3}{2}}=\frac{5}{4}\times\frac{2}{3}=\frac{5}{6}$

(ii) $\alpha^2\beta^2$

Solution:
$\alpha^2\beta^2$
$4x^2-5x+6=0$
Here a=4, b=-5, c=6

Let α , β be the roots of the given equation

Then $\alpha+\beta=-\frac{b}{a}$ *and* $\alpha\beta=\frac{c}{a}$

$=-\frac{(-5)}{4}$ $\alpha\beta=\frac{6}{4}$

$=\frac{5}{4}$ $=\frac{3}{2}$

Now $\alpha^2\beta^2=(\alpha\beta)^2=\left(\frac{3}{2}\right)^2=\frac{9}{4}$

(iii) $\frac{1}{\alpha^2\beta}+\frac{1}{\alpha\beta^2}$

Solution:
$4x^2-5x+6=0$
Here a=4, b=-5, c=6

Let α , β be the roots of the given equation

Then $\alpha+\beta=-\frac{b}{a}$ *and* $\alpha\beta=\frac{c}{a}$

$=-\frac{(-5)}{4}$ $\alpha\beta=\frac{6}{4}$

$=\frac{5}{4}$ $=\frac{3}{2}$

Now $\frac{1}{\alpha^2\beta}+\frac{1}{\alpha\beta^2}=\frac{\alpha+\beta}{\alpha^2\beta^2}=\frac{\alpha+\beta}{(\alpha\beta)^2}$

$=\frac{\frac{5}{4}}{\left(\frac{3}{2}\right)^2}=\frac{5}{9}\times\frac{4}{9}=\frac{5}{9}$

(iv) $\frac{\alpha^2}{\beta}+\frac{\beta^2}{\alpha}$

Solution:
$4x^2-5x+6=0$
Here a=4, b=-5, c=6

Let α , β be the roots of the given equation

Then $\alpha+\beta=-\frac{b}{a}$ *and* $\alpha\beta=\frac{c}{a}$

$=-\frac{(-5)}{4}$ $\alpha\beta=\frac{6}{4}$

$=\frac{5}{4}$ $=\frac{3}{2}$

Now $\frac{\alpha^3}{\beta}+\frac{\beta^3}{\alpha}=\frac{\alpha^4+\beta^4}{\alpha\beta}=\frac{(\alpha+\beta)^2-3\alpha\beta((\alpha+\beta)}{\alpha\beta}$

$=\frac{\left(\frac{5}{4}\right)^2-\frac{3}{2}\left(\frac{5}{4}\right)}{\frac{3}{2}}=\frac{\frac{125}{64}}{\frac{3}{2}}$

$=\frac{125-360}{64}\times\frac{2}{3}=-\frac{235}{64}\times\frac{2}{3}$

$=\frac{235}{96}$

3. If α , β are the roots of the equation lx²+mx+n=0 $(l\neq 0)$**, then find the values of:**

(i) $\alpha^2\beta^2+\alpha^2\beta^3$

Solution:
$lx^2+mx+n=0$
Here a=l, b=m, c=n

Let α , β be the roots of the given equation

Then $\alpha+\beta=-\frac{b}{a}$ *and* $\alpha\beta=\frac{c}{a}$

$=-\frac{m}{l}$ $\alpha\beta=\frac{n}{l}$

Now $\alpha^2\beta^2+\alpha^2\beta^3=\alpha^2\beta^2(\alpha+\beta)=(\alpha\beta)^2(\alpha+\beta)$

$=\left(\frac{n}{l}\right)^2\left(-\frac{m}{l}\right)=-\frac{mn^2}{l^3}$

(ii) $\frac{1}{\alpha^2}+\frac{1}{\beta^2}$

Solution:
$lx^2+mx+n=0$
Here a=l, b=m, c=n

Let α , β be the roots of the given equation

Then $\alpha+\beta=-\frac{b}{a}$ *and* $\alpha\beta=\frac{c}{a}$

$=-\frac{m}{l}$ $\alpha\beta=\frac{n}{l}$

Now $\frac{1}{\alpha^2}+\frac{1}{\beta^2}=\frac{\alpha^2+\beta^2}{\alpha^2\beta^2}=\frac{(\alpha+\beta)^2-2\alpha\beta}{(\alpha\beta)^2}$

$=\frac{\left(-\frac{m}{l}\right)^2-2\left(\frac{n}{l}\right)}{\left(\frac{n}{l}\right)^2}=\frac{\frac{m^2}{l^2}-\frac{2n}{l}}{\frac{n^2}{l^2}}$

$=\frac{m^2-2nl}{l^2}\times\frac{l^2}{n^2}=\frac{m^2-2nl}{n^2}$

$=\frac{1}{n^2}\left[m^2-2nl\right]$

Formation of a quadratic equation:
If α and β are the roots of the required quadratic equation.

Let $\quad \alpha + \alpha \quad$ and $\quad \alpha + \beta$
i.e., $\quad \alpha + \alpha = 0, \quad \alpha \cdot \beta = 0$
and $\quad (x - \alpha)(x - \beta) = 0$
$\Rightarrow x^2 - (\alpha + \beta)x + \alpha\beta = 0$
which is the required quadratic equation in standard form.

Find a quadratic equation from given roots and establish the formula

x^2 - (sum of the roots) x + product of the roots = 0

Let $\quad \alpha, \beta$ be the roots of the given equation
$\alpha x^2 + bx + c = 0 \quad \ldots (i) \quad (a \neq 0) \quad \ldots \ldots (ii)$
Then $\quad \alpha + \beta = -\frac{b}{a} \quad$ and $\quad \alpha\beta = \frac{c}{a}$
Rewrite eq.(i) as $\quad x^2 + \frac{b}{a}x + \frac{c}{a} = 0$
or $\quad x^2 - \left(-\frac{b}{a}\right)x + \frac{c}{a} = 0$
or $\quad x^2 - (\alpha + \beta)x + \alpha\beta = 0$
or $\quad x^2 -$ (sum of the roots) x + product of the roots =0, that is
$x^2 - Sx + P = 0$ where S= $\alpha + \beta$ and P= $\alpha\beta$

Exercise 2.5

1. Write the quadratic equations having following roots.
(a) 1 3
Solution:
Since 1 and 3 are the roots of the required quadratic equation, therefore,

S = Sum of roots = 1 + 3 = 4
P = Product of roots = (1) (3) = 3
Thus, the required quadratic equation is
x^2 - Sx + P = 0
$x^2 - 4x + 3 = 0$

(b) 4 5
Solution:
Since 4 and 5 are the roots of the required quadratic equation, therefore,
S = Sum of roots = 4 + 5 = 9
P = Product of roots = (4) (5) = 36
Thus, the required quadratic equation is
x^2 - Sx + P = 0
$x^2 - 9x + 36 = 0$

(c) -3 3
Solution:
Since -3 and 3 are the roots of the required quadratic equation, therefore,
S = Sum of roots = -3 + 3 = 0
P = Product of roots = (-3) (3) = -9
Thus, the required quadratic equation is
x^2 - Sx + P = 0
$x^2 - (0)x + (-9) = 0$
$x^2 - 9 = 0$

(d) 5 - 6
Solution:
Since 5 and -6 are the roots of the required quadratic equation, therefore,
S = Sum of roots = 5 + (-6) = 5 - 6 = -1
P = Product of roots = (5) (-6) = -30
Thus, the required quadratic equation is
x^2 - Sx + P = 0
$x^2 - (-1)x + (-30) = 0$
$x^2 + x - 30 = 0$

Solution:
Since 0 and - 3 are the roots of the required quadratic equation, therefore,
S = Sum of roots = 0 + (-3) = 0 - 3 = -3
P = Product of roots = (0) (-3) = 0
Thus, the required quadratic equation is
x^2 - Sx + P = 0
$x^2 - (-3)x + 0 = 0$
$x^2 + 3x = 0$

(e) 3 - 1 + i
Solution:
Since 3 and - i are the roots of the required quadratic equation, therefore,
S = Sum of roots = 3 + (-i) = 3 - i
P = Product of roots = (3) (-i) = -3i
Thus, the required quadratic equation is
x^2 - Sx + P = 0
$x^2 - (3 - i)x + (-3i) = 0$
$x^2 - 3x + ix - 3i = 0$

(f) -1 - i 7
Solution:
Since -1 and - 7 are the roots of the required quadratic equation, therefore,
S = Sum of roots = (-1) + (-7) = -1 - 7 = -8
P = Product of roots = (-1) (-7) = 7
Thus, the required quadratic equation is
x^2 - Sx + P = 0

$x^2 - (-8)x + 7 = 0$
$x^2 + 8x + 7 = 0$

(g) 3+i 1 + i
Solution:
Since 1+ i and 1 − i are the roots of the required quadratic equation, therefore,
S = Sum of roots = (1+ i) + (1 − i) = 1 + i + 1 − i = 2
P = Product of roots = (1+ i) (1 − i) = 1 − i + i − i = 1 − (−1) = 2
Thus, the required quadratic equation is
x^2 - Sx + P = 0
$x^2 - 2x + 2 = 0$

(h) 2+ $\sqrt{3}$, 2- $\sqrt{3}$
Solution:
Since 2+ $\sqrt{3}$ and 2- $\sqrt{3}$ are the roots of the required quadratic equation, therefore,
S = Sum of roots = $2 + \sqrt{3} + 2 - \sqrt{3} = 4$
P = Product of roots = $(2 + \sqrt{3})(2 - \sqrt{3}) = (2)^2 - (\sqrt{3})^2 = 4 - 3 = 1$
Thus, the required quadratic equation is
x^2 - Sx + P = 0
$x^2 - 4x + 1 = 0$

2. If α, β are the roots of the equation $x^2 - 3x + 6 = 0$
form equations whose roots are

a) $2\alpha + 1, 2\beta + 1$
Solution:

$x^2 - 3x + 6 = 0$
Here $a = 1, b = -3, c = 6$.

Let α, β be the roots of the given equation
Then $\quad \alpha + \beta = -\frac{b}{a} \quad$ and $\quad \alpha\beta = \frac{c}{a}$
$= -\frac{(-3)}{1} \qquad\qquad = \frac{6}{1}$
$= 3 \qquad\qquad = 6$

Let roots of the new equation be $2\alpha + 1, 2\beta + 1$.

S = Sum of roots	P = Product of roots
$= (2\alpha + 1) + (2\beta + 1)$	$= (2\alpha + 1)(2\beta + 1)$
$= 2\alpha + 1 + 2\beta + 1$	$= (2\alpha + 1)(2\beta + 1)$
$= 2\alpha + 2\beta + 2$	$= 4\alpha\beta + 2\alpha + 2\beta + 1$
$= 2(\alpha + \beta) + 2$	$= 4\alpha\beta + 2(\alpha + \beta) + 1$
$= 2(3) + 2$	$= 4(6) + 2(3) + 1$
$= 6 + 2$	$= 24 + 6 + 1$
$= 8$	$= 31$

Thus, the required quadratic equation is
$x^2 - Sx + P = 0$
$x^2 - (8)x + 31 = 0$
$x^2 - 8x + 31 = 0$

b) α^2, β^2
Solution:

$x^2 - 3x + 6 = 0$
Here $a = 1, b = -3, c = 6$.

Let α, β be the roots of the given equation
Then $\quad \alpha + \beta = -\frac{b}{a} \quad$ and $\quad \alpha\beta = \frac{c}{a}$
$= -\frac{(-3)}{1} \qquad\qquad = \frac{6}{1}$
$= 3 \qquad\qquad = 6$

Let roots of the new equation be α^2, β^2.

S = Sum of roots	P = Product of roots
$= \alpha^2 + \beta^2$	$= (\alpha^2)(\beta^2)$
$= (\alpha + \beta)^2 - 2\alpha\beta$	$= (\alpha\beta)^2$
$= (3)^2 - 2(6)$	$= (6)^2$
$= 9 - 12$	$= 36$
$= -3$	

Thus, the required quadratic equation is
$x^2 - Sx + P = 0$
$x^2 - (-3)x + 36 = 0$
$x^2 + 3x + 36 = 0$

c) $\frac{1}{\alpha}, \frac{1}{\beta}$
Solution:
$x^2 - 3x + 6 = 0$
Here $a = 1, b = -3, c = 6$

Let α, β be the roots of the given equation
Then $\quad \alpha + \beta = -\frac{b}{a} \quad$ and $\quad \alpha\beta = \frac{c}{a}$
$= -\frac{(-3)}{1} \qquad\qquad = \frac{6}{1}$
$= 3 \qquad\qquad = 6$

Let roots of the new equation be $\frac{1}{\alpha}, \frac{1}{\beta}$.

S = Sum of roots	P = Product of roots
$= \frac{1}{\alpha} + \frac{1}{\beta}$	$= \left(\frac{1}{\alpha}\right)\left(\frac{1}{\beta}\right)$
$= \frac{\alpha + \beta}{\alpha\beta}$	$= \frac{1}{\alpha\beta}$
$= \frac{3}{6}$	$= \frac{1}{6}$

Thus, the required quadratic equation is
$x^2 - Sx + P = 0$
$x^2 - \left(\frac{1}{2}\right)x + \frac{1}{6} = 0$
$6x^2 - 3x + 1 = 0$

d) $\frac{\alpha}{\beta}, \frac{\beta}{\alpha}$
Solution:
$x^2 - 3x + 6 = 0$
Here $a = 1, b = -3, c = 6$.

Let α, β be the roots of the given equation
Then $\quad \alpha + \beta = -\frac{b}{a} \quad$ and $\quad \alpha\beta = \frac{c}{a}$
$= -\frac{(-3)}{1} \qquad\qquad = \frac{6}{1}$
$= 3 \qquad\qquad = 6$

Let roots of the new equation be $\frac{\alpha}{\beta}, \frac{\beta}{\alpha}$.

S = Sum of roots	P = Product of roots
$= \frac{\alpha}{\beta} + \frac{\beta}{\alpha}$	$= \left(\frac{\alpha}{\beta}\right)\left(\frac{\beta}{\alpha}\right)$
$= \frac{\alpha^2 + \beta^2}{\alpha\beta}$	$= 1$
$= \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta}$	
$= \frac{(3)^2 - 2(6)}{6}$	
$= \frac{9 - 12}{6}$	
$= \frac{-3}{6}$	
$= -\frac{1}{2}$	

Thus, the required quadratic equation is

$x^2 - Sx + P = 0$
$x^2 - \left(-\frac{1}{2}\right)x + 1 = 0$
$2x^2 + x + 2 = 0$

e) $\alpha + \frac{1}{\beta}, \beta + \frac{1}{\alpha}$
Solution:
$x^2 - 3x + 6 = 0$
Here $a = 1, b = -3, c = 6$.

Let α, β be the roots of the given equation
Then $\quad \alpha + \beta = -\frac{b}{a} \quad$ and $\quad \alpha\beta = \frac{c}{a}$
$= -\frac{(-3)}{1} \qquad\qquad = \frac{6}{1}$
$= 3 \qquad\qquad = 6$

Let roots of the new equation be $\alpha + \frac{1}{\beta}, \beta + \frac{1}{\alpha}$ P = Product of roots

S = Sum of roots	
$= \alpha + \beta + \frac{1}{\alpha} + \frac{1}{\beta}$	$= (\alpha + \beta)\left(\frac{1}{\alpha} + \frac{1}{\beta}\right)$
$= (\alpha + \beta)\left(\frac{\alpha + \beta}{\alpha\beta}\right)$	$= (\alpha + \beta)\left(\frac{\alpha + \beta}{\alpha\beta}\right)$
$= 3 + \frac{3}{6}$	$= 1 + \left(\frac{3}{6}\right)^2$
$= 3 + \frac{1}{2}$	$= 1 + \frac{9}{36}$
$= \frac{7}{2}$	$= \frac{5}{4}$

Thus, the required quadratic equation is

3. If α, β are the roots of the equation $x^2 + px + q = 0$
form equations whose roots are

a) α^2, β^2
Solution:
$x^2 + px + q = 0$
Here $a = 1, b = p, c = q$.

Let α, β be the roots of the given equation
Then $\quad \alpha + \beta = -\frac{b}{a} \quad$ and $\quad \alpha\beta = \frac{c}{a}$
$= -\frac{p}{1} \qquad\qquad = \frac{q}{1}$
$= -p \qquad\qquad = q$

Let roots of the new equation be α^2, β^2.

S = Sum of roots	P = Product of roots
$= \alpha^2 + \beta^2$	$= (\alpha^2)(\beta^2)$
$= (\alpha + \beta)^2 - 2\alpha\beta$	$= (\alpha\beta)^2$
$= (-p)^2 - 2q$	$= q^2$
$= p^2 - 2q$	

Thus, the required quadratic equation is
$x^2 - Sx + P = 0$
$x^2 - (p^2 - 2q)x + q^2 = 0$

b) $\frac{\alpha}{\beta}, \frac{\beta}{\alpha}$
Solution:
$x^2 + px + q = 0$
Here $a = 1, b = p, c = q$

Let α, β be the roots of the given equation

Then $\quad \alpha + \beta = -\frac{b}{a} \quad$ and $\quad \alpha\beta = \frac{c}{a}$

$= -\frac{p}{1} \qquad\qquad = \frac{q}{1}$
$= -p \qquad\qquad = q$

Let roots of the new equation be $\frac{\alpha}{\beta}, \frac{\beta}{\alpha}$.

S = Sum of roots	P = Product of roots
$= \frac{\alpha}{\beta} + \frac{\beta}{\alpha}$	$= \left(\frac{\alpha}{\beta}\right)\left(\frac{\beta}{\alpha}\right)$
$= \frac{\alpha^2 + \beta^2}{\alpha\beta}$	$= 1$
$= \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta}$	
$= \frac{(-p)^2 - 2q}{q}$	
$= \frac{p^2 - 2q}{q}$	

Thus, the required quadratic equation is
$x^2 - Sx + P = 0$
$x^2 - \left(\frac{p^2 - 2q}{q}\right)x + 1 = 0$
$qx^2 - (p^2 - 2q)x + q = 0$

Synthetic Division:
Synthetic division is the process of finding the *quotient* and *remainder*, when a polynomial is divided by a linear polynomial. In fact, synthetic division is simply a shortcut of long division method.

Exercise 2.6

1. Use synthetic division to find the quotient and the remainder, when

(i) $\left(x^3-7x-1\right)\div\left(x+1\right)$

Solution:

$\left(x^3-7x-1\right)\div\left(x+1\right)$

As $x+1=x-(-1)$, so $a=-1$

Now write the coefficients of dividend in a row and $a=-1$ on the left side,,

$$\begin{array}{r|rrrr} -1 & 1 & 0 & -7 & -1 \\ & & -1 & 1 & 6 \\ \hline & 1 & -1 & -6 & 5 \end{array}$$

Quotient $=x^2-x-6$

Remainder $=5$

(ii) $\left(6x^3-5x+15\right)\div\left(x+3\right)$

Solution:

$\left(6x^3-5x+15\right)\div\left(x+3\right)$

As $x+3=x-(-3)$, so $a=-3$

Now write the coefficients of dividend in a row and $a=-3$ on the left side.

$$\begin{array}{r|rrrr} -3 & 6 & 0 & -5 & 15 \\ & & -18 & 54 & -147 \\ \hline & 6 & -18 & 49 & -132 \end{array}$$

Quotient $=6x^2-18x+49$

Remainder $=-132$

(iii) $\left(6x^3+x^2-3x+2\right)\div\left(x-2\right)$

Solution:

$\left(6x^3+x^2-3x+2\right)\div\left(x-2\right)$

As $x-2=x+(-2)$, so $a=2$

Now write the coefficients of dividend in a row and $a=2$ on the left side.

$$\begin{array}{r|rrrr} 2 & 6 & 1 & -3 & 2 \\ & & 12 & 26 & 46 \\ \hline & 6 & 13 & 23 & 48 \end{array}$$

Quotient $=6x^2+13x+23$

Remainder $=48$

2. Find the value of k using synthetic division, if

(i) 3 is the zero of the polynomial $2x^3-3kx^2+9$

Solution:

$P(x)=2x^3-3kx^2+9$ and 3 is zero is 3.

Then by synthetic division,

$$\begin{array}{r|rrrr} 3 & 2 & -3k & 0 & 9 \\ & & 6 & 18-9k & 54-27k \\ \hline & 2 & 6-3k & 18-9k & 63-27k \end{array}$$

Remainder $=63-27k$

Since 3 is the zero of the polynomial, therefore

Remainder $=0$, that is,

$$\begin{aligned} 63-27k&=0\\ 27k&=63\\ k&=\frac{63}{27}\\ k&=\frac{7}{3} \end{aligned}$$

(ii) 1 is the zero of the polynomial $3^{x^3}-2kx^2+11$

Solution:

$P(x)=x^3-2kx^2+11$ and its zero is 1.

Then by synthetic division.

$$\begin{array}{r|rrrr} 1 & 3 & -2k & 0 & 11 \\ & & 3 & 3-2k & 3-2k \\ \hline & 3 & 3-2k & 3-2k & 13-2k \end{array}$$

Remainder $=23-2k$

Since 1 is the zero of the polynomial, therefore

Remainder $=0$, that is

$$\begin{aligned} 23-2k&=0\\ 2k&=23\\ k&=\frac{23}{2}=k \end{aligned}$$

(iii) 1 is the zero of the polynomial $2x^3+5kx-23$.

Solution:

$P(x)=2x^3+5kx-23$ and its zero is 1.

Then by synthetic division.

$$\begin{array}{r|rrrr} 1 & 2 & 0 & 5k & -23 \\ & & 2 & 2 & 5k+2 \\ \hline & 2 & 2 & 5k+2 & 5k-23 \end{array}$$

Remainder $=5k-23$

Since 1 is the zero of the polynomial, therefore

Remainder $=0$, that is

$$\begin{aligned} 5k-23&=0\\ 5k&=23\\ k&=\frac{23}{5}\\ k&=5\frac{3}{5} \end{aligned}$$

3. Use synthetic division to find the values of l and m, if

(i) (x+3) and (x-2) are the factors of the polynomial $x^3-4x^2+2lx+m$

Solution:

Since $(x+3)$ and $(x-2)$ are the factors of $p(x)=x^3+4x^2+2lx+m$. Therefore 2 and 3 are zeroes of polynomial $p(x)$.

Now by synthetic division,

$$\begin{array}{r|rrrr} 2 & 1 & -4 & 2l & m \\ & & 2 & -4 & 4l-8 \\ \hline & 1 & -2 & 2l-4 & m+4l-8 \end{array}$$

Remainder $=m-4l+8$

Since -2 is the zero of the polynomial, therefore,

Remainder $=0$, that is,

$m-4l+8=0$ ……(i)

and

$$\begin{array}{r|rrrr} -3 & 1 & -4 & 2l & m \\ & & -3 & 21 & -6l+36 \\ \hline & 1 & -7 & 2l+21 & m-6l+36 \end{array}$$

Remainder $=m-6l+36$

Since -3 is the zero of the polynomial, therefore,

Remainder $=0$, that is,

$m-6l+36=0$ ……(ii)

$$\begin{aligned} m-4l+8&=0\\ \frac{(m=6l+36)}{}&=0\\ -2l-28&=0\\ -2l&=28\\ l&=\frac{28}{-2}\\ l&=-\frac{14}{1} \end{aligned}$$

Put $l=-\frac{14}{1}$ in eq.(i), we get

$$\begin{aligned} m-4\left(-\frac{14}{1}\right)+8&=0\\ m+\frac{56}{1}+8&=0\\ m+64&=0\\ m&=-64 \end{aligned}$$

Thus, $l=-\frac{14}{1}$ & $m=-64$

(ii) (x-1) and (x+1) are the factors of the polynomial $x^3+3lx^2+2mx+8$.

Solution:

Since $(x-1)$ and $(x+1)$ are the factors of $p(x)=x^3+3lx^2+2mx+8$. Therefore 1 and -1 are zeroes of polynomial $p(x)$.

Now by synthetic division,

$$\begin{array}{r|rrrr} 1 & 1 & 3l & 2m & 8 \\ & & 1 & 3l+1 & 2m+3l+1 \\ \hline & 1 & 3l+1 & 2m+3l+1 & 2m+3l+9 \end{array}$$

Remainder $=2m+3l+9$

Since 1 is the zero of the polynomial, therefore,

Remainder $=0$, that is,

$2m+3l+9=0$ ……(i)

Again by synthetic division,

$$\begin{array}{r|rrrr} -1 & 1 & 3l & 2m & 8 \\ & & -1 & 1-3l & -2m+3l-1 \\ \hline & 1 & 3l-1 & 2m-3l+1 & -2m+3l+7 \end{array}$$

Remainder $=-2m+3l+7$

Since -1 is the zero of the polynomial, therefore,

Remainder $=0$, that is,

$-2m+3l+7=0$ ……(ii)

and

$$\begin{array}{r|rrrr} -1 & 1 & 3l & 2m & 8 \\ & & -1 & 3l+1 & -2m-3l-1 \\ \hline & 1 & 3l-1 & 2m+3l+1 & -2m-3l+7 \end{array}$$

Remainder $=-2m-3l+7$

Since -1 is the zero of the polynomial, therefore,

Remainder $=0$, that is ,

$-2m-3l+7=0$ ……(ii)

Add eq(i) in eq(ii), we get

$$\begin{aligned} 2m-3l+7&=0\\ \frac{-2m+3l+9=0}{}&\\ 16&=0\\ m+1&=0\\ m&=\frac{16}{1}\\ m&=-1 \end{aligned}$$

Put $l=2$ in eq.(i), we get

$$\begin{aligned} 2m+3(2)+7&=0\\ 2m+6+7&=0\\ 2m+13&=0\\ 2m&=-1\\ m&=-\frac{1}{2} \end{aligned}$$

Thus $l=2,m=-\frac{1}{2}$

4. Solve by using synthetic division, if

(i) 3 is the root of the equation $x^3-28x+48=0$

Solution:

Since 3 is the root of the equation $x^3-28x+48=0$

Then by synthetic division, we get

$$\begin{array}{r|rrrr} 3 & 1 & 0 & -28 & 48 \\ & & 3 & 9 & -57 \\ \hline & 1 & 3 & -19 & -9 \end{array}$$

The depressed equation is

$x^2+3x-19=0$

$x^2+3x-19=0$

$x(x+3)-6(x+3)=0$

$(x-6)(x+3)=0$

Hence, $-6,3$ and 4 are the roots of the given equation.

(ii) 3 is the root of the equation $2x^3-3x^2-11x+6=0$.

Solution:

Since 3 is the root of the equation $2x^3-3x^2-11x+6=0$

Then by synthetic division, we get

$$\begin{array}{r|rrrr} 3 & 2 & -3 & -11 & 6 \\ & & 6 & 9 & -6 \\ \hline & 2 & 3 & -2 & 0 \end{array}$$

The depressed equation is

$2x^2+3x-2=0$

$6x^2-4x+3x-2=0$

$2x(3x-1)+1(3x-1)=0$

$(2x+1)(3x-1)=0$

Either $2x+1=0$ or $3x-1=0$

Either $x=-\frac{1}{2}$ or $x=\frac{1}{3}$

Hence, $-\frac{1}{2}$ and $\frac{1}{3}$ are the roots of the given equation.

Classmate, Inc. © 2020

(iii) -1 is the root of the equation $6x^3-x^2-11x+6=0$.

Solution:

Since -1 is the root of the equation $6x^3-x^2-11x+6=0$

Then by synthetic division, we get

$$\begin{array}{r|rrrr} -1 & 6 & -1 & -11 & 6 \\ & & -6 & 7 & 4 \\ \hline & 6 & -7 & -4 & 10 \end{array}$$

The depressed equation is

$6x^2-7x-4=0$

$6x^2-8x+x-4=0$

$6x(x-2)+1(x-2)=0$

$(6x+1)(x-2)=0$

Either $6x+1=0$ or $x-2=0$

Either $x=-\frac{1}{6}$ or $x=2$

Hence, $-\frac{1}{6}$, 2 and 3 are the roots of the given equation.

5. Solve by using synthetic division, if

(i) 3 and 2 the roots of the equation $x^3-10x^2+9=0$.

Solution:

Since 3 and 2 are the root of the equation

$x^3-10x^2+9=0$

Then by synthetic division, we get

$$\begin{array}{r|rrrr} 3 & 1 & -10 & 0 & 9 \\ & & 3 & -21 & -63 \\ \hline 2 & 1 & -7 & -21 & -54 \\ & & 2 & -10 & -62 \\ \hline & 1 & -5 & -31 & \boxed{-116} \end{array}$$

The depressed equation is

$x^2-5x-31=0$

$x^2-5x+(-31)=0$

$x(x+3)-31(x+3)=0$

$(x-31)(x+3)=0$

Either $x+3=0$ or $x-31=0$

Either $x=-3$ or $x=31$

Thus, -3, 2 and 3 are the roots of the given equation.

(ii) 3 and -4 are the roots of the equation $x^4+2x^3-13x^2-14x+24=0$.

Solution:

Since 3 and -4 are the root of the equation

$x^4+2x^3-13x^2-14x+24=0$

Then by synthetic division, we get

$$\begin{array}{r|rrrrr} 3 & 1 & 2 & -13 & -14 & 24 \\ & & 3 & 15 & 6 & -24 \\ \hline -4 & 1 & 5 & 2 & -8 & \boxed{0} \\ & & -4 & -4 & 8 & \\ \hline & 1 & 1 & -2 & \boxed{0} & \end{array}$$

The depressed equation is

$x^3+x-2=0$

$x^3+2x-x-2=0$

$x(x+1)-2(x+1)=0$

$(x-2)(x+1)=0$

Either $x+1=0$ or $x-2=0$

$x=-1$ $x=2$

Thus, -4, -1, 2 and 3 are the roots of the given equation.

Simultaneous equations:

A system of equations having a common solution is called a system of simultaneous equations.

The set of all the ordered pairs (x,y), which satisfies the system of equations is called the solution set of the system.

Exercise 2.7

Solve the following simultaneous equations.

1. $x + y = 5$; $x^2 - 2y - 14 = 0$

Solution:

$x+y=5$ ....(i)
$x^2-2y-14=0$ ....(ii)

From eq. (i), we have
$y=5-x$ ....(iii)

Put value of y in eq.(ii), we get
$x^2-2(5-x)-14=0$
$x^2-10+2x-14=0$
$x^2+2x-24=0$
$x^2+6x-4x-24=0$
$x(x+6)-4(x+6)=0$
$(x-4)(x+6)=0$

Either $x-4=0$ or $x+6=0$
$x=4$ $x=-6$

Put $x=4$ in eq.(iii), we get Put $x=-6$ in eq.(iii), we get

$y=5-4$ $y=5-(-6)$
$=1$ $=5+6$
$=11$

The ordered pairs are $(4,1)$ and $(-6,11)$.

Thus, solution set $=\{(4,1),(-6,11)\}$

2. $3x - 2y = 1$; $x^2 + xy - y^2 = 1$

Solution:

$3x-2y=1$ ....(i)
$x^2+xy-y^2=1$ ....(ii)

From eq. (i), we have
$2y=3x-1$
$y=\frac{1}{2}(3x-1)$
$y=\frac{3}{2}x-\frac{1}{2}$ ....(iii)

Put value of y in eq.(ii), we get
$x^2+x\left(\frac{3}{2}x-\frac{1}{2}\right)-\left(\frac{3}{2}x-\frac{1}{2}\right)^2=1$
$x^2+\frac{3}{2}x^2-\frac{1}{2}x-\frac{9}{4}x^2+\frac{3}{2}x-\frac{1}{4}=1$

Multiplying both sides by '4' we get
$4x^2+6x^2-2x-9x^2+6x-1=4$
$4x^2+6x^2-9x^2-2x+6x-1-4=0$
$x^2+4x-5=0$
$x^2+5x-x-5=0$
$x(x+5)-1(x+5)=0$
$(x-1)(x+5)=0$

Either $x-1=0$ or $x+5=0$
$x=1$ $x=-5$

Put $x=1$ in eq.(iii), we get Put $x=-5$ in eq.(iii), we get
$y=\frac{3}{2}(1)-\frac{1}{2}$ $y=\frac{3}{2}(-5)-\frac{1}{2}$
$=\frac{3}{2}-\frac{1}{2}$ $=-\frac{15}{2}-\frac{1}{2}$
$=\frac{2}{2}$ $=-\frac{16}{2}$
$=1$ $=-8$

The ordered pairs are $(1,1)$ and $(-5,-8)$

Thus, solution set $=\{(1,1),(-5,-8)\}$

3. $x-y=7$; $\frac{2}{x}-\frac{5}{y}=2$

Solution:

$x-y=7$ ....(i)
$\frac{2}{x}-\frac{5}{y}=2$
$\frac{2y-5x}{xy}=2$
$2y-5x=2xy$ ....(ii)

From eq. (i), we have
$y=x-7$ ....(iii)

Put value of y in eq.(ii), we get
$2(x-7)-5x=2x(x-7)$
$2x-14-5x=2x^2-14x$
$-3x-14=2x^2-14x$
$2x^2-14x+3x+14=0$

$2x^2-11x+14=0$
$2x^2-7x-4x+14=0$
$x(2x-7)-2(2x-7)=0$
$(x-2)(2x-7)=0$

Either $x-2=0$ or $2x-7=0$
$x=2$ $2x=7$
$x=\frac{7}{2}$

Put $x=2$ in eq.(iii), we get Put $x=\frac{7}{2}$ in eq.(iii), we get

$y=2-7$ $y=\frac{7}{2}-7$
$=-5$ $=-\frac{7}{2}$

The ordered pairs are $(2,-5)$ and $\left(\frac{7}{2},-\frac{7}{2}\right)$

Thus, solution set $=\left\{(2,-5),\left(\frac{7}{2},-\frac{7}{2}\right)\right\}$

4. $x+y=a-b$; $\frac{a}{x}-\frac{b}{y}=2$

Solution:

$x+y=a-b$ ....(i)
$\frac{a}{x}-\frac{b}{y}=2$
$\frac{ay-bx}{xy}=2$
$ay-bx=2xy$........(ii)

From eq.(i), we have
$y=a-b-x$ ....(iii)

Put value of y in eq.(ii), we get
$a(a-b-x)-bx=2x(a-b-x)$
$a(a-b-x)-bx-2x(a-b-x)=0$
$a^2-ab-ax-bx-2ax+2bx+2x^2=0$
$2x^2-2ax-ax+2bx-bx+a^2-ab=0$
$2x^2-3ax+bx+a^2-ab=0$
$2x^2-(3a-b)x+(a^2-ab)=0$

Using quadratic formula
$x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}=\frac{-[-(3a-b)]\pm\sqrt{[-(3a-b)]^2-4(2)(a^2-ab)}}{2(2)}$
$=\frac{(3a-b)\pm\sqrt{9a^2-6ab+b^2-8a^2+8ab}}{4}$
$=\frac{(3a-b)\pm\sqrt{a^2+2ab+b^2}}{4}$
$=\frac{(3a-b)\pm\sqrt{(a+b)^2}}{4}=\frac{(3a-b)\pm(a+b)}{4}$

Either $x=\frac{(3a-b)+(a+b)}{4}$ or $\frac{(3a-b)-(a+b)}{4}$
$x=\frac{3a-b+a+b}{4}$ or $\frac{3a-b-a-b}{4}$
$x=\frac{4a}{4}$ or $\frac{2a-2b}{4}$
$x=a$ or $\frac{2(a-b)}{4}=\frac{a-b}{2}$

Put $x=a$ in eq.(iii), we get Put $x=\frac{a-b}{2}$ in eq.(iii), we get
$y=a-b-a$ $y=(a-b)-\left(\frac{a-b}{2}\right)$
$=-b$ $=\frac{2a-2b-a+b}{2}$
$=\frac{a-b}{2}$

$\therefore$ The ordered pairs are $(a,-b)$ and $\left(\frac{a-b}{2},\frac{a-b}{2}\right)$.

Thus, solution set $=\left\{(a,-b),\left(\frac{a-b}{2},\frac{a-b}{2}\right)\right\}$

5. $x^2+(y-1)^2=10$; $x^2+y^2+4x=1$

Solution:

$x^2+(y-1)^2=10$
$x^2+y^2-2y+1=10$
$x^2+y^2-2y=9$ ....(i)
$x^2+y^2+4x=1$ ....(ii)

Subtract eq.(ii) from eq.(i), we have
$x^2+y^2-2y=9$
$\pm x^2\pm y^2\pm 4x=\pm 1$
$-4x-2y=8$
$-2(2x+y)=8$
$2x+y=-4$
$y=-2x-4$ ......(iii)

Put value of y in eq.(ii), we have
$x^2+(-2x-4)^2+4x=1$
$x^2+\{-(2x+4)\}^2+4x=1$
$x^2+4x^2+16x+16+4x-1=0$
$5x^2+20x+15=0$
$5(x^2+4x+3)=0$
$\Rightarrow x^2+4x+3=0$
$x^2+3x+x+3=0$
$x(x+3)+1(x+3)=0$
$(x+1)(x+3)=0$

Either $x+3=0$ or $x+1=0$
$x=-3$ or $x=-1$

Put $x=-3$ in eq.(iii), we get Put $x=-1$ in eq.(iii), we get
$y=-2(-3)-4$ $y=-2(-1)-4$
$=6-4$ $=2-4$
$=2$ $=-2$

$\therefore$ The ordered pairs are $(-3,2),(-1,-2)$.

Thus, solution set $=\{(-3,2),(-1,-2)\}$

6. $(x+1)^2+(y+1)^2=5$; $(x+2)^2+y^2=5$

Solution:

$(x+1)^2+(y+1)^2=5$
$x^2+2x+1+y^2+2y+1=5$
$x^2+y^2+2x+2y=5-1-1$
$x^2+y^2+2x+2y=3$ ....(i)
$(x+2)^2+y^2=5$
$x^2+4x+4+y^2=5$
$x^2+y^2+4x=5-4$
$x^2+y^2+4x=1$ ....(ii)

Subtract eq.(ii) from eq.(i), we have
$x^2+y^2+2x+2y=3$
$\pm x^2\pm y^2\pm 4x=\pm 1$
$-2x+2y=2$
$2(-x+y)=2$
$-x+y=1$
$y=x+1$ ......(iii)

Put value of y in eq.(ii), we have
$x^2+(x+1)^2+4x=1$
$x^2+x^2+2x+1+4x=1$
$2x^2+6x+1-1=0$
$2x^2+6x=0$

$2x(x+3)=0$

Either $2x=0$ or $x+3=0$
$x=0$ or $x=-3$

Put $x=0$ in eq.(iii), we get Put $x=-3$ in eq.(iii), we get
$y=0+1$ $y=-3+1$
$=1$ $=-2$

$\therefore$ The ordered pairs are $(0,1),(-3,-2)$.

Thus, solution set $=\{(0,1),(-3,-2)\}$

7. $x^2+2y^2=22$; $5x^2+y^2=29$

Solution:

$x^2+2y^2=22$ ....(i)
$5x^2+y^2=29$ ....(ii)

Multiply eq.(ii) by '2' and then Subtract eq.(ii) from eq.(i), we get
$x^2+2y^2=22$
$\pm 10x^2\pm 2y^2=\pm 58$
$-9x^2=-36$
$\Rightarrow x^2=4$
$x=\pm 2$

Put value of $x^2=4$ in eq.(i), we get
$4+2y^2=22$
$2y^2=22-4$
$2y^2=18$
$y^2=9$
$y=\pm 3$

Thus, solution set $=\{(\pm 2,\pm 3)\}$

8. $4x^2-5y^2=6$; $3x^2+y^2=14$

Solution:

$4x^2-5y^2=6$ ....(i)
$3x^2+y^2=14$ ....(ii)

Multiply eq.(ii) by 5 and then add eq.(i) in eq.(ii), we get
$4x^2-5y^2=6$
$15x^2+5y^2=70$
$19x^2=76$
$\Rightarrow x^2=4$
$x=\pm 2$

Put value of $x^2=4$ in eq.(ii), we get
$3(4)+y^2=14$
$12+y^2=14$
$y^2=14-12$
$y^2=2$
$y=\pm\sqrt{2}$

Thus, solution set $=\{(\pm 2,\pm\sqrt{2})\}$

9. $7x^2-3y^2=4$; $2x^2+5y^2=7$

Solution:

$7x^2-3y^2=4$ ....(i)
$2x^2+5y^2=7$ ....(ii)

Multiply eq.(i) by '5' and eq.(ii) by '3' then add eq.(i) and eq.(ii), we get
$35x^2-15y^2=20$
$6x^2+15y^2=21$
$41x^2=41$
$\Rightarrow x^2=1$
$x=\pm 1$

Put value of $x^2=1$ in eq.(ii), we get

$2(1)+5y^2=7$
$2+5y^2=7$
$5y^2=7-2$
$5y^2=5$
$y^2=1$
$y=\pm\sqrt{1}$

Thus, solution set $=\{(\pm 1,\pm 1)\}$

10. $x^2+2y^2=3$; $x^2+4xy-5y^2=0$

Solution:

$x^2+2y^2=3$ ....(i)
$x^2+4xy-5y^2=0$
$x^2+5xy-xy-5y^2=0$
$x(x+5y)-y(x+5y)=0$
$(x-y)(x+5y)=0$
$x-y=0$ or $x+5y=0$
or $y=x$ $x=-5y$

Put y = x in eq.(i), we get Put $x=-5y$ in eq.(i), we get
$x^2+2x^2=3$ $(-5y)^2+2y^2=3$
$3x^2=3$ $25y^2+2y^2=3$
$\Rightarrow x^2=1$ $27y^2=3$
$x=\pm 1$ $y^2=\frac{1}{9}$
$y=\pm\frac{1}{3}$

Put $x=1$ in $y=x$, we get Put $y=\frac{1}{3}$ in $x=-5y$, we get
y = 1 $x=-5\left(\frac{1}{3}\right)$
Put $x=-1$ in $y=x$, we get $=-\frac{5}{3}$

$y=-1$ Put $y=-\frac{1}{3}$ in $x=-5y$, we get
$x=-5\left(-\frac{1}{3}\right)$
$=\frac{5}{3}$

$\therefore$ The ordered pairs are $(1,1),(-1,-1)$. $\therefore$ The ordered pairs are $\left(\frac{5}{3},-\frac{1}{3}\right),\left(-\frac{5}{3},\frac{1}{3}\right)$

Thus, solution set $=\left\{(1,1),(-1,-1),\left(\frac{5}{3},-\frac{1}{3}\right),\left(-\frac{5}{3},\frac{1}{3}\right)\right\}$

11. $3x^2-y^2=26$; $3x^2-5xy-12y^2=0$

Solution:

$3x^2-y^2=26$ ....(i)
$3x^2-5xy-12y^2=0$
$3x^2-9xy+4xy-12y^2=0$
$3x(x-3y)+4y(x-3y)=0$
$(3x+4y)(x-3y)=0$

Either $3x+4y=0$ or $x-3y=0$
$3x=-4y$ $x=3y$
$x=-\frac{4}{3}y$

Put $x=3y$ in eq.(i), we get
$3(3y)^2-y^2=26$
$3(9y^2)-y^2=26$
$27y^2-y^2=26$
$26y^2=26$
$y^2=1$
$y=\pm 1$

Put $y=1$ in $x=2y$, we get
$x=3(1)$

$x=3$
Put $y=-1$ in $x=2y$, we get
$x=3(-1)$
$x=-3$

$\therefore$ The ordered pairs are $(3,1),(-3,-1)$

Put $x=-\frac{4}{3}y$ in eq.(i), we get
$3\left(-\frac{4}{3}y\right)^2-y^2=26$
$3\left(\frac{16}{9}y^2\right)-y^2=26$
$\frac{16}{3}y^2-y^2=26$
$\frac{16y^2-3y^2}{3}=26$
$13y^2=26\times 3$
$y^2=\frac{78}{13}$
$y^2=6$
$y=\pm\sqrt{6}$

Put $y=\sqrt{6}$ in $x=-\frac{4}{3}y$, we get
$x=-\frac{4}{3}(\sqrt{6})$
$x=\frac{-4\sqrt{6}}{3}$
Put $y=-\sqrt{6}$ in $x=-\frac{4}{3}y$, we get
$x=-\frac{4}{3}(-\sqrt{6})$
$x=\frac{4\sqrt{6}}{3}$

$\therefore$ The ordered pairs are $\left(\frac{-4\sqrt{6}}{3},\sqrt{6}\right),\left(\frac{4\sqrt{6}}{3},-\sqrt{6}\right)$

Thus, solution set $=\left\{(3,1),(-3,-1),\left(\frac{-4\sqrt{6}}{3},\sqrt{6}\right),\left(\frac{4\sqrt{6}}{3},-\sqrt{6}\right)\right\}$

12. $x^2+xy=5$; $y^2+xy=3$

Solution:

$x^2+xy=5$ ....(i)
$y^2+xy=3$ ....(ii)

Multiply eq.(i) by '3' and eq.(ii) by '5' then subtract eq.(ii) from eq.(i), we get
$3x^2+3xy=15$
$\pm 5y^2\pm 5xy=\pm 15$
$3x^2-2xy-5y^2=0$
$3x^2-5xy+3xy-5y^2=0$
$x(3x-5y)+y(3x-5y)=0$
$(x+y)(3x-5y)=0$

Either $x+y=0$ or $3x-5y=0$
$y=-x$ $3x=5y$
$x=\frac{5}{3}y$

Put y = −x in eq.(i), we get Put x $=\frac{5}{3}y$ in eq.(i), we get
$x^2+x(-x)=5$ $\left(\frac{5}{3}y\right)^2+\left(\frac{5}{3}y\right)y=5$
$x^2-x^2=5$ $\frac{25}{9}y^2+\frac{5}{3}y^2=5$
$0=5$ $\frac{25y^2+15y^2}{9}=5$
Which is not possible $\frac{40y^2}{9}=5$
$40y^2=45$
$y^2=\frac{45}{40}$
$y^2=\frac{9}{8}$

$y=\pm\frac{3}{2\sqrt{2}}$

Put y $=\frac{3}{2\sqrt{2}}$ in $x=\frac{5}{3}y$, we get
$x=\frac{5}{3}\left(\frac{3}{2\sqrt{2}}\right)$
$x=\frac{5}{2\sqrt{2}}$
Put y $=-\frac{3}{2\sqrt{2}}$ in $x=\frac{5}{3}y$, we get
$x=\frac{5}{3}\left(-\frac{3}{2\sqrt{2}}\right)$
$x=-\frac{5}{2\sqrt{2}}$

$\therefore$ The ordered pairs are $\left(\frac{5}{2\sqrt{2}},\frac{3}{2\sqrt{2}}\right),\left(-\frac{5}{2\sqrt{2}},-\frac{3}{2\sqrt{2}}\right)$

Thus, solution set $=\left\{\left(\frac{5}{2\sqrt{2}},\frac{3}{2\sqrt{2}}\right),\left(-\frac{5}{2\sqrt{2}},-\frac{3}{2\sqrt{2}}\right)\right\}$

13. $x^2-2xy=7$; $xy+3y^2=2$

Solution:

$x^2-2xy=7$ ....(i)
$xy+3y^2=2$ ....(ii)

Multiply eq.(i) by '2' and eq.(ii) by '7' then subtract eq.(ii) from eq.(i), we get
$2x^2-4xy=14$
$\pm 21y^2\pm 7xy=\pm 14$
$2x^2-11xy-21y^2=0$
$2x^2-14xy+3xy-21y^2=0$
$2x(x-7y)+3y(x-7y)=0$
$(2x+3y)(x-7y)=0$

Either $2x+3y=0$ or $x-7y=0$
$2x=-3y$ $x=7y$
$x=-\frac{3}{2}y$

Put $x=-\frac{3}{2}y$ in eq.(ii), we get Put $x=7y$ in eq.(ii), we get
$\left(-\frac{3}{2}y\right)y+3y^2=2$ $(7y)y+3y^2=2$
$-\frac{3}{2}y^2+3y^2=2$ $7y^2+3y^2=2$
$\frac{-3y^2+6y^2}{2}=2$ $10y^2=2$
$3y^2=4$ $y^2=\frac{2}{10}$
$y^2=\frac{4}{3}$ $y=\pm\frac{\sqrt{2}}{\sqrt{10}}$
$y=\pm\frac{2}{\sqrt{3}}$ $y=\pm\frac{\sqrt{2}}{\sqrt{2}\times\sqrt{5}}$
$y=\pm\frac{1}{\sqrt{5}}$

Put y $=\frac{2}{\sqrt{3}}$ in $x=-\frac{3}{2}y$, we get
$x=-\frac{3}{2}\left(\frac{2}{\sqrt{3}}\right)$
$x=-\frac{3}{\sqrt{3}}\times\frac{\sqrt{3}}{\sqrt{3}}$
$x=-\frac{3\sqrt{3}}{(\sqrt{3})^2}=-\frac{3\sqrt{3}}{3}$
$x=-\sqrt{3}$

Put y $=-\frac{2}{\sqrt{3}}$ in $x=-\frac{3}{2}y$, we get
$x=-\frac{3}{2}\left(-\frac{2}{\sqrt{3}}\right)$
$x=\frac{3}{\sqrt{3}}\times\frac{\sqrt{3}}{\sqrt{3}}$
$x=\frac{3\sqrt{3}}{(\sqrt{3})^2}=\frac{3\sqrt{3}}{3}$

$x=\sqrt{3}$
$\therefore$ The ordered pairs are $\left(-\sqrt{3},\frac{2}{\sqrt{3}}\right),\left(\sqrt{3},-\frac{2}{\sqrt{3}}\right)$

Put y $=\frac{1}{\sqrt{5}}$ in $x=7y$, we get
$x=7\left(\frac{1}{\sqrt{5}}\right)$
$x=\frac{7}{\sqrt{5}}$
Put y $=-\frac{1}{\sqrt{5}}$ in $x=7y$, we get
$x=7\left(-\frac{1}{\sqrt{5}}\right)$
$x=-\frac{7}{\sqrt{5}}$

$\therefore$ The ordered pairs are $\left(\frac{7}{\sqrt{5}},\frac{1}{\sqrt{5}}\right),\left(-\frac{7}{\sqrt{5}},-\frac{1}{\sqrt{5}}\right)$

Thus, solution set $=\left\{\left(\frac{7}{\sqrt{5}},\frac{1}{\sqrt{5}}\right),\left(-\frac{7}{\sqrt{5}},-\frac{1}{\sqrt{5}}\right),\left(-\sqrt{3},\frac{2}{\sqrt{3}}\right),\left(\sqrt{3},-\frac{2}{\sqrt{3}}\right)\right\}$

Exercise 2.8

1. The product of two positive consecutive numbers is 182. Find the numbers.

Solution:

Let the numbers be $x, x+1$

Then, $x(x+1) = 182$

$x^2 + x - 182 = 0$

$x^2 + 14x - 13x - 182 = 0$

$x(x+14) - 13(x+14) = 0$

$(x-13)(x+14) = 0$

$x - 13 = 0 \quad \text{gives} \quad x = 13$

Numbers are: $x = 13$

$x + 1 = 13 + 1 = 14$

13,14

Now, $x + 14 = 0$

gives $x = -14$

we ignore this value because it is negative

2. The sum of squares of three positive consecutive numbers is 77. Find the numbers.

Solution:

Let the numbers be $x, x+1, x+2$

Applying the given conditions

$x^2 + (x+1)^2 + (x+2)^2 = 77$

$x^2 + x^2 + 2x + 1 + x^2 + 4x + 4 = 77$

$3x^2 + 6x + 5 = 77$

$3x^2 + 6x - 5 - 77 = 0$

$3x^2 + 6x - 72 = 0$

$x^2 + 2x - 24 = 0 \quad \text{(Dividing by 3)}$

$x^2 + 6x - 4x - 24 = 0$

$x(x+6) - 4(x+6) = 0$

$(x-4)(x+6) = 0$

$x - 4 = 0 \quad \text{gives} \quad x = 4$

Numbers are: $x = 4$

$x + 1 = 4 + 1 = 5$

$x + 2 = 4 + 2 = 6$

4,5,6

$x + 6 = 0$

$x = -6$, we ignore the negative number.

3. The sum of five times a number and the square of the number is 204. Find the number.

Solution:

Let the numbers be

According to the given conditions,

$x^2 + 5x = 204$

$x^2 + 5x - 204 = 0$

$x^2 + 17x - 12x - 204 = 0$

$x(x+17) - 12(x+17) = 0$

$(x+17)(x-12) = 0$

$x - 12 = 0 \quad \text{gives} \quad x = 12$

or $x + 17 \quad \text{gives} \quad x = -17$

Number is 12 or -17.

4. The product of five less than three times a certain number and one less than four times the number is 7. Find the number.

Solution:

Let the numbers be x

According to the given conditions,

$(3x-5)(4x-1) = 7$

$12x^2 - 3x - 20x + 5 = 7$

$12x^2 - 23x + 5 = 7$

$12x^2 - 23x + 5 - 7 = 0$

$12x^2 - 23x - 2 = 0$

$12x^2 - 24x + x - 2 = 0$

$12x(x-2) + 1(x-2) = 0$

$(12x+1)(x-2) = 0$

$x - 2 = 0 \quad \text{gives} \quad x = 2$

and $12x + 1 = 0 \quad \text{gives} \quad x = -\frac{1}{12}$

Number is 2 or $-\frac{1}{12}$

5. The difference of a number and its reciprocal is $\frac{15}{4}$. Find the number.

Solution:

Let the numbers be x

Then, According to the given conditions,

$x - \frac{1}{x} = \frac{15}{4}$

$4x^2 - 4 = 15x \quad \text{(Multiplying by 4x)}$

$4x^2 - 15x - 4 = 0$

$4x^2 - 16x + x - 4 = 0$

$4x(x-4) + 1(x-4) = 0$

$(x-4)(4x+1) = 0$

$x - 4 = 0 \quad \text{gives} \quad x = 4$

and $4x + 1 = 0 \quad \text{gives} \quad x = -\frac{1}{4}$

Number is 4 and $-\frac{1}{4}$

6. The sum of squares of two digits of a positive integral number is 65 and the number is 9 times the sum of its digits. Find the number.

Solution:

Let xy be the number, where unit digit is y and tens digit is x.

According the given condition,

$x^2 + y^2 = 65 \quad \ldots\ldots(A)$

Number $= y + 10x$

$y + 10x = 9(x + y)$

$y + 10x = 9x + 9y$

$10x - 9x = 9y - y$

$x = 8y \quad \ldots\ldots(B)$

Putting $x = 8y$ *in* (A)

$(8y)^2 + y^2 = 65$

$64y^2 + y^2 = 65$

$65y^2 = 65$

$y^2 = 1$

$y = 1$

$y = 1 \quad \text{(taking +ve value)}$

Putting $y = 1$ *in* (B)

$x = 8(1) = 8$

Now,

Number $= y + 10x$

$= 8 + 10(8)$

$= 81$

Therefore, number is 81.

7. The sum of the co-ordinates of a point is 7 and sum of their squares is 49. Find the co-ordinates of the point.

Solution:

Let $P(x, y)$ be the point.

According the given condition,

$x + y = 9 \quad \ldots\ldots(A)$

and $x^2 + y^2 = 49 \quad \ldots\ldots(B)$

From A $\quad x = 9 - y \quad \ldots\ldots(C)$

Putting $x = 9 - y$ in (B)

$(9-y)^2 + y^2 = 45$

$81 - 18y + y^2 + y^2 = 45$

$2y^2 - 18y + 81 - 45 = 0$

$2y^2 - 18y + 36 = 0$

$y^2 - 9y + 18 = 0 \quad \text{(Dividing by 2)}$

$y^2 - 6y - 3y + 18 = 0$

$y(y-6) - 3(y-6) = 0$

$(y-6)(y-3) = 0$

$y - 6 = 0 \quad \text{gives} \quad y = 6$

Then from (C)

$x = 9 - 6 = 3$

Point is $P(3,6)$

When $y = 3$ then $y = 3$

From (C)

$x = 9 - 3 = 6$

Point is $P(6,3)$

8. Find two integers whose sum is 9 and the difference of their squares is also 9.

Solution:

Let the integers be x, y.

Then,

$x + y = 9 \quad \ldots\ldots(A)$

and $x^2 - y^2 = 9 \quad \ldots\ldots(B)$

From A $\quad x = 9 - y$

Putting $x = 9 - y$ in (B)

$(9-y)^2 - y^2 = 9$

$81 - 18y + y^2 - y^2 = 9$

$-18y = 9 - 81$

$-18y = -72$

$y = \frac{-72}{-18}$

$y = 4$

Putting $y = 4$ in (A)

$x + 4 = 9$

$x = 9 - 4 = 5$

Integers are 5,4

9. Find two integers whose difference is 4 and whose squares differ by 72.

Solution:

Let the integers be x, y.

Then,

$x - y = 4 \quad \ldots\ldots(A)$

and $x^2 - y^2 = 72 \quad \ldots\ldots(B)$

From A $\quad x - y = 4$

$x = 4 + y$

Putting $x = 4 + y$ in (B), we get

$(4+y)^2 - y^2 = 72$

$16 + y^2 + 8y - y^2 = 72$

$8y = 72 - 16$

$8y = 56$

$y = \frac{56}{8}$

$y = 7$

Putting $y = 7$ in (A)

$x - 7 = 4$

$x = 4 + 7 = 11$

Integers are 11 and 7.

10. Find the dimensions of a rectangle, whose perimeter is 80cm and its area is 375cm².



Solution:

Let x and y be the length and width respectively of the rectangle.

According to the given conditions,

Perimeter $\quad 2x + 2y = 80$

$or \quad x + y = 40 \quad \ldots\ldots(A)$

Area $\quad xy = 375 \quad \ldots\ldots(B)$

From A $\quad x = 40 - y$

Putting $x = 40 - y$ in (B), we get

$(40 - y)y = 375$

$40y - y^2 = 375$

$\Rightarrow y^2 - 40y + 375 = 0$

$y^2 - 25y - 15y + 375 = 0$

$y(y - 25) - 15(y - 25) = 0$

$(y - 15)(y - 25) = 0$

$y - 15 = 0 \quad \text{gives} \quad y = 15$

Putting $y = 15$ in (A)

$x + 15 = 40$

$x = 40 - 15 = 25$

Length = 25cm Breadth = 15cm

Miscellaneous Exercise 2

1. Multiple Choice Questions

Four possible answers are given for the following questions. Tick (✓) the correct answer.

(i) If x_1 is the root of $x^2 + x + 1 = 0$ then x is

(a) $\frac{1}{x_1}$ (b) x_1^2 (c) $\frac{1}{x_1^2}$ (d) $\frac{x_1}{2}$

(ii) If x_1, x_2 are the roots of $x^2 + x + 1 = 0$, then $x_1 \cdot x_2$ is

(a) $\frac{1}{2}$ (b) $-\frac{1}{2}$ (c) 1 (d) -1

(iii) Nature of equation $4x^2 - 3x + 2 = 0$ is

(a) Irrational (b) Imaginary (c) Rational (d) None

(iv) Sum of roots of $-x^2 + 3x + 5 = 0$ is

(a) 3 (b) -3 (c) 5 (d) -5

(v) Sum of the cube roots of unity is

(a) 1 (b) 0 (c) -1 (d) 2

(vi) Product of cube roots of unity is

(a) 1 (b) 0 (c) -1 (d) 2

(vii) If $3x^2 - 4x + 12 = 0$ then the roots of $x^2 + 4x + 12 = 0$ are

(a) Irrational (b) Imaginary (c) Rational (d) None

(viii) If $3x^2 - 4x + 12 = 0$ and not a perfect square then roots of $x^2 + 4x + 12 = 0$ are

(a) Imaginary (b) Rational (c) Irrational (d) None

(ix) $\begin{vmatrix} \frac{1}{x} & 1 \\ 1 & x \end{vmatrix}$ **is equal to**

(a) $\begin{vmatrix} x & 1 \\ 1 & \frac{1}{x} \end{vmatrix}$ (b) $\begin{vmatrix} \frac{1}{x} & 1 \\ 1 & x \end{vmatrix}$ (c) $\begin{vmatrix} \frac{1}{x} & 1 \\ 1 & x \end{vmatrix}$ (d) $\begin{vmatrix} x & 1 \\ 1 & \frac{1}{x} \end{vmatrix}$

(x) $\begin{vmatrix} a & b \\ c & d \end{vmatrix}$ **is equal to**

(a) $\begin{vmatrix} a & c \\ b & d \end{vmatrix}$ (b) $\begin{vmatrix} -a & -b \\ -c & -d \end{vmatrix}$

(c) $-\begin{vmatrix} a & b \\ c & d \end{vmatrix}$ (d) $\begin{vmatrix} -a & b \\ c & -d \end{vmatrix}$

(xi) If the cube roots of unity are

(a) $1, -1$ (b) $1, 0$ (c) $1, \omega, \omega^2$ (d) $0, \omega, \omega^2$

(xii) Roots of equation $3x^2 - 5x + 7 = 0$ are

(a) Rational (b) Irrational (c) Imaginary (d) Irrational

(xiii) If ω_1, ω_2 are roots of $x^2 + x + 1 = 0$ then sum of roots ω_1 and ω_2 is

(a) $\frac{1}{2}$ (b) $-\frac{1}{2}$ (c) $\frac{1}{3}$ (d) $-\frac{1}{3}$

(xiv) If x_1, x_2 are the roots of $x^2 - x + 1 = 0$ then product of roots x_1 and x_2 is

(a) 1 (b) 0 (c) -1 (d) 2

(xv) The nature of roots of equation $ax^2 + bx + c = 0$ is determined by

(a) Sum of roots (b) Product of roots

(c) Discriminant (d) Environment

Just Plus and Minimal (Level 1 to 3 Ex.)

Solutions												
(i)	(a)	(ii)	(b)	(iii)	(b)	(iv)	(b)	(v)	(b)	(vi)	(a)	
(vii)	(b)	(viii)	(c)	(ix)	(c)	(x)	(c)	(xi)	(c)	(xii)	(b)	
(xiii)	(b)	(xiv)	(a)	(xv)	(c)	(xvi)	(d)	(xvii)	(c)	(xviii)	(c)	
(xix)	(c)	(xx)	(c)									

Write brief answers of the following questions.

(i) Discuss the nature of the roots of the following equations.

(a) $x^2 - 7x + 5 = 0$ (b) $3x^2 - 7x + 5 = 0$

(c) $2x^2 - 7x + 5 = 0$ (d) $25x^2 - 70x + 49 = 0$

(e) $x^2 + 3x + 5 = 0$

Here $a = 1, b = -7, c = 5$

Disc. $= b^2 - 4ac$

$= (-7)^2 - 4(1)(5)$

$= 49 - 20$

$= 29$

The, it roots is Rationa, real an integers.

(b) $3x^2 - 7x + 5 = 0$

Here $a = 3, b = -7, c = 5$

Disc. $= b^2 - 4ac$

$= (-7)^2 - 4(3)(5)$

$= 49 - 60$

$= -11$

The, it roots is rationa, Irrationa, real an, not, rational and equal.

(c) $2x^2 - 7x + 5 = 0$

Here $a = 2, b = -7, c = 5$

Disc. $= b^2 - 4ac$

$= (-7)^2 - 4(2)(5)$

$= 49 - 40$

$= 9$

The, it roots is rationa, Irrationa, real an, not, rational and equal.

(d) $25x^2 - 70x + 49 = 0$

Here $a = 25, b = -70, c = 49$

Disc. $= b^2 - 4ac$

$= (-70)^2 - 4(25)(49)$

$= 4900 - 4900$

$= 0$ (real)

Rone and, rationa and equal.

(e) $x^2 + 3x + 5 = 0$

Here $a = 1, b = 3, c = 5$

Disc. $= b^2 - 4ac$

$= (3)^2 - 4(1)(5)$

$= 9 - 20$

$= -11$

The, it roots, rationa, Irrationa, real an, not, rational and equal.

(ii) If $\alpha^2 - \beta^2 = \alpha + \frac{\beta}{\alpha}$ **, find α.**

Solution:

Given $\alpha^2 - \beta^2 = \alpha + \frac{\beta}{\alpha}$

$\alpha^2 - \beta^2 = \frac{\alpha^2 + \beta}{\alpha}$

$\alpha(\alpha^2 - \beta^2) = \alpha^2 + \beta$

$\alpha^3 - \alpha\beta^2 = \alpha^2 + \beta$

$\alpha^3 - \alpha^2 - \alpha\beta^2 - \beta = 0$

$\alpha^2(\alpha - 1) - \beta(\alpha^2 + 1) = 0$

$\alpha^2(\alpha - 1) = \beta(\alpha^2 + 1)$

$\frac{\alpha^2(\alpha - 1)}{\alpha^2 + 1} = \beta$

$\frac{\alpha^2(\alpha - 1)}{\alpha^2 + 1} = \beta$

$\alpha = \beta$

(iii) Find the product of complex cube roots of unity.

Solution:

Here we,

$\omega = \frac{-1 + \sqrt{3}i}{2}$, $\omega^2 = \frac{-1 - \sqrt{3}i}{2}$

Product $= \left(\frac{-1 + \sqrt{3}i}{2}\right)\left(\frac{-1 - \sqrt{3}i}{2}\right)$

$= \frac{(-1)^2 - (\sqrt{3}i)^2}{4}$

$= \frac{1 - 3i^2}{4}$

$= \frac{1 + 3}{4}$

$= \frac{4}{4}$

$= 1$

(iv) Prove that the sum of all cube roots of unity is zero.

Solution:

We have the cube roots of unity are

$1, \omega, \omega^2$

Now, $1 + \omega + \omega^2 = 1 + \frac{-1 + \sqrt{3}i}{2} + \frac{-1 - \sqrt{3}i}{2}$

$= \frac{2 - 1 + \sqrt{3}i - 1 - \sqrt{3}i}{2}$

$= \frac{0}{2}$

$= 0$

$1 + \omega + \omega^2 = 0$

(v) Find the value of $\left(\frac{-1 + \sqrt{3}i}{2}\right)^3 + \left(\frac{-1 - \sqrt{3}i}{2}\right)^3$

Solution:

Let $\omega = \frac{-1 + \sqrt{3}i}{2}$, $\omega^2 = \frac{-1 - \sqrt{3}i}{2}$

Now $\omega^3 + (\omega^2)^3 = \omega^3 + \omega^6 = \omega^3(1 + \omega^3)$

$= \omega^3(1 + 1)$

$= 2\omega^3$

$= 2(1)$

$= 2$

(vi) Prove that $x^2 + x^2(y+z) + y^2(z+x) + z^2(x+y) + 3xyz = (x+y+z)^3$

Solution:

Here L.S

$= x^3 + x^2y + x^2z + y^2z + y^2x + z^2x + z^2y + 3xyz$

$= x^3 + x^2y + x^2z + xyz + y^2z + y^2x + xyz + z^2x + z^2y + xyz$

$= x(x^2 + xy + xz + yz) + y(xy + y^2 + yz + xz) + z(xz + yz + z^2 + xy)$

$= x(x+y)(x+z) + y(y+x)(y+z) + z(z+x)(z+y)$

$= x^3 + x^2z + x^2y + xyz + y^3 + y^2z + y^2x + xyz + z^3 + z^2x + z^2y + xyz$

$= x^3 + y^3 + z^3 + 3xyz$

(vii) Evaluate $\begin{vmatrix} 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 \\ \omega^2 & 1 & \omega \end{vmatrix}$

Solution:

$= 1(\omega^2 - 1) - \omega(\omega - \omega^2) + \omega^2(1 - \omega^2)$

$= \omega^2 - 1 - \omega^2 + \omega^3 + \omega^2 - \omega^4$

$= \omega^2 - 1 - \omega^2 + 1 + \omega^2 - \omega$

$= \omega^2 - \omega$

(viii) Evaluate $\begin{vmatrix} 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 \\ \omega^2 & 1 & \omega \end{vmatrix}$

Solution:

$= 1(\omega^2 - 1) - \omega(\omega - \omega^2) + \omega^2(1 - \omega^2)$

$= \omega^2 - 1 - \omega^2 + \omega^3 + \omega^2 - \omega^4$

$= \omega^2 - 1 - \omega^2 + 1 + \omega^2 - \omega$

$= \omega^2 - \omega$

(ix) If x is one root of unity, form an equation whose roots are $2x$ and $3x^2$

Solution:

Since x is root

$\therefore$ Value of roots $= 1$

$\therefore$ Product of roots $= 1$

$2x \cdot 3x^2 = 1$

$6x^3 = 1$

$x^3 = \frac{1}{6}$

$x^3 - \frac{1}{6} = 0$

(x) If ω is root of unity, find the remainder and quotient when $\left(x^3 + 3x^2 + 3x + 1\right) \div \left(x + \frac{1}{\omega}\right)$

Solution:

Let $f(x) = x^3 + 3x^2 + 3x + 1$

$\therefore x + \frac{1}{\omega} = 0$

$x = -\frac{1}{\omega}$

Using synthetic division,

	1	3	3	1
$-\frac{1}{\omega}$		$-\frac{1}{\omega}$	$-\frac{2}{\omega}$	$-\frac{1}{\omega}$
	1	$3 - \frac{1}{\omega}$	$3 - \frac{2}{\omega}$	$1 - \frac{1}{\omega}$

Quotient $= x^2 + \left(3 - \frac{1}{\omega}\right)x + \left(3 - \frac{2}{\omega}\right)$

Remainder $= 1 - \frac{1}{\omega}$

(xi) Find the cube root of unity, hence show that $(1 - \omega + \omega^2)^3 = -8$

Solution:

Let $x^3 = 1$

$x^3 - 1 = 0$

$(x - 1)(x^2 + x + 1) = 0$

$x - 1 = 0$ or $x^2 + x + 1 = 0$

$x = 1$ or $x = \frac{-1 \pm \sqrt{1^2 - 4(1)(1)}}{2(1)}$

$x = \frac{-1 \pm \sqrt{-3}}{2}$

$x = \frac{-1 \pm \sqrt{3}i}{2}$

$x = 1, \frac{-1 + \sqrt{3}i}{2}, \frac{-1 - \sqrt{3}i}{2}$

$x = 1, \omega, \omega^2$

Now $(1 - \omega + \omega^2)^3 = (-\omega - \omega)^3$

$= (-2\omega)^3$

$= -8\omega^3$

$= -8(1)$

$= -8$

(xii) Find the remainder and quotient of the roots of the equation $\left(x^3 - 7x + 6\right) \div \left(x - \frac{1}{\omega}\right)$

Solution:

Let $f(x) = x^3 - 7x + 6$

$\therefore x - \frac{1}{\omega} = 0$

$x = \frac{1}{\omega}$

$x = \frac{\omega^3}{\omega}$

$x = \omega^2$

Using synthetic division,

	1	0	-7	6
ω^2		ω^2	$\omega^2 + \omega^4$	$-6 - 6\omega^2$
	1	ω^2	$-6 + \omega$	0

Quotient $= x^2 + \omega^2 x + (-6 + \omega)$

Remainder $= 0$

$\therefore x - \frac{1}{\omega}$ is factor of $f(x)$

(xiii) Find the sum and product of the roots of the equation $\left(ax^2 + bx + c\right) = 0$

Solution:

Let $ax^2 + bx + c = 0$

Here $a = a, b = b, c = c$

$\alpha + \beta = \frac{-b}{a}$

$\alpha\beta = \frac{c}{a}$

(xiv) Find $\frac{1}{\alpha} + \frac{1}{\beta}$ **of the roots of the equation** $x^2 - 4x + 12 = 0$.

Solution:

Here $a = 1, b = -4, c = 12$

$\frac{1}{\alpha} + \frac{1}{\beta} = \frac{\beta + \alpha}{\alpha\beta}$

$= \frac{-b}{a} \div \frac{c}{a}$

$= \frac{-b}{a} \times \frac{a}{c}$

$= \frac{-b}{c}$

Here $a = 1, b = -4, c = 12$

$\frac{1}{\alpha} + \frac{1}{\beta} = \frac{-b}{c}$

$= \frac{-(-4)}{12}$

$= \frac{4}{12}$

$= \frac{1}{3}$

(xv) If α, β are the roots of $x^2 - 3x + 7 = 0$, find an equation whose roots are

(i) $\alpha + 2$ (ii) $3\alpha, 3\beta$

(i) $\alpha + 2$

Solution:

Given $x^2 - 3x + 7 = 0$

Equation $= x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$

$\therefore$ Sum of roots $= \alpha + 2 + \beta + 2$

$= \alpha + \beta + 4$

$= 3 + 4$

$= 7$

Product of roots $= (\alpha + 2)(\beta + 2)$

$= \alpha\beta + 2\alpha + 2\beta + 4$

$= \alpha\beta + 2(\alpha + \beta) + 4$

$= 7 + 2(3) + 4$

$= 7 + 6 + 4$

$= 17$

Equation $= x^2 - 7x + 17 = 0$

(ii) $3\alpha, 3\beta$

Here $a = 1, b = -3, c = 7$

$\alpha + \beta = \frac{-b}{a}$

$= \frac{-(-3)}{1}$

$= 3$

$\alpha\beta = \frac{c}{a}$

$= \frac{7}{1}$

$= 7$

Now $3\alpha + 3\beta = 3(\alpha + \beta)$

$= 3(3)$

$= 9$

$3\alpha \cdot 3\beta = 9\alpha\beta$

$= 9(7)$

$= 63$

Equation $= x^2 - 9x + 63 = 0$

(xvi) If α, β are the roots of $x^2 - 3x + 7 = 0$, find an equation whose roots are

(i) $\alpha - \frac{2}{\beta}$ (ii) $\alpha, \frac{1}{\beta}$

(i) $\alpha - \frac{2}{\beta}$

Solution:

Given $x^2 - 3x + 7 = 0$

Equation $= x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$

$\therefore$ Sum of roots $= \alpha - \frac{2}{\beta} + \beta - \frac{2}{\alpha}$

$= \alpha + \beta - \frac{2}{\beta} - \frac{2}{\alpha}$

$= \alpha + \beta - 2\left(\frac{1}{\beta} + \frac{1}{\alpha}\right)$

$= \alpha + \beta - 2\left(\frac{\alpha + \beta}{\alpha\beta}\right)$

$= 3 - 2\left(\frac{3}{7}\right)$

$= 3 - \frac{6}{7}$

$= \frac{21 - 6}{7}$

$= \frac{15}{7}$

Product of roots $= \left(\alpha - \frac{2}{\beta}\right)\left(\beta - \frac{2}{\alpha}\right)$

$= \alpha\beta - 2 - 2 + \frac{4}{\alpha\beta}$

$= \alpha\beta - 4 + \frac{4}{\alpha\beta}$

$= 7 - 4 + \frac{4}{7}$

$= 3 + \frac{4}{7}$

$= \frac{21 + 4}{7}$

$= \frac{25}{7}$

Equation $= x^2 - \frac{15}{7}x + \frac{25}{7} = 0$

$7x^2 - 15x + 25 = 0$

(ii) $\alpha, \frac{1}{\beta}$

Here $a = 1, b = -3, c = 7$

$\alpha + \beta = \frac{-b}{a}$

$= \frac{-(-3)}{1}$

$= 3$

$\alpha\beta = \frac{c}{a}$

$= \frac{7}{1}$

$= 7$

Now $\alpha + \frac{1}{\beta} = \frac{\alpha\beta + 1}{\beta}$

$= \frac{7 + 1}{\beta}$

$= \frac{8}{\beta}$

$\alpha \cdot \frac{1}{\beta} = \frac{\alpha}{\beta}$

$= \frac{\alpha^2}{\alpha\beta}$

$= \frac{\alpha^2}{7}$

Equation $= x^2 - \left(\frac{8}{\beta}\right)x + \frac{\alpha^2}{7} = 0$

$7\beta x^2 - 8\alpha x + \alpha^2 = 0$

(xvii) If α, β are the roots of $x^2 - 3x + 7 = 0$, find an equation whose roots are α^2, β^2

Solution:

Given $x^2 - 3x + 7 = 0$

Equation $= x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$

$\therefore$ Sum of roots $= \alpha^2 + \beta^2$

$= (\alpha + \beta)^2 - 2\alpha\beta$

$= (3)^2 - 2(7)$

$= 9 - 14$

$= -5$

Product of roots $= \alpha^2\beta^2$

$= (\alpha\beta)^2$

$= (7)^2$

$= 49$

Equation $= x^2 - (-5)x + 49 = 0$

$x^2 + 5x + 49 = 0$

(xviii) If α, β are the roots of $x^2 - 3x + 7 = 0$, find an equation whose roots are $\frac{\alpha}{\beta}, \frac{\beta}{\alpha}$

Solution:

Given $x^2 - 3x + 7 = 0$

Equation $= x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$

$\therefore$ Sum of roots $= \frac{\alpha}{\beta} + \frac{\beta}{\alpha}$

$= \frac{\alpha^2 + \beta^2}{\alpha\beta}$

$= \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta}$

$= \frac{(3)^2 - 2(7)}{7}$

$= \frac{9 - 14}{7}$

$= -\frac{5}{7}$

Product of roots $= \frac{\alpha}{\beta} \cdot \frac{\beta}{\alpha}$

$= 1$

Equation $= x^2 - \left(-\frac{5}{7}\right)x + 1 = 0$

$7x^2 + 5x + 7 = 0$

(xix) If α, β are the roots of $x^2 - 3x + 7 = 0$, find an equation whose roots are

(i) α^3, β^3 (ii) $\alpha^3\beta, \alpha\beta^3$

(i) α^3, β^3

Solution:

Given $x^2 - 3x + 7 = 0$

Here $a = 1, b = -3, c = 7$

$\alpha + \beta = \frac{-b}{a}$

$= \frac{-(-3)}{1}$

$= 3$

$\alpha\beta = \frac{c}{a}$

$= \frac{7}{1}$

$= 7$

Equation $= x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$

$\therefore$ Sum of roots $= \alpha^3 + \beta^3$

$= (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$

$= (3)^3 - 3(7)(3)$

$= 27 - 63$

$= -36$

Product of roots $= \alpha^3\beta^3$

$= (\alpha\beta)^3$

$= (7)^3$

$= 343$

Equation $= x^2 - (-36)x + 343 = 0$

$x^2 + 36x + 343 = 0$

(ii) $\alpha^3\beta, \alpha\beta^3$

Here $a = 1, b = -3, c = 7$

$\alpha + \beta = \frac{-b}{a}$

$= \frac{-(-3)}{1}$

$= 3$

$\alpha\beta = \frac{c}{a}$

$= \frac{7}{1}$

$= 7$

Now $\alpha^3\beta + \alpha\beta^3 = \alpha\beta(\alpha^2 + \beta^2)$

$= \alpha\beta((\alpha + \beta)^2 - 2\alpha\beta)$

$= 7((3)^2 - 2(7))$

$= 7(9 - 14)$

$= 7(-5)$

$= -35$

$\alpha^3\beta \cdot \alpha\beta^3 = \alpha^4\beta^4$

$= (\alpha\beta)^4$

$= (7)^4$

$= 2401$

Equation $= x^2 - (-35)x + 2401 = 0$

$x^2 + 35x + 2401 = 0$

Fill in the blanks

(i) Sum of the roots of $ax^2 + bx + c = 0$ is _____.

(ii) $x^2 - 4x + 12$ has a pair of real roots if Δ is _____.

(iii) $x^2 - 4x + 12$ has Real roots of $ax^2 + bx + c = 0$ if Δ _____.

(iv) $x^2 - 4x + 12$ has Rational roots of $ax^2 + bx + c = 0$ if Δ _____.

(v) $x^2 - 4x + 12$ has Imaginary roots, Then the roots of $ax^2 + bx + c = 0$ are _____.

(vi) $x^2 - 4x + 12$ is perfect square, then roots of $ax^2 + bx + c = 0$ are _____.

(vii) The roots of $ax^2 + bx + c = 0$ are not perfect square, then the roots of $ax^2 + bx + c = 0$ are _____.

(viii) If α, β are the roots of $ax^2 + bx + c = 0$, then product of the roots is _____.

(ix) If α, β are the roots of $ax^2 + bx + c = 0$, then sum of the roots is _____.

(x) The in quadratic equation $ax^2 + bx + c = 0$, Δ is called _____.

(xi) $x^2 - 4x + 12$ has _____ roots.

(xii) $x^2 - 4x + 12$ has _____ roots.

Hint If Δ is not 0 and the roots of an equation are rational, then Δ is _____.

Answers

(i)	$-\frac{b}{a}$	(ii)	< 0	(iii)	$= 0$	(iv)	$= k^2$	(v)	Imaginary	(vi)	rational
(vii)	Irrational	(viii)	$\frac{c}{a}$	(ix)	$-\frac{b}{a}$	(x)	Discriminant	(xi)	2	(xii)	Imaginary
(xiii)	$\frac{1}{3}$	(xiv)	$\frac{1}{3}$	(xv)	$\frac{15}{7}$	(xvi)	$\frac{25}{7}$	(xvii)	$-\frac{5}{7}$	(xviii)	$\frac{1}{3}$
(xix)	$\frac{1}{3}$	(xx)	$\frac{1}{3}$								

SUMMARY

A **Discriminant** of the quadratic equation $ax^2 + bx + c = 0^{\circ}$, $b^2 - 4ac$.

If **the cube roots** of unity are $1, \omega, \omega^2$ then $1 + \omega + \omega^2 = 0$.

A **Complex cube roots** of unity are $\omega = \frac{-1 + \sqrt{3}i}{2}$, $\omega^2 = \frac{-1 - \sqrt{3}i}{2}$.

The product of complex cube roots of unity is 1.

The sum of the cube roots of unity is zero.

The product of the cube roots of unity is 1.

The sum of the roots of $ax^2 + bx + c = 0$ is $-\frac{b}{a}$.

The **roots** of the quadratic equation $ax^2 + bx + c = 0$ are

$\alpha = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$, $\beta = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$

A **The sum and Then product** of the roots of $ax^2 + bx + c = 0$ are $\alpha + \beta = -\frac{b}{a}$ and $\alpha\beta = \frac{c}{a}$ respectively.

Polynomial Synthetic Division is a process of finding the quotient and remainder, when a polynomial is divided by a linear polynomial.

Synthetic division is a method of finding the quotient and remainder, when a polynomial is divided by a linear polynomial.

Synthetic division Dividing a common factor is called a system of **simultaneous equations.**