EXERCISE 3.1

1. Express the following as a ratio: a : b and as a fraction in its simplest common form.

(i) Rs.750, Rs.1250

Solution:

$$\begin{aligned} &\text{The ratio} \\ &= \text{Rs. } 750 : \text{Rs. } 1250 \\ &= 750 : 1250 \quad \text{(divided by 10)} \\ &= 75 : 125 \\ &= \frac{75}{125} \quad \text{(divided by 5)} \\ &= 15 : 25 \quad \text{(divided by 5)} \\ &= \frac{15}{25} \\ &= 3 : 5 \\ &= \frac{3}{5} \end{aligned}$$

(ii) 400cm, 2m

Solution:

$$\begin{aligned} &\text{400cm : 200cm} \qquad \because 1\text{m} = 100\text{cm} \\ &= 400 : 200 \\ &= 200 : 100 \quad \text{(divided by 10)} \\ &= 2 : 1 \\ &= \frac{2}{1} \end{aligned}$$

(iii) 2kg, 2kg 750gm

Solution:

$$\begin{aligned} &\text{2kg : 2kg 750gm} \\ &= \text{2000g : 2750g} \qquad \because 1\text{kg} = 1000\text{g} \\ &= 2000 : 2750 \\ &= \frac{2000}{2750} \quad \text{(divided by 10)} \\ &= \frac{200}{275} \quad \text{(divided by 5)} \\ &= \frac{40}{55} \quad \text{(divided by 5)} \\ &= 40 : 55 \\ &= \frac{40}{55} \quad \text{(divided by 5)} \\ &= 8 : 11 \\ &= \frac{8}{11} \end{aligned}$$

(iv) 2hrs. 30 sec, 1 hour

Solution:

$$\begin{aligned} &\text{2 hrs. 30sec : 1 hour} \\ &= 7230 \text{ sec : 3600 sec} \qquad (1\text{ hr} = 60 \times 60 = 3600\text{ sec}) \\ &= \frac{7230}{3600} \quad \text{(divided by 10)} \\ &= \frac{723}{360} \\ &= \frac{241}{120} \end{aligned}$$

$$\begin{aligned} &= 9 : 4 \\ &= \frac{9}{4} : 1 \\ &= \frac{9}{4} \\ &= 2\frac{1}{4} \end{aligned}$$

(Divided by 5)

(ii) Rs. 2kg, 750gm

Solution:

$$\begin{aligned} &\text{Rs. 2kg : 750gm} \\ &= \text{Rs. 2000g : 750g} \qquad (1\text{kg} = 1000\text{g}) \\ &= 2000 : 750 \\ &= \frac{2000}{750} \quad \text{(divided by 10)} \\ &= \frac{200}{75} \quad \text{(divided by 5)} \\ &= 200 : 75 \\ &= \frac{200}{75} \quad \text{(divided by 5)} \\ &= 40 : 15 \\ &= \frac{40}{15} \quad \text{(divided by 5)} \\ &= 8 : 3 \\ &= \frac{8}{3} \end{aligned}$$

(iii) 2m, 20 cm

Solution:

$$\begin{aligned} &\text{2m : 20cm} \\ &= 200\text{cm : 20cm} \qquad (1\text{m} = 100\text{cm}) \\ &= 200 : 20 \\ &= 20 : 2 \quad \text{(divided by 10)} \\ &= 10 : 1 \quad \text{(divided by 2)} \\ &= \frac{10}{1} \\ &= 10 \end{aligned}$$

(iv) 1hr. 20 min, 1 hour

Solution:

$$\begin{aligned} &\text{1hr. 20min : 1 hour} \\ &= 4800\text{ sec : 3600 sec} \qquad (1\text{ hr} = 60 \times 60 = 3600\text{ sec}) \\ &= 4800 : 3600 \\ &= \frac{4800}{3600} \quad \text{(divided by 10)} \\ &= \frac{480}{360} \\ &= 480 : 360 \\ &= \frac{480}{360} \quad \text{(divided by 5)} \\ &= 48 : 36 \\ &= \frac{48}{36} \quad \text{(divided by 5)} \\ &= 4 : 3 \\ &= \frac{4}{3} \end{aligned}$$

$$\begin{aligned} &= 625 : 500 \\ &= 125 : 100 \quad \text{(Divided by 5)} \\ &= \frac{125}{100} \\ &= 25 : 20 \quad \text{(Divided by 5)} \\ &= \frac{25}{20} \\ &= 5 : 4 \\ &= \frac{5}{4} \end{aligned}$$

(v) 75°, 225°

Solution:

$$\begin{aligned} &75° : 225° \\ &= 75 : 225 \quad \text{(divided by 5)} \\ &= \frac{75}{225} \\ &= 15 : 45 \quad \text{(divided by 5)} \\ &= \frac{15}{45} \\ &= 1 : 3 \quad \text{(divided by 15)} \\ &= \frac{1}{3} \\ &= 1 : 3 \\ &= \frac{1}{3} \end{aligned}$$

2. In a class of 40 students, 25 students are girls and remaining students are boys. Compute the ratio of

(i) Boys to total students

Solution:

Total students = 40

Number of girls students = 25

Number of boys students = 40 − 25 = 15

$$\begin{aligned} &\text{15 : 40} = \frac{15}{40} \quad \text{(divided by 5)} \\ &= \frac{3}{8} \\ &= 3 : 8 \end{aligned}$$

(ii) Boys to girls

Solution:

$$\begin{aligned} &\text{15 : 25} = \frac{15}{25} \quad \text{(divided by 5)} \\ &= \frac{3}{5} \\ &= 3 : 5 \end{aligned}$$

3. If 2.5a : 3y = 2y : 7y. Find the ratio x : y.

Solution:

$$\begin{aligned} &\text{5}(2x : 2y) = 2y : 7y \\ &\text{5}(2x : 2y) = 2y : 7y \\ &\text{10}x : \text{5}x = 2y : 7y \\ &\frac{10x}{5x} = \frac{2y}{7y} \\ &\frac{2}{1} = \frac{2}{7} \\ &2 \times 7 = 1 \times 2 \\ &14 = 2 \\ &\text{which is not true.} \end{aligned}$$

4. Find the value of y, if the ratio 2y + 1 : 3y + 4 and 3 : 4 are equal.

Solution:

As the given ratios are equal, so

$$\begin{aligned} &2y + 1 : 3y + 4 = 3 : 4 \\ &\frac{2y + 1}{3y + 4} = \frac{3}{4} \\ &4(2y + 1) = 3(3y + 4) \\ &8y + 4 = 9y + 12 \\ &8y - 9y = 12 - 4 \\ &-y = 8 \\ &y = -8 \end{aligned}$$

$$\begin{aligned} &5y + 3y = 30 - 12 \\ &8y = 18 \\ &y = \frac{18}{8} \\ &y = \frac{9}{4} \end{aligned}$$

5. If the ratios 3x + 7 : 4 + 4x and 2 : 3 are equal. Find the value of x.

Solution:

As the given ratios are equal, so

$$\begin{aligned} &3x + 7 : 4 + 4x = 2 : 3 \\ &\frac{3x + 7}{4 + 4x} = \frac{2}{3} \\ &3(3x + 7) = 2(4 + 4x) \\ &9x + 21 = 8 + 8x \\ &9x - 8x = 8 - 21 \\ &x = -13 \end{aligned}$$

Dividing throughout by 7, we get

$$x = 2$$

6. Two numbers are in the ratio 2 : 5. If 5 is added to each number, we get a new ratio 3:7. Find the numbers.

Solution:

Because the ratio of two numbers is 2 : 5

Multiply each number of the ratio with x. Then the numbers be 2x, 5x and the ratio becomes 2x : 5x.

Now according to the given condition, we have

$$\begin{aligned} &\frac{2x + 5}{5x + 5} = \frac{3}{7} \\ &7(2x + 5) = 3(5x + 5) \\ &14x + 35 = 15x + 15 \\ &14x - 15x = 15 - 35 \\ &-x = -20 \\ &x = 20 \end{aligned}$$

Dividing both sides by 7, we get

$$x = 2$$

The required numbers are

$$\begin{aligned} &2(x) = 2(2) = 4 \\ &5(x) = 5(2) = 10 \end{aligned}$$

Dividing throughout by 7, we get

$$x = 3$$

The required numbers are

$$\begin{aligned} &3(x) = 3(3) = 9 \\ &4(x) = 4(3) = 12 \end{aligned}$$

7. If 2/3 is added to each number of the ratio 4 : 13, we get a new ratio 1 : 2. Find the numbers.

Solution:

Because the ratio of two numbers is 4 : 13

multiply each number of the ratio with x. Then the numbers be 4x, 13x and the ratio becomes 4x : 13x.

Now according to the given condition, we have

$$\begin{aligned} &\frac{4x + \frac{2}{3}}{13x + \frac{2}{3}} = \frac{1}{2} \\ &\frac{\frac{12x + 2}{3}}{\frac{39x + 2}{3}} = \frac{1}{2} \\ &\frac{12x + 2}{39x + 2} = \frac{1}{2} \\ &2(12x + 2) = 1(39x + 2) \\ &24x + 4 = 39x + 2 \\ &24x - 39x = 2 - 4 \\ &-15x = -2 \\ &x = \frac{-2}{-15} \\ &x = \frac{2}{15} \end{aligned}$$

(Dividing numerator and denominator by 5)

Dividing throughout by 7, we get

$$x = 2$$

The required numbers are

$$\begin{aligned} &4x = 4(2) = 8 \\ &13x = 13(2) = 26 \end{aligned}$$

8. If 2/3 is added to each number of the ratio 3 : 5, we get a new ratio 7 : 12. Find the numbers.

Solution:

Because the ratio of two numbers is 3 : 5

multiply each number of the ratio with x. Then the numbers be 3x, 5x and the ratio becomes 3x : 5x.

Now according to the given condition, we have

$$\begin{aligned} &\frac{3x + \frac{2}{3}}{5x + \frac{2}{3}} = \frac{7}{12} \\ &\frac{\frac{9x + 2}{3}}{\frac{15x + 2}{3}} = \frac{7}{12} \\ &\frac{9x + 2}{15x + 2} = \frac{7}{12} \\ &12(9x + 2) = 7(15x + 2) \\ &108x + 24 = 105x + 14 \\ &108x - 105x = 14 - 24 \\ &3x = -10 \\ &x = \frac{-10}{3} \end{aligned}$$

$$\begin{aligned} &\frac{3x + 2y}{3x - 2y} = \frac{5}{1} \\ &\frac{3x + 2y}{3x - 2y} = \frac{5}{1} \\ &\text{Hence,} \\ &3x + 2y : 3x - 2y = 5 : 1 \end{aligned}$$

10. Complete the following:

(i) $\frac{a}{b} = \frac{c}{d}$ **, then** $\frac{a+b}{a-b} =$ _____

Solution:

$$\begin{aligned} &\frac{a}{b} = \frac{c}{d} \\ &a(d) = c(b) \quad \text{(Multiplying by } bd\text{)} \\ &ad = bc \quad \text{(Multiplying by } \frac{1}{ab}\text{)} \\ &\frac{ad}{ab} = \frac{bc}{ab} \\ &\frac{d}{b} = \frac{c}{a} \end{aligned}$$

(ii) $\frac{3x}{5y} = \frac{7z}{2w}$ **, then** $xw : yz =$ _____

Solution:

$$\begin{aligned} &\frac{3x}{5y} = \frac{7z}{2w} \\ &\frac{x}{y} = \frac{7z}{2w} \times \frac{5}{3} \\ &\frac{x}{y} = \frac{35z}{6w} \\ &xw : yz = 35 : 6 \end{aligned}$$

(iii) $\frac{x}{y} = \frac{7w}{2w}$ **, then** $5x + 2y =$ _____

Solution:

$$\begin{aligned} &\frac{x}{y} = \frac{7w}{2w} \\ &\frac{5x}{5y} = \frac{7w}{2w} \\ &\frac{5x + 2y}{5y - 2y} = \frac{7w + 2w}{7w - 2w} \\ &\frac{5x + 2y}{3y} = \frac{9w}{5w} \\ &\frac{5x + 2y}{3y} = \frac{9}{5} \\ &5x + 2y : 3y = 9 : 5 \end{aligned}$$

11. Find a in the following proportions.

(i) 3a : 4 :: 2a + 1 : 7

Solution:

$$\begin{aligned} &3a : 4 :: 2a + 1 : 7 \\ &3a : 4 = 2a + 1 : 7 \\ &\frac{3a}{4} = \frac{2a + 1}{7} \end{aligned}$$

Product of extremes = Product of means

$$\begin{aligned} &21a = 4(2a + 4) \\ &21a = 8a + 4 \\ &21a - 8a = 4 \\ &13a = 4 \end{aligned}$$

Dividing both sides by 13, we get

$$a = \frac{4}{13}$$

(ii) $\frac{a-2}{a+2} :: \frac{a-1}{a+4}$

Solution:

$$\begin{aligned} &\frac{a-2}{a+2} :: \frac{a-1}{a+4} \\ &\frac{a-2}{a+2} = \frac{a-1}{a+4} \end{aligned}$$

Product of extremes = Product of means

$$\begin{aligned} &(a-2)(a+4) = (a+2)(a-1) \\ &a(a+4) - 2(a+4) = a(a-1) + 2(a-1) \\ &a^2 + 4a - 2a - 8 = a^2 - a + 2a - 2 \\ &a^2 + 2a - 8 = a^2 + a - 2 \\ &a^2 + 2a - a^2 - a = -2 + 8 \\ &a = 6 \end{aligned}$$

(iii) $5 : x :: 11 : x + 24 :: a : 25 + x$

Solution:

$$\begin{aligned} &5 : x :: 11 : x + 24 \\ &5 : x = 11 : x + 24 \\ &\frac{5}{x} = \frac{11}{x + 24} \end{aligned}$$

Product of extremes = Product of means

$$\begin{aligned} &5(x + 24) = 11x \\ &5x + 120 = 11x \\ &5x - 11x = -120 \\ &-6x = -120 \\ &x = \frac{-120}{-6} \\ &x = 20 \end{aligned}$$

$$\begin{aligned} &x : 11 :: a : x + 24 \\ &\frac{x}{11} = \frac{a}{x + 24} \end{aligned}$$

Product of extremes = Product of means

$$\begin{aligned} &x(x + 24) = 11a \\ &x^2 + 24x = 11a \\ &20^2 + 24(20) = 11a \quad (\because x = 20) \\ &400 + 480 = 11a \\ &880 = 11a \\ &\frac{880}{11} = a \\ &80 = a \\ &a = 80 \end{aligned}$$

$$\begin{aligned} &x : 11 :: a : x + 24 \\ &\frac{x}{11} = \frac{a}{x + 24} \end{aligned}$$

Product of extremes = Product of means

$$\begin{aligned} &x(x + 24) = 11a \\ &x^2 + 24x = 11a \\ &(-30)^2 + 24(-30) = 11a \quad (\because x = -30) \\ &900 - 720 = 11a \\ &180 = 11a \\ &\frac{180}{11} = a \\ &a = \frac{180}{11} \end{aligned}$$

$$\begin{aligned} &5(3x - 6) = 3(x + 4) \\ &15x - 30 = 3x + 12 \\ &15x - 3x = 30 + 12 \\ &12x = 42 \end{aligned}$$

$$x = \frac{42}{12}$$

$$x = \frac{7}{2}$$

$$\begin{aligned} &y^2 + pq + q^2 = \frac{p^2 - q^2}{p + q}(y - q) \\ &\text{(ii)} \\ &\text{Solution} \end{aligned}$$

$$\begin{aligned} &y^2 + pq + q^2 = \frac{p^2 - q^2}{p + q}(y - q) \\ &y^2 + pq + q^2 = \frac{(p + q)(p - q)}{p + q}(y - q) \end{aligned}$$

Product of means = Product of extremes

$$\begin{aligned} &(p + q)\left(y^2 + pq + q^2\right) = (p - q)(y - q) \\ &(p + q)\left(y^2 + pq + q^2\right) = (p - q)(y - q) \\ &(p + q)\left(y^2 + pq + q^2\right) = p(y - q) - q(y - q) \\ &(p + q)\left(y^2 + pq + q^2\right) = py - pq - qy + q^2 \\ &py - pq - qy + q^2 - (p + q)\left(y^2 + pq + q^2\right) = 0 \\ &py - pq - qy + q^2 - py^2 - p^2q - pq^2 - qy^2 - pq^2 - q^3 = 0 \\ &-py^2 - qy^2 - p^2q - 2pq^2 - q^3 = 0 \\ &-y^2(p + q) - pq(p + 2q) - q^3 = 0 \\ &y^2(p + q) + pq(p + 2q) + q^3 = 0 \end{aligned}$$

(iv) $5 : x :: 11 : x + 24 :: a : 25 + x$

Solution:

$$\begin{aligned} &5 : x :: 11 : x + 24 \\ &5 : x = 11 : x + 24 \\ &\frac{5}{x} = \frac{11}{x + 24} \end{aligned}$$

Product of extremes = Product of means

$$\begin{aligned} &5(x + 24) = 11x \\ &5x + 120 = 11x \\ &5x - 11x = -120 \\ &-6x = -120 \\ &x = \frac{-120}{-6} \\ &x = 20 \end{aligned}$$

$$\begin{aligned} &x : 11 :: a : x + 24 \\ &\frac{x}{11} = \frac{a}{x + 24} \end{aligned}$$

Product of extremes = Product of means

$$\begin{aligned} &x(x + 24) = 11a \\ &x^2 + 24x = 11a \\ &20^2 + 24(20) = 11a \quad (\because x = 20) \\ &400 + 480 = 11a \\ &880 = 11a \\ &\frac{880}{11} = a \\ &80 = a \\ &a = 80 \end{aligned}$$

$$\begin{aligned} &x : 11 :: a : x + 24 \\ &\frac{x}{11} = \frac{a}{x + 24} \end{aligned}$$

Product of extremes = Product of means

$$\begin{aligned} &x(x + 24) = 11a \\ &x^2 + 24x = 11a \\ &(-30)^2 + 24(-30) = 11a \quad (\because x = -30) \\ &900 - 720 = 11a \\ &180 = 11a \\ &\frac{180}{11} = a \\ &a = \frac{180}{11} \end{aligned}$$

(c) Variation

Two or more variables is frequently used in all sciences. There are two types of variations:

(i) Direct variation **(ii)** Inverse variation.

(i) Direct Variation

Two variables are related in such a way that increase (decrease) in one quantity causes increase (decrease) in the other quantity. Then the variation is called direct variation.

In other words, if a quantity y varies directly with regard to a quantity x, We say that y is directly proportional to x and is written as $y \propto x$ or $y = kx$, $k \neq 0$.

The sign ' $\propto$ ' stand as "varies as" is called the sign of proportionality or variation, where k is known as constant of variation.

Note

(i) Consider the speed of a car, longer the distance it covers.

(ii) The smaller the radius of the circle, smaller the circumference is.

(ii) Inverse Variation

If two quantities are related in such a way that when one quantity increases, the other decreases, then the variation is called inverse variation.

In other words, if a quantity y varies inversely with regard to quantity x, We say that y is inversely proportional to x or y varies inversely as x and is written as $y \propto \frac{1}{x}$

i.e., $xy = k$, where k is it the constant of variation.

Exercise 3.2

1. If y varies directly as x, and y = 6 when x = 1, find

(i) y in terms of x.

Solution:

Given that y varies directly as x.

Therefore $y \propto x$

$\Rightarrow \quad y = kx \quad \ldots\ldots(i)$

Where k is constant of variation.

Put $x = 2$ and $y = 8$ in eq. (i), we have

$8 = k(2)$

$\Rightarrow \quad 2k = 8$

$\Rightarrow \quad k = 4$

Put $k = 4$ in eq. (i), we get

$y = 4x$

(ii) y when x = 8

Solution:

Given that y varies directly as x.

Therefore $y \propto x$

$\Rightarrow \quad y = kx \quad \ldots\ldots(i)$

Where k is constant of variation.

Put $x = 2$ and $y = 8$ in eq. (i), we have

$8 = k(2)$

$\Rightarrow \quad 2k = 8$

$\Rightarrow \quad k = 4$

Put $k = 4$ and $x = 8$ in eq. (i), we get

$y = 4(8)$

$\Rightarrow \quad y = 32$

2. If $x \propto \sqrt{y}$ and $y = 7$ when $x = 28$ find

(i) x in terms of y

Solution:

Given that x varies directly as $\sqrt{y}$.

Therefore $x \propto \sqrt{y}$

$\Rightarrow \quad x = k\sqrt{y} \quad \ldots\ldots(i)$

Where k is constant of variation.

Put $y = 7$ and $x = 28$ in eq. (i), we get

$28 = k\sqrt{7}$

$\Rightarrow \quad k = \frac{28}{\sqrt{7}}$

$\Rightarrow \quad k = \frac{28\sqrt{7}}{7}$

$\Rightarrow \quad k = 4\sqrt{7}$

Put $k = 4\sqrt{7}$ in eq. (i), we get

$x = 4\sqrt{7}\sqrt{y}$

$x = 4\sqrt{7y}$

$\Rightarrow \quad y = 7 \times 49 \quad \ldots\ldots(i)$

Where k is constant of variation.

Put $x = 3$ and $y = 7 \times 9$ in eq. (i), we have

$7 \times 9 = k(3)$

$\Rightarrow \quad 3k = 7 \times 9$

$\Rightarrow \quad k = \frac{7 \times 9}{3}$

Put $k = \frac{7 \times 9}{3}$ and $x = 49$ in eq. (i), we get

$y = \frac{7 \times 9}{3}(49)$

$y = \frac{3 \times 147}{1}$

$y = 441$

(ii) x when y = 20 and y when x = 12

Solution:

Given that y varies directly as x.

Therefore $y \propto x$

$\Rightarrow \quad y = kx \quad \ldots\ldots(i)$

Where k is constant of variation.

Put $x = 3$ and $y = 7 \times 9$ in eq. (i), we have

$7 \times 9 = k(3)$

$\Rightarrow \quad 3k = 7 \times 9$

$\Rightarrow \quad k = \frac{7 \times 9}{3}$

Put $k = \frac{7 \times 9}{3}$ and $y = 20$ in eq. (i), we get

$20 = \frac{7 \times 9}{3}x$

$\Rightarrow \quad 3 \times 20 = 7x$

$\Rightarrow \quad 60 = 7x$

$\Rightarrow \quad x = \frac{60}{7}$

Put $x = \frac{60}{7}$ and $k = \frac{7 \times 9}{3}$ in eq. (i), we get

$y = \frac{7 \times 9}{3}\left(\frac{60}{7}\right)$

$\Rightarrow \quad y = \frac{3 \times 60}{1}$

$\Rightarrow \quad y = 180$

3. If $R \propto \frac{1}{T}$ and $R = 5$ when $T = 6$, find the equation connecting R and T. Also find R when $T = 20$ and T when $R = 25$.

Solution:

Given that $R \propto \frac{1}{T}$

$\Rightarrow \quad R = k\frac{1}{T} \quad \ldots\ldots(i)$

Put $R = 5$ and $T = 6$ in eq. (i), we get

$5 = k\left(\frac{1}{6}\right)$

$\Rightarrow \quad 5k = 1$

$\Rightarrow \quad k = \frac{1}{5}$

Put $k = \frac{1}{5}$ in eq. (i), we get

$R = \frac{1}{5}\left(\frac{1}{T}\right) \quad \ldots\ldots(ii)$

Put $T = 20$ and $R = 25$ in eq. (ii), we get

$R = \frac{1}{5}\left(\frac{1}{20}\right)$

$\Rightarrow \quad R = \frac{1}{100}$

Put $R = \frac{1}{100}$ and $k = \frac{1}{5}$ in eq. (i), we get

$\frac{1}{100} = \frac{1}{5}\left(\frac{1}{T}\right)$

$\Rightarrow \quad \frac{1}{100} = \frac{1}{5T}$

$\Rightarrow \quad 5T = 100$

$\Rightarrow \quad T = \frac{100}{5}$

$\Rightarrow \quad T = 20$

4. If $R \propto \frac{1}{T^2}$ and $R = 4$ when $T = 5$. find R when T = 4.

Solution:

Given that $R \propto \frac{1}{T^2}$

$\Rightarrow \quad R = k\frac{1}{T^2} \quad \ldots\ldots(i)$

Put $R = 4$ and $T = 5$ in eq. (i), we get

$4 = k\left(\frac{1}{5^2}\right)$

$\Rightarrow \quad 4 = k\left(\frac{1}{25}\right)$

$\Rightarrow \quad 4 \times 25 = k$

$\Rightarrow \quad k = 100$

Put $k = 100$ in eq. (i), we get

$R = 100\left(\frac{1}{T^2}\right)$

$\Rightarrow \quad R = \frac{100}{T^2}$

Put $T = 4$ in eq. (ii), we get

$R = \frac{100}{4^2}$

$\Rightarrow \quad R = \frac{100}{16}$

$\Rightarrow \quad R = 6.25$

5. If $Y \propto \frac{1}{X}$ and $Y = 4$ when $X = 3$, find X, when $Y = 125$.

Solution:

Given that $Y \propto \frac{1}{X}$

$\Rightarrow \quad Y = k\frac{1}{X} \quad \ldots\ldots(i)$

Put $Y = 4$ and $X = 3$ in eq. (i), we get

$4 = k\left(\frac{1}{3}\right)$

$\Rightarrow \quad k = 12$

Put $k = 12$ in eq. (i), we get

$Y = 12\left(\frac{1}{X}\right)$

$\Rightarrow \quad Y = \frac{12}{X}$

Put $Y = 125$ in eq. (ii), we get

$125 = \frac{12}{X}$

$\Rightarrow \quad 125X = 12$

$\Rightarrow \quad X = \frac{12}{125}$

6. If $y \propto \frac{1}{x^2}$ and $y = 9$ when $x = 2$, find

(i) y when $x = \frac{1}{3}$

Solution:

Given that $y \propto \frac{1}{x^2}$

$\Rightarrow \quad y = k\frac{1}{x^2} \quad \ldots\ldots(i)$

Put $y = 9$ and $x = 2$ in eq. (i), we get

$9 = k\left(\frac{1}{2^2}\right)$

$\Rightarrow \quad 9 = k\left(\frac{1}{4}\right)$

$\Rightarrow \quad 9 \times 4 = k$

$\Rightarrow \quad k = 36$

Put $k = 36$ and $x = \frac{1}{3}$ in eq. (i), we get

$y = 36\left(\frac{1}{\frac{1}{3}}\right)^2$

$\Rightarrow \quad y = 36(3)^2$

$\Rightarrow \quad y = 36 \times 9$

$\Rightarrow \quad y = 324$

(ii) x when $y = \frac{1}{4}$

Solution:

Given that $y \propto \frac{1}{x^2}$

$\Rightarrow \quad y = k\frac{1}{x^2} \quad \ldots\ldots(i)$

Put $y = 9$ and $x = 2$ in eq. (i), we get

$9 = k\left(\frac{1}{2^2}\right)$

$\Rightarrow \quad 9 = k\left(\frac{1}{4}\right)$

$\Rightarrow \quad 9 \times 4 = k$

$\Rightarrow \quad k = 36$

Put $k = 36$ and $y = \frac{1}{4}$ in eq. (i), we get

$\frac{1}{4} = 36\left(\frac{1}{x^2}\right)$

$\Rightarrow \quad \frac{1}{4} = \frac{36}{x^2}$

$\Rightarrow \quad x^2 = 4 \times 36$

$\Rightarrow \quad x^2 = 144$

$\Rightarrow \quad x = \sqrt{144}$

$\Rightarrow \quad x = \pm 12$

7. If $y \propto \frac{1}{x}$ and $x = 4$ when $y = 3$, find x when $y = 6$.

Solution:

Given that $y \propto \frac{1}{x}$

$\Rightarrow \quad y = k\frac{1}{x} \quad \ldots\ldots(i)$

Put $y = 3$ and $x = 4$ in eq. (i), we get

$3 = k\left(\frac{1}{4}\right)$

$\Rightarrow \quad 3 = \frac{k}{4}$

$\Rightarrow \quad k = 12$

Put $k = 12$ and $y = 6$ in eq. (i), we get

$6 = \frac{12}{x}$

$\Rightarrow \quad 6x = 12$

$\Rightarrow \quad x = \frac{12}{6}$

$\Rightarrow \quad x = 2$

8. If $y \propto \frac{1}{x}$ and $y = 4$ when $x = 2$, find x when $y = 25$.

Solution:

Given that $y \propto \frac{1}{x}$

$\Rightarrow \quad y = k\frac{1}{x} \quad \ldots\ldots(i)$

Put $y = 4$ and $x = 2$ in eq. (i), we get

$4 = k\left(\frac{1}{2}\right)$

$\Rightarrow \quad 4 = \frac{k}{2}$

$\Rightarrow \quad k = 8$

Put $k = 8$ and $y = 25$ in eq. (i), we get

$25 = \frac{8}{x}$

$\Rightarrow \quad 25x = 8$

$\Rightarrow \quad x = \frac{8}{25}$

9. If $y \propto \frac{1}{x^2}$ and $y = 4$ when $x = 3$, find x when $y = \frac{4}{9}$.

Solution:

Given that $y \propto \frac{1}{x^2}$

$\Rightarrow \quad y = k\frac{1}{x^2} \quad \ldots\ldots(i)$

Put $y = 4$ and $x = 3$ in eq. (i), we get

$4 = k\left(\frac{1}{3^2}\right)$

$\Rightarrow \quad 4 = k\left(\frac{1}{9}\right)$

$\Rightarrow \quad 4 \times 9 = k$

$\Rightarrow \quad k = 36$

Put $k = 36$ and $y = \frac{4}{9}$ in eq. (i), we get

$\frac{4}{9} = 36\left(\frac{1}{x^2}\right)$

$\Rightarrow \quad \frac{4}{9} = \frac{36}{x^2}$

$\Rightarrow \quad 4x^2 = 9 \times 36$

$\Rightarrow \quad x^2 = \frac{9 \times 36}{4}$

$\Rightarrow \quad x^2 = 81$

$\Rightarrow \quad x = \sqrt{81}$

$\Rightarrow \quad x = \pm 9$

10. If $x \propto \frac{1}{y^2}$ and $x = 4$ when $y = 3$, find y when $x = 72$.

Solution:

Given that $x \propto \frac{1}{y^2}$

$\Rightarrow \quad x = k\frac{1}{y^2} \quad \ldots\ldots(i)$

Put $x = 4$ and $y = 3$ in eq. (i), we get

$4 = k\left(\frac{1}{3^2}\right)$

$\Rightarrow \quad 4 = k\left(\frac{1}{9}\right)$

$\Rightarrow \quad 4 \times 9 = k$

$\Rightarrow \quad k = 36$

Put $k = 36$ and $x = 72$ in eq. (i), we get

$72 = \frac{36}{y^2}$

$\Rightarrow \quad y^2 = \frac{36}{72}$

$\Rightarrow \quad y^2 = \frac{1}{2}$

$\Rightarrow \quad y = \sqrt{\frac{1}{2}}$

11. If $x \propto \frac{1}{y^2}$ and $x = 9$ when $y = 4$. find x when $y = 6$.

Solution:

Given that $x \propto \frac{1}{y^2}$

$\Rightarrow \quad x = k\frac{1}{y^2} \quad \ldots\ldots(i)$

Put $x = 9$ and $y = 4$ in eq. (i), we get

$9 = k\left(\frac{1}{4^2}\right)$

$\Rightarrow \quad 9 = k\left(\frac{1}{16}\right)$

$\Rightarrow \quad 9 \times 16 = k$

$\Rightarrow \quad k = 144$

Put $k = 144$ and $y = 6$ in eq. (i), we get

$x = \frac{144}{6^2}$

$\Rightarrow \quad x = \frac{144}{36}$

$\Rightarrow \quad x = 4$

12. If $x \propto \frac{1}{y^2}$ and $x = 1$ when $y = 3$, find x when $y = 12$.

Solution:

Given that $x \propto \frac{1}{y^2}$

$\Rightarrow \quad x = k\frac{1}{y^2} \quad \ldots\ldots(i)$

Put $x = 1$ and $y = 3$ in eq. (i), we get

$1 = k\left(\frac{1}{3^2}\right)$

$\Rightarrow \quad 1 = k\left(\frac{1}{9}\right)$

$\Rightarrow \quad 1 \times 9 = k$

$\Rightarrow \quad k = 9$

Put $k = 9$ and $y = 12$ in eq. (i), we get

$x = \frac{9}{12^2}$

$\Rightarrow \quad x = \frac{9}{144}$

$\Rightarrow \quad x = \frac{1}{16}$

13. If $x \propto \frac{1}{y^2}$ and $x = 9$ when $y = 4$. find x when $y = 6$ and y when $x = 400$.

Solution:

Given that $x \propto \frac{1}{y^2}$

$\Rightarrow \quad x = k\frac{1}{y^2} \quad \ldots\ldots(i)$

Put $x = 9$ and $y = 4$ in eq. (i), we get

$9 = k\left(\frac{1}{4^2}\right)$

$\Rightarrow \quad 9 = k\left(\frac{1}{16}\right)$

$\Rightarrow \quad 9 \times 16 = k$

$\Rightarrow \quad k = 144$

Put $k = 144$ and $x = 4$ in eq. (i), we get

$x = \frac{144}{6^2}$

$\Rightarrow \quad x = \frac{144}{36}$

$\Rightarrow \quad x = 4$

Put $x = 400$ and $k = 144$ in eq. (i), we get

$400 = \frac{144}{y^2}$

$\Rightarrow \quad 400y^2 = 144$

$\Rightarrow \quad y^2 = \frac{144}{400}$

$\Rightarrow \quad y = \sqrt{\frac{144}{400}}$

$\Rightarrow \quad y = \pm\frac{12}{20}$

14. If $x \propto \frac{1}{y^2}$ and $x = 1$ when $y = 12$. find x when $y = 120$.

Solution:

Given that $x \propto \frac{1}{y^2}$

$\Rightarrow \quad x = k\frac{1}{y^2} \quad \ldots\ldots(i)$

Put $x = 1$ and $y = 12$ in eq. (i), we get

$1 = k\left(\frac{1}{12^2}\right)$

$\Rightarrow \quad 1 = k\left(\frac{1}{144}\right)$

$\Rightarrow \quad 1 \times 144 = k$

$\Rightarrow \quad k = 144$

Put $k = 144$ and $y = 120$ in eq. (i), we get

$x = \frac{144}{120^2}$

$\Rightarrow \quad x = \frac{144}{14400}$

$\Rightarrow \quad x = \frac{1}{100}$

$\Rightarrow \quad x = \frac{1}{y^2}$

$\Rightarrow \quad x = \frac{1}{y^2}$

$\Rightarrow \quad x = \frac{1}{y^2}$

Put $x = 100$ and $k = 144$ in eq. (i), we get

$100 = \frac{144}{y^2}$

$\Rightarrow \quad 100y^2 = 144$

$\Rightarrow \quad y^2 = \frac{144}{100}$

$\Rightarrow \quad y = \sqrt{\frac{144}{100}}$

$\Rightarrow \quad y = \pm\frac{12}{10}$

15. If $x \propto \frac{1}{y^2}$ and $x = 4$ when $y = 3$. find x when $y = 6$.

Solution:

Given that $x \propto \frac{1}{y^2}$

$\Rightarrow \quad x = k\frac{1}{y^2} \quad \ldots\ldots(i)$

Put $x = 4$ and $y = 3$ in eq. (i), we get

$4 = k\left(\frac{1}{3^2}\right)$

$\Rightarrow \quad 4 = k\left(\frac{1}{9}\right)$

$\Rightarrow \quad 4 \times 9 = k$

$\Rightarrow \quad k = 36$

Put $k = 36$ and $y = 6$ in eq. (i), we get

$x = \frac{36}{6^2}$

$\Rightarrow \quad x = \frac{36}{36}$

$\Rightarrow \quad x = 1$

16. If $x \propto \frac{1}{y^2}$ and $x = 1$ when $y = 12$. find x when $y = 120$.

Solution:

Given that $x \propto \frac{1}{y^2}$

$\Rightarrow \quad x = k\frac{1}{y^2} \quad \ldots\ldots(i)$

Put $x = 1$ and $y = 12$ in eq. (i), we get

$1 = k\left(\frac{1}{12^2}\right)$

$\Rightarrow \quad 1 = k\left(\frac{1}{144}\right)$

$\Rightarrow \quad 1 \times 144 = k$

$\Rightarrow \quad k = 144$

Put $k = 144$ and $y = 120$ in eq. (i), we get

$x = \frac{144}{120^2}$

$\Rightarrow \quad x = \frac{144}{14400}$

$\Rightarrow \quad x = \frac{1}{100}$

$\Rightarrow \quad x = \frac{1}{y^2}$

$\Rightarrow \quad x = \frac{1}{y^2}$

$\Rightarrow \quad x = \frac{1}{y^2}$

Put $x = 100$ and $k = 144$ in eq. (i), we get

$100 = \frac{144}{y^2}$

$\Rightarrow \quad 100y^2 = 144$

$\Rightarrow \quad y^2 = \frac{144}{100}$

$\Rightarrow \quad y = \sqrt{\frac{144}{100}}$

$\Rightarrow \quad y = \pm\frac{12}{10}$

17. If $x \propto \frac{1}{y^2}$ and $x = 9$ when $y = 4$. find x when $y = 6$ and y when $x = 400$.

Solution:

Given that $x \propto \frac{1}{y^2}$

$\Rightarrow \quad x = k\frac{1}{y^2} \quad \ldots\ldots(i)$

Put $x = 9$ and $y = 4$ in eq. (i), we get

$9 = k\left(\frac{1}{4^2}\right)$

$\Rightarrow \quad 9 = k\left(\frac{1}{16}\right)$

$\Rightarrow \quad 9 \times 16 = k$

$\Rightarrow \quad k = 144$

Put $k = 144$ and $x = 4$ in eq. (i), we get

$x = \frac{144}{6^2}$

$\Rightarrow \quad x = \frac{144}{36}$

$\Rightarrow \quad x = 4$

Put $x = 400$ and $k = 144$ in eq. (i), we get

$400 = \frac{144}{y^2}$

$\Rightarrow \quad 400y^2 = 144$

$\Rightarrow \quad y^2 = \frac{144}{400}$

$\Rightarrow \quad y = \sqrt{\frac{144}{400}}$

$\Rightarrow \quad y = \pm\frac{12}{20}$

Find 3^{rd}, 4^{th}, mean and continued proportion:

We are already familiar with proportions that if quantities a, b, c and d are in proportion, then a : b :: c : d

i.e., product of extremes = product of means

Third Proportional

If three quantities a, b and c are related as a : b :: b : c, then c is called the third proportion.

Fourth Proportional

If four quantities a, b, c and d are related as a: b :: c : d

Then d is called the fourth proportional.

Mean Proportional

If three quantities a, b and c are related as a : b :: b : c, then b is called the mean proportional.

Continued Proportion

If three quantities a, b and c are related as

a : b :: b : c

where a is first, b is the mean and c is the third proportional, then a, b and c are in continued proportion.

Exercise 3.3

1. Find a third proportional to

(i) 6, 12

Solution:

Let C be the third proportional, then

6:12 :: 12:C

∴ Product of extremes = Product of means

6C = 12 x 12

6C = 144

$C = \frac{144}{6}$

C = 24

(ii) $a^3, 3a^2$

Solution:

Let C be the third proportional, then

$a^3 : 3a^2 :: 3a^2 : C$

∴ Product of extremes = Product of means

$(C)(a^3) = (3a^2)(3a^2)$

$C = \frac{(3a^2)(3a^2)}{a^3}$

$C = \frac{9a^4}{a^3}$

$C = 9a$

(iii) $a^2 - b^2, a - b$

Solution:

Let C be the third proportional, then

$a^2 - b^2 : a - b :: a - b : C$

∴ Product of extremes = Product of means

$(a^2 - b^2)(C) = (a-b)(a-b)$

$C = \frac{(a-b)(a-b)}{(a^2-b^2)}$

$C = \frac{(a-b)(a-b)}{(a-b)(a+b)}$

$C = \frac{(a-b)}{(a+b)}$

(iv) $(x-y)^2, x^3 - y^3$

Solution:

Let C be the third proportional, then

$(x-y)^2 : x^3 - y^3 :: x^3 - y^3 : C$

∴ Product of extremes = Product of means

$(C)(x-y)^2 = (x^3-y^3)(x^3-y^3)$

$C = \frac{(x-y)(x^2+xy+y^2)(x-y)(x^2+xy+y^2)}{(x-y)^2}$

$C = \frac{(x-y)^2(x^2+xy+y^2)^2}{(x-y)^2}$

$C = (x^2+xy+y^2)^2$

(v) $(x+y)^2, x^2 - xy - 2y^2$

Solution:

Let C be the third proportional, then

$(x+y)^2 : x^2 - xy - 2y^2 :: x^2 - xy - 2y^2 : C$

∴ Product of extremes = Product of means

$(C)(x+y)^2 = (x^2-xy-2y^2)(x^2-xy-2y^2)$

$C = \frac{(x^2-xy-2y^2)^2}{(x+y)^2}$

$C = \frac{(x^2-2xy+xy-2y^2)^2}{(x+y)^2}$

$C = \frac{[x(x-2y)+y(x-2y)]^2}{(x+y)^2}$

$C = \frac{[(x-2y)(x+y)]^2}{(x+y)^2}$

$C = \frac{(x-2y)^2(x+y)^2}{(x+y)^2}$

$C = (x-2y)^2$

(vi) $\frac{p^2-q^2}{p^3+q^3}, \frac{p-q}{p^2-pq+q^2}$

Solution:

Let C be the third proportional, then

$\frac{p^2-q^2}{p^3+q^3} : \frac{p-q}{p^2-pq+q^2} :: \frac{p-q}{p^2-pq+q^2} : C$

∴ Product of extremes = Product of means

$(C)\left(\frac{p^2-q^2}{p^3+q^3}\right) = \left(\frac{p-q}{p^2-pq+q^2}\right)\left(\frac{p-q}{p^2-pq+q^2}\right)$

$C = \frac{(p-q)^2}{(p^2-pq+q^2)^2} \times \frac{p^3+q^3}{p^2-q^2}$

$C = \frac{(p-q)^2}{(p^2-pq+q^2)^2} \times \frac{(p+q)(p^2-pq+q^2)}{(p+q)(p-q)}$

$C = \frac{p-q}{p^2-pq+q^2}$

2. Find a fourth proportional to

(i) 5, 8, 15

Solution:

Let x be the fourth proportional, then

$5 : 8 :: 15 : x$

∴ Product of extremes = Product of means

$(5)(x) = (8)(15)$

$x = \frac{(8)(15)}{(5)}$

$x = 8 \times 3 = 24$

(ii) $4x^4, 2x^3, 18x^5$

Solution:

[redacted]

Let C be the fourth proportional, then

$4x^4 : 2x^3 :: 18x^5 : C$

∴ Product of extremes = Product of means

$(4x^4)(C) = (2x^3)(18x^5)$

$C = \frac{36x^8}{4x^4}$

$C = 9x^4$

(iii) $15a^5b^6, 10a^2b^5, 21a^3b^3$

Solution:

Let x be the fourth proportional, then

$15a^5b^6 : 10a^2b^5 :: 21a^3b^3 : x$

∴ Product of extremes = Product of means

$x(15a^5b^6) = (10a^2b^5)(21a^3b^3)$

$x = \frac{10 \times 21a^5b^8}{15a^5b^6}$

$x = 2 \times 7b^2$

$x = 14b^2$

(iv) $x^2 - 11x + 24, (x-3), 5x^4 - 40x^3$

Solution:

Let x be the fourth proportional, then

$x^2 - 11x + 24 : (x-3) :: 5x^4 - 40x^3 : C$

∴ Product of extremes = Product of means

$C(x^2-11x+24) = (x-3)(5x^4-40x^3)$

$C = \frac{(x-3)[5x^3(x-8)]}{x^2-11x+24}$

$C = \frac{5x^3(x-3)(x-8)}{x^2-8x-3x+24}$

$C = \frac{5x^3(x-3)(x-8)}{x(x-8)-3(x-8)}$

$C = \frac{5x^3(x-3)(x-8)}{(x-3)(x-8)}$

$C = 5x^3$

(v) $p^3+q^3, p^2-q^2, p^2-pq+q^2$

Solution:

Let C be the fourth proportional, then

$p^3+q^3 : p^2-q^2 :: p^2-pq+q^2 : C$

∴ Product of extremes = Product of means

$C(p^3+q^3) = (p^2-q^2)(p^2-pq+q^2)$

$C = \frac{(p-q)(p+q)(p^2-pq+q^2)}{(p^3+q^3)}$

$C = \frac{(p-q)(p^3+q^3)}{(p^3+q^3)}$

$C = p - q$

(vi) $(p^2-q^2)(p^2+pq+q^2), p^3+q^3, p^3-q^3$

Solution:

Let x be the fourth proportional, then

$(p^2-q^2)(p^2+pq+q^2) : p^3+q^3 :: p^3-q^3 : x$

∴ Product of extremes = Product of means

$x(p^2-q^2)(p^2+pq+q^2) = (p^3+q^3)(p^3-q^3)$

$x = \frac{(p^3+q^3)(p^3-q^3)}{(p^2-q^2)(p^2+pq+q^2)}$

$x = \frac{(p^3+q^3)(p^3-q^3)}{(p+q)(p-q)(p^2+pq+q^2)}$

$C = \frac{(p+q)(p^2-pq+q^2)(p^3-q^3)}{(p+q)(p^3-q^3)}$

$C = p^2 - pq + q^2$

3. Find a mean proportional between

(i) 20,45

Solution:

Let m be the mean proportional, then

$20 : m :: m : 45$

∴ Product of means = Product of extremes

$m \times m = 20 \times 45$

$m^2 = 900$

$m = \pm\sqrt{900}$

$m = \pm 30$

(ii) $20x^3y^5, 5x^7y$

Solution:

Let m be the mean proportional, then

$20x^3y^5 : m :: m : 5x^7y$

∴ Product of means = Product of extremes

$m \times m = 20x^3y^5 \times 5x^7y$

$m^2 = 100x^{10}y^6$

$m = \pm\sqrt{100x^{10}y^6}$

$m = \pm(100x^{10}y^6)^{\frac{1}{2}}$

$m = \pm(10^2)^{\frac{1}{2}}(x^{10})^{\frac{1}{2}}(y^6)^{\frac{1}{2}}$

$m = \pm 10x^5y^3$

(iii) $15p^4qr^3, 135q^5r^7$

Solution:

Let m be the mean proportional, then

$15p^4qr^3 : m :: m : 135q^5r^7$

∴ Product of means = Product of extremes

$m \times m = 15p^4qr^3 \times 135q^5r^7$

$m \times m = 15 \times 135p^4q^6r^{10}$

$m^2 = 2025p^4q^6r^{10}$

$m = \pm\sqrt{2025p^4q^6r^{10}}$

$m = \pm(2025p^4q^6r^{10})^{\frac{1}{2}}$

$m = \pm(45^2)^{\frac{1}{2}}(p^4)^{\frac{1}{2}}(q^6)^{\frac{1}{2}}(r^{10})^{\frac{1}{2}}$

$m = \pm 45p^2q^3r^5$

(iv) $x^2 - y^2, \frac{x-y}{x+y}$

Solution:

Let m be the mean proportional, then

$x^2 - y^2 : m :: m : \frac{x-y}{x+y}$

∴ Product of means = Product of extremes

$m \times m = x^2 - y^2 \times \frac{x-y}{x+y}$

$m^2 = (x-y)(x+y) \times \frac{x-y}{x+y}$

$m^2 = (x-y)(x-y)$

$m^2 = (x-y)^2$

$m = \pm\sqrt{(x-y)^2}$

$m = \pm(x-y)$

4. Find the values of the letter involved in the following continued proportions.

(i) 5, p, 45

Solution:

Since 5, p and 45 are in continued proportions.

$5 : p :: p : 45$

∴ Product of means = Product of extremes

$p \times p = 5 \times 45$

$p^2 = 225$

$p = \pm\sqrt{225}$

$p = \pm 15$

(ii) 8, x, 18

Solution:

Since 8, x and 18 are in continued proportions.

$8 : x :: x : 18$

∴ Product of means = Product of extremes

$x \times x = 8 \times 18$

$x^2 = 144$

$x = \pm\sqrt{144}$

$x = \pm 12$

(iii) 12, 3p-6, 27

Solution:

Since 12, $3p - 6$ and 27 are in continued proportions.

$12 : 3p - 6 :: 3p - 6 : 27$

∴ Product of means = Product of extremes

$(3p-6)(3p-6) = 12 \times 27$

$(3p-6)^2 = 324$

$\sqrt{(3p-6)^2} = \pm\sqrt{324}$

$3p - 6 = \pm 18$

$3p - 6 = -18$ or $3p - 6 - 18$

$3p = 6 - 18$ or $3p = 18 + 6$

$3p = -12$ or $3p = 24$

$p = -4$ or $p = 8$

(iv) 7, m-3, 28

Solution:

Since 7, m − 3 and 28 are in continued proportions.

$7 : m - 3 :: m - 3 : 28$

∴ Product of means = Product of extremes

$(m-3)(m-3) = 7 \times 28$

$(m-3)^2 = 196$

$\sqrt{(m-3)^2} = \pm\sqrt{196}$

$m - 3 = \pm 14$

$m - 3 = -14$ or $m - 3 = 14$

$m = 3 - 14$ or $m = 3 + 14$

$m = -11$ or $m = 17$

Mathematics

Properties on Proportions
If four quantities a, b, c and d form a proportion, then cross other varied proportions may be derived by the properties of fractions.

(1) Theorem of Componendo
If $\frac{a}{b} = \frac{c}{d}$, then $\frac{a+b}{b} = \frac{c+d}{d}$

(2) Theorem of Dividendo
If $\frac{a}{b} = \frac{c}{d}$, then $\frac{a-b}{b} = \frac{c-d}{d}$

(3) Theorem of Componendo-Dividendo
If $\frac{a}{b} = \frac{c}{d}$, then $\frac{a+b}{a-b} = \frac{c+d}{c-d}$

(4) Theorem of Invertendo
If $\frac{a}{b} = \frac{c}{d}$, then $\frac{b}{a} = \frac{d}{c}$

(5) Theorem of Alternendo
If $\frac{a}{b} = \frac{c}{d}$, then $\frac{a}{c} = \frac{b}{d}$

(6) Theorem of Componendo-Alternendo
If $\frac{a}{b} = \frac{c}{d}$, then $\frac{a+c}{b+d} = \frac{a-c}{b-d}$

Exercise 3.4

Mathematics

1. Find x, if (i) $\frac{x}{3} = \frac{4}{5}$ (ii) $\frac{x}{3} = \frac{5}{4}$

Solution

Given $\frac{x}{3} = \frac{4}{5}$

Multiplying both sides by 3

$\frac{x}{3} \times 3 = \frac{4}{5} \times 3$

$x = \frac{4}{5} \times 3$

$x = \frac{12}{5}$

Hence proved

Given $\frac{x}{3} = \frac{5}{4}$

Multiplying both sides by 3

$\frac{x}{3} \times 3 = \frac{5}{4} \times 3$

$x = \frac{5}{4} \times 3$

$x = \frac{15}{4}$

Hence proved

2. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{x+y}{x-y}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x+y}{x-y} = \frac{3+4}{3-4}$

$\frac{x+y}{x-y} = \frac{7}{-1}$

$\frac{x+y}{x-y} = -7$

Hence proved

3. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{x-y}{x+y}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a-b}{a+b} = \frac{c-d}{c+d}$

$\frac{x-y}{x+y} = \frac{5-6}{5+6}$

$\frac{x-y}{x+y} = \frac{-1}{11}$

Hence proved

4. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{3x+5y}{3x-5y}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

Multiplying both sides by $\frac{5}{3}$, we get

$\frac{x}{y} \times \frac{5}{3} = \frac{3}{4} \times \frac{5}{3}$

$\frac{5x}{3y} = \frac{5}{4}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{5x+3y}{5x-3y} = \frac{5+4}{5-4}$

$\frac{5x+3y}{5x-3y} = \frac{9}{1}$

$\frac{5x+3y}{5x-3y} = 9$

Hence proved

5. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{3x-2y}{3x+2y}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

Multiplying both sides by $\frac{3}{2}$, we get

$\frac{x}{y} \times \frac{3}{2} = \frac{5}{6} \times \frac{3}{2}$

$\frac{3x}{2y} = \frac{5}{4}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a-b}{a+b} = \frac{c-d}{c+d}$

$\frac{3x-2y}{3x+2y} = \frac{5-4}{5+4}$

$\frac{3x-2y}{3x+2y} = \frac{1}{9}$

Hence proved

6. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{x^2+y^2}{x^2-y^2}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

Squaring both sides, we get

$\left(\frac{x}{y}\right)^2 = \left(\frac{3}{4}\right)^2$

$\frac{x^2}{y^2} = \frac{9}{16}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^2+y^2}{x^2-y^2} = \frac{9+16}{9-16}$

$\frac{x^2+y^2}{x^2-y^2} = \frac{25}{-7}$

$\frac{x^2+y^2}{x^2-y^2} = -\frac{25}{7}$

Hence proved

7. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{x^2+y^2}{x^2-y^2}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

Squaring both sides, we get

$\left(\frac{x}{y}\right)^2 = \left(\frac{5}{6}\right)^2$

$\frac{x^2}{y^2} = \frac{25}{36}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^2+y^2}{x^2-y^2} = \frac{25+36}{25-36}$

$\frac{x^2+y^2}{x^2-y^2} = \frac{61}{-11}$

$\frac{x^2+y^2}{x^2-y^2} = -\frac{61}{11}$

Hence proved

8. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{x^3+y^3}{x^3-y^3}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

Cubing both sides, we get

$\left(\frac{x}{y}\right)^3 = \left(\frac{3}{4}\right)^3$

$\frac{x^3}{y^3} = \frac{27}{64}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{27+64}{27-64}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{91}{-37}$

$\frac{x^3+y^3}{x^3-y^3} = -\frac{91}{37}$

Hence proved

9. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{x^3+y^3}{x^3-y^3}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

Cubing both sides, we get

$\left(\frac{x}{y}\right)^3 = \left(\frac{5}{6}\right)^3$

$\frac{x^3}{y^3} = \frac{125}{216}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{125+216}{125-216}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{341}{-91}$

$\frac{x^3+y^3}{x^3-y^3} = -\frac{341}{91}$

Hence proved

10. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{x^4+y^4}{x^4-y^4}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

Squaring both sides, we get

$\left(\frac{x}{y}\right)^2 = \left(\frac{3}{4}\right)^2$

$\frac{x^2}{y^2} = \frac{9}{16}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^2+y^2}{x^2-y^2} = \frac{9+16}{9-16}$

$\frac{x^2+y^2}{x^2-y^2} = \frac{25}{-7}$

$\frac{x^2+y^2}{x^2-y^2} = -\frac{25}{7}$

Squaring both sides, we get

$\left(\frac{x^2+y^2}{x^2-y^2}\right)^2 = \left(-\frac{25}{7}\right)^2$

$\frac{x^4+y^4+2x^2y^2}{x^4+y^4-2x^2y^2} = \frac{625}{49}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^4+y^4+2x^2y^2+x^4+y^4-2x^2y^2}{x^4+y^4+2x^2y^2-x^4-y^4+2x^2y^2} = \frac{625+49}{625-49}$

$\frac{2x^4+2y^4}{4x^2y^2} = \frac{674}{576}$

$\frac{x^4+y^4}{2x^2y^2} = \frac{337}{288}$

$\frac{x^4+y^4}{x^4-y^4} = \frac{337}{49}$

Hence proved

11. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{x^4+y^4}{x^4-y^4}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

Squaring both sides, we get

$\left(\frac{x}{y}\right)^2 = \left(\frac{5}{6}\right)^2$

$\frac{x^2}{y^2} = \frac{25}{36}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^2+y^2}{x^2-y^2} = \frac{25+36}{25-36}$

$\frac{x^2+y^2}{x^2-y^2} = \frac{61}{-11}$

$\frac{x^2+y^2}{x^2-y^2} = -\frac{61}{11}$

Squaring both sides, we get

$\left(\frac{x^2+y^2}{x^2-y^2}\right)^2 = \left(-\frac{61}{11}\right)^2$

$\frac{x^4+y^4+2x^2y^2}{x^4+y^4-2x^2y^2} = \frac{3721}{121}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^4+y^4+2x^2y^2+x^4+y^4-2x^2y^2}{x^4+y^4+2x^2y^2-x^4-y^4+2x^2y^2} = \frac{3721+121}{3721-121}$

$\frac{2x^4+2y^4}{4x^2y^2} = \frac{3842}{3600}$

$\frac{x^4+y^4}{2x^2y^2} = \frac{1921}{1800}$

$\frac{x^4+y^4}{x^4-y^4} = \frac{1921}{121}$

Hence proved

12. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{x^5+y^5}{x^5-y^5}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

Cubing both sides, we get

$\left(\frac{x}{y}\right)^3 = \left(\frac{3}{4}\right)^3$

$\frac{x^3}{y^3} = \frac{27}{64}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{27+64}{27-64}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{91}{-37}$

$\frac{x^3+y^3}{x^3-y^3} = -\frac{91}{37}$

Squaring both sides, we get

$\left(\frac{x^3+y^3}{x^3-y^3}\right)^2 = \left(-\frac{91}{37}\right)^2$

$\frac{x^6+y^6+2x^3y^3}{x^6+y^6-2x^3y^3} = \frac{8281}{1369}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^6+y^6+2x^3y^3+x^6+y^6-2x^3y^3}{x^6+y^6+2x^3y^3-x^6-y^6+2x^3y^3} = \frac{8281+1369}{8281-1369}$

$\frac{2x^6+2y^6}{4x^3y^3} = \frac{9650}{6912}$

$\frac{x^6+y^6}{2x^3y^3} = \frac{4825}{3456}$

$\frac{x^6+y^6}{x^6-y^6} = \frac{4825}{1569}$

Hence proved

13. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{x^5+y^5}{x^5-y^5}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

Cubing both sides, we get

$\left(\frac{x}{y}\right)^3 = \left(\frac{5}{6}\right)^3$

$\frac{x^3}{y^3} = \frac{125}{216}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{125+216}{125-216}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{341}{-91}$

$\frac{x^3+y^3}{x^3-y^3} = -\frac{341}{91}$

Squaring both sides, we get

$\left(\frac{x^3+y^3}{x^3-y^3}\right)^2 = \left(-\frac{341}{91}\right)^2$

$\frac{x^6+y^6+2x^3y^3}{x^6+y^6-2x^3y^3} = \frac{116281}{8281}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^6+y^6+2x^3y^3+x^6+y^6-2x^3y^3}{x^6+y^6+2x^3y^3-x^6-y^6+2x^3y^3} = \frac{116281+8281}{116281-8281}$

$\frac{2x^6+2y^6}{4x^3y^3} = \frac{124562}{108000}$

$\frac{x^6+y^6}{2x^3y^3} = \frac{62281}{54000}$

$\frac{x^6+y^6}{x^6-y^6} = \frac{62281}{7719}$

Hence proved

14. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{x^6+y^6}{x^6-y^6}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

Cubing both sides, we get

$\left(\frac{x}{y}\right)^3 = \left(\frac{3}{4}\right)^3$

$\frac{x^3}{y^3} = \frac{27}{64}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{27+64}{27-64}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{91}{-37}$

$\frac{x^3+y^3}{x^3-y^3} = -\frac{91}{37}$

Squaring both sides, we get

$\left(\frac{x^3+y^3}{x^3-y^3}\right)^2 = \left(-\frac{91}{37}\right)^2$

$\frac{x^6+y^6+2x^3y^3}{x^6+y^6-2x^3y^3} = \frac{8281}{1369}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^6+y^6+2x^3y^3+x^6+y^6-2x^3y^3}{x^6+y^6+2x^3y^3-x^6-y^6+2x^3y^3} = \frac{8281+1369}{8281-1369}$

$\frac{2x^6+2y^6}{4x^3y^3} = \frac{9650}{6912}$

$\frac{x^6+y^6}{2x^3y^3} = \frac{4825}{3456}$

$\frac{x^6+y^6}{x^6-y^6} = \frac{4825}{1569}$

Hence proved

15. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{x^6+y^6}{x^6-y^6}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

Cubing both sides, we get

$\left(\frac{x}{y}\right)^3 = \left(\frac{5}{6}\right)^3$

$\frac{x^3}{y^3} = \frac{125}{216}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{125+216}{125-216}$

$\frac{x^3+y^3}{x^3-y^3} = \frac{341}{-91}$

$\frac{x^3+y^3}{x^3-y^3} = -\frac{341}{91}$

Squaring both sides, we get

$\left(\frac{x^3+y^3}{x^3-y^3}\right)^2 = \left(-\frac{341}{91}\right)^2$

$\frac{x^6+y^6+2x^3y^3}{x^6+y^6-2x^3y^3} = \frac{116281}{8281}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^6+y^6+2x^3y^3+x^6+y^6-2x^3y^3}{x^6+y^6+2x^3y^3-x^6-y^6+2x^3y^3} = \frac{116281+8281}{116281-8281}$

$\frac{2x^6+2y^6}{4x^3y^3} = \frac{124562}{108000}$

$\frac{x^6+y^6}{2x^3y^3} = \frac{62281}{54000}$

$\frac{x^6+y^6}{x^6-y^6} = \frac{62281}{7719}$

Hence proved

16. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{x^7+y^7}{x^7-y^7}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

Raising both sides to the power 7, we get

$\left(\frac{x}{y}\right)^7 = \left(\frac{3}{4}\right)^7$

$\frac{x^7}{y^7} = \frac{2187}{16384}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^7+y^7}{x^7-y^7} = \frac{2187+16384}{2187-16384}$

$\frac{x^7+y^7}{x^7-y^7} = \frac{18571}{-14197}$

$\frac{x^7+y^7}{x^7-y^7} = -\frac{18571}{14197}$

Hence proved

17. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{x^7+y^7}{x^7-y^7}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

Raising both sides to the power 7, we get

$\left(\frac{x}{y}\right)^7 = \left(\frac{5}{6}\right)^7$

$\frac{x^7}{y^7} = \frac{78125}{279936}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^7+y^7}{x^7-y^7} = \frac{78125+279936}{78125-279936}$

$\frac{x^7+y^7}{x^7-y^7} = \frac{358061}{-201811}$

$\frac{x^7+y^7}{x^7-y^7} = -\frac{358061}{201811}$

Hence proved

18. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{x^8+y^8}{x^8-y^8}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

Raising both sides to the power 8, we get

$\left(\frac{x}{y}\right)^8 = \left(\frac{3}{4}\right)^8$

$\frac{x^8}{y^8} = \frac{6561}{65536}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^8+y^8}{x^8-y^8} = \frac{6561+65536}{6561-65536}$

$\frac{x^8+y^8}{x^8-y^8} = \frac{72097}{-58975}$

$\frac{x^8+y^8}{x^8-y^8} = -\frac{72097}{58975}$

Hence proved

19. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{x^8+y^8}{x^8-y^8}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

Raising both sides to the power 8, we get

$\left(\frac{x}{y}\right)^8 = \left(\frac{5}{6}\right)^8$

$\frac{x^8}{y^8} = \frac{390625}{1679616}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^8+y^8}{x^8-y^8} = \frac{390625+1679616}{390625-1679616}$

$\frac{x^8+y^8}{x^8-y^8} = \frac{2070241}{-1289991}$

$\frac{x^8+y^8}{x^8-y^8} = -\frac{2070241}{1289991}$

Hence proved

20. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{x^9+y^9}{x^9-y^9}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

Raising both sides to the power 9, we get

$\left(\frac{x}{y}\right)^9 = \left(\frac{3}{4}\right)^9$

$\frac{x^9}{y^9} = \frac{19683}{262144}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^9+y^9}{x^9-y^9} = \frac{19683+262144}{19683-262144}$

$\frac{x^9+y^9}{x^9-y^9} = \frac{281827}{-242461}$

$\frac{x^9+y^9}{x^9-y^9} = -\frac{281827}{242461}$

Hence proved

21. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{x^9+y^9}{x^9-y^9}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

Raising both sides to the power 9, we get

$\left(\frac{x}{y}\right)^9 = \left(\frac{5}{6}\right)^9$

$\frac{x^9}{y^9} = \frac{1953125}{10077696}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^9+y^9}{x^9-y^9} = \frac{1953125+10077696}{1953125-10077696}$

$\frac{x^9+y^9}{x^9-y^9} = \frac{12030821}{-8124571}$

$\frac{x^9+y^9}{x^9-y^9} = -\frac{12030821}{8124571}$

Hence proved

22. If $\frac{x}{y} = \frac{3}{4}$, find the value of $\frac{x^{10}+y^{10}}{x^{10}-y^{10}}$

Solution

Given $\frac{x}{y} = \frac{3}{4}$

Raising both sides to the power 10, we get

$\left(\frac{x}{y}\right)^{10} = \left(\frac{3}{4}\right)^{10}$

$\frac{x^{10}}{y^{10}} = \frac{59049}{1048576}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^{10}+y^{10}}{x^{10}-y^{10}} = \frac{59049+1048576}{59049-1048576}$

$\frac{x^{10}+y^{10}}{x^{10}-y^{10}} = \frac{1107625}{-989527}$

$\frac{x^{10}+y^{10}}{x^{10}-y^{10}} = -\frac{1107625}{989527}$

Hence proved

23. If $\frac{x}{y} = \frac{5}{6}$, find the value of $\frac{x^{10}+y^{10}}{x^{10}-y^{10}}$

Solution

Given $\frac{x}{y} = \frac{5}{6}$

Raising both sides to the power 10, we get

$\left(\frac{x}{y}\right)^{10} = \left(\frac{5}{6}\right)^{10}$

$\frac{x^{10}}{y^{10}} = \frac{9765625}{60466176}$

By componendo-dividendo theorem, if $\frac{a}{b} = \frac{c}{d}$

$\frac{a+b}{a-b} = \frac{c+d}{c-d}$

$\frac{x^{10}+y^{10}}{x^{10}-y^{10}} = \frac{9765625+60466176}{9765625-60466176}$

$\frac{x^{10}+y^{10}}{x^{10}-y^{10}} = \frac{70231801}{-50700551}$

$\frac{x^{10}+y^{10}}{x^{10}-y^{10}} = -\frac{70231801}{50700551}$

Hence proved

(i) Joint variation

A combination of direct and inverse variations of one or more than one variable forms joint variation.

If a variable y varies directly as x and varies inversely as z.

Then $y \propto x$ and $y \propto \frac{1}{z}$

In joint variation, we write it as

$$y \propto \frac{x}{z}$$

i.e, $y = k\frac{x}{z}$

Where k ≠ 0 is the constant of variation.

Exercise 3.5

1. If s varies directly as u^2 and inversely as v and t = 7 when u = 3, v = 2. Find the value of s when u = 4 and v = 10.

Solution:

Given that s varies directly as u^2, so

$s \propto u^2$

Also given that S varies inversely as v,so

$s \propto \frac{1}{v}$

In joint variation, we can write

$s \propto \frac{u^2}{v}$

$\Rightarrow \quad s = k\frac{u^2}{v}$ ……(i)

Put s = 7, u = 3 and v = 2 in eq (i), we get

$7 = k\frac{(3)^2}{2}$

$7 = \frac{9k}{2}$

$or \ \frac{9k}{2} = 7$

Multiplying both sides by $\frac{2}{9}$, we get

$k = 7 \times \frac{2}{9}$

$k = \frac{14}{9}$

Put $k = \frac{14}{9}$ in eq (i), we get

$s = \frac{14u^2}{9v}$ ……(ii)

Put u = 4 and v = 10 in eq (ii), we get

$S = \frac{14 \times (4)^2}{9(10)}$

$= \frac{14 \times 16}{9 \times 10} = \frac{504}{90} = \frac{28}{5}$

2. If w varies jointly as x, y^2 and 2 and w = 5 when x == 2, y = 3, z = 10. Find w When x = 4, y = 7 and z = 3.

Solution:

Given that s varies directly as x, y^2 and z.

Therefore $w \propto xy^2z$

$\Rightarrow \quad w = kxy^2z$ ……(i)

Put w =5, x = 2 and y = 3 and z = 10 in eq (i), we get

$5 = k(2)(3)^2(10)$

$5 = k(2)(9)(10)$

$5 = 180k$

$180k = 5$

$k = \frac{5}{180} = \frac{1}{36}$

Put $k = \frac{1}{36}$ in eq (i), we get

$w = \frac{1}{36}xy^2z$ ……(ii)

Put x = 4 and y = 7 and z = 3 in eq (ii), we get

$w = \frac{1}{36}(4)(7^2)(3)$

$= \frac{1}{36}(4)(49)(3)$

$= \frac{588}{36} = \frac{49}{3}$

3. If y varies directly as x^2 and inversely as z^2 and t, and y = 16 when x = 4, z = 2, t = 3. Find the value of y when x = 2, z = 3 and t = 4.

Solution:

Given that s varies directly as x^2.

Therefore $y \propto x^2$

Also given that y varies inversely as z^2 and t.

Therefore $y \propto \frac{1}{z^2t}$

In joint variation, we can write

$y \propto \frac{x^2}{z^2t}$

$\Rightarrow \quad y = k\frac{x^2}{z^2t}$ ……(i)

Put y = 16, x = 4 and z = 2 and t = 3 in eq (i), we get

$16 = k\frac{(4)^2}{(2)^2(3)}$

$16 = k\frac{16}{4 \times 3}$

$16 = k\frac{16}{12}$

$or \ \frac{16}{12}k = 16$

$k = 16 \times \frac{12}{16}$

$k = 3$

Put k = z in eq (i), we get

$y = \frac{3x^2}{z^2t}$ ……(ii)

Put x = 2 and z = 3 and t = 4 in eq (ii), we get

$y = \frac{3(2)^2}{(3)^2(4)}$

$= \frac{3 \times 8}{9 \times 4} = \frac{2}{3}$

4. If u varies directly as x^2 and inversely as the product ys^3, and u = 2 when x = 8, y = 7, z = 2. Find the value of u when x = 6, y = 2, z = 2.

Solution:

Given that u varies directly as x^2.

Therefore $u \propto x^2$

Also given that u varies inversely as yz^3.

Therefore $u \propto \frac{1}{yz^3}$

In joint variation, we can write

$u \propto \frac{x^2}{yz^3}$

$\Rightarrow \quad u = k\frac{x^2}{yz^3}$ ……(i)

Put u = 2, x = 8 and y = 7 and z = 2, we get

$2 = k\frac{(8)^2}{(7)(2)^3}$

$2 = \frac{64}{56}k$

$\frac{64}{56}k = 2$

$k = 2 \times \frac{56}{64} = \frac{7}{4}$

Put $k = \frac{7}{4}$ in eq (i), we get

$u = \frac{7x^2}{4yz^3}$ ……(ii)

Put x = 6 and y = 3 and z = 2 in eq (ii), we get

$u = \frac{7(6)^2}{4(3)(2)^3}$

$= \frac{7 \times 36}{4 \times 3 \times 8} = \frac{252}{96} = \frac{21}{8}$

5. If v varies directly as the product xy^2 and inversely as z^2 and v = 27 when x = 7, y = 6, z = 7. Find the value of v when x = 4, y = 2, z = 3.

Solution:

Given that v varies directly as xy^2.

Therefore $v \propto xy^2$

Also given that u varies inversely as z^2.

Therefore $v \propto \frac{1}{z^2}$

In joint variation, we can write

$v \propto \frac{xy^2}{z^2}$

$\Rightarrow \quad v = k\frac{xy^2}{z^2}$ ……(i)

Put v = 27, x = 7 and y = 6 and z = 7, we get

$27 = \frac{k(7)(6)^2}{(7)^2}$

$27 = \frac{k(7)(216)}{49}$

$27 = \frac{1512k}{49}$

$or \ k = 27 \times \frac{49}{1512}$

$k = \frac{1323}{1512} = \frac{7}{8}$

Put $k = \frac{7}{8}$ in eq (i), we get

$v = \frac{7xy^2}{8z^2}$ ……(ii)

Put x = 4 and y = 2 and z = 3 in eq (ii), we get

$v = \frac{7(4)(2)^2}{8(3)^2}$

$= \frac{336}{72} = \frac{14}{3}$

6. If w varies inversely as the cube of u, and w = 5 when u = 3. Find w, when u = 4.

Solution:

Given that w varies inversely as u^3.

Therefore $w \propto \frac{1}{u^3}$

$\Rightarrow \quad w = \frac{k}{u^3}$ ……(i)

Put w =5,u = 3 in eq (i), we get

$5 = \frac{k}{(3)^3}$

$k = 27 \times 5 = 135$

Put $k = 135$ in eq (i), we get

$w = \frac{135}{u^3}$ ……(ii)

Put u = 6 in eq (ii), we get

$w = \frac{135}{(6)^3}$

$= \frac{135}{216} = \frac{5}{8}$

6. Method:
(1) Use k – method to prove conditional equalities involving proportions.
If a / b = c / d is a proportion, then putting each ratio equal to k

$$\frac{a}{b} = \frac{c}{d} = k$$
$$\frac{a}{b} = k \text{ and } \frac{c}{d} = k$$
$$a = bk \text{ and } c = dk$$

Using the above equations, we can solve certain problems relating to proportions more easily. This method's known as a method.

Exercise 3.6

1. If a, b ≠ c , d, (a, b, c, d ≠ 0), then show that

(i) $\frac{4a-7b}{4a+5b} = \frac{4c-7d}{4c+5d}$

Solution:

Given $a, b \neq c, d$

Let $\frac{a}{b} = \frac{c}{d} = k$

$\Rightarrow \frac{a}{b} = k$ and $\frac{c}{d} = k$

$a = bk$ and $c = dk$ ……(i)

L.H.S. $= \frac{4a-7b}{4a+5b} = \frac{4bk-7b}{4bk+5b}$

$= \frac{b(4k-7)}{b(4k+5)}$

$= \frac{4k-7}{4k+5}$ ……(ii)

R.H.S. $= \frac{4c-7d}{4c+5d} = \frac{4dk-7d}{4dk+5d}$

$= \frac{d(4k-7)}{d(4k+5)}$

$= \frac{4k-7}{4k+5}$ ……(iii)

From (ii) and (iii) we have

L.H.S. = R.H.S

Hence $\frac{4a-7b}{4a+5b} = \frac{4c-7d}{4c+5d}$

(ii) $\frac{6a-5b}{6a+5b} = \frac{6c-5d}{6c+5d}$

Solution:

Given $a, b \neq c, d$

Let $\frac{a}{b} = \frac{c}{d} = k$

$\Rightarrow \frac{a}{b} = k$ and $\frac{c}{d} = k$

$a = bk$ and $c = dk$ ……(i)

L.H.S. $= \frac{6a-5b}{6a+5b} = \frac{6bk-5b}{6bk+5b}$

$= \frac{b(6k-5)}{b(6k+5)}$

$= \frac{6k-5}{6k+5}$ ……(ii)

R.H.S. $= \frac{6c-5d}{6c+5d} = \frac{6dk-5d}{6dk+5d}$

$= \frac{d(6k-5)}{d(6k+5)}$

$= \frac{6k-5}{6k+5}$ ……(iii)

From (ii) and (iii) we have

L.H.S. = R.H.S

Hence $\frac{6a-5b}{6a+5b} = \frac{6c-5d}{6c+5d}$

(iii) $\frac{a^2}{b^2} = \frac{\sqrt{a^2+c^2}}{\sqrt{b^2+d^2}}$

Solution:

Given $a, b \neq c, d$

Let $\frac{a}{b} = \frac{c}{d} = k$

$\Rightarrow \frac{a}{b} = k$ and $\frac{c}{d} = k$

$a = bk$ and $c = dk$ ……(i)

L.H.S. $= \frac{a^2}{b^2} = \frac{(bk)^2}{b^2}$ R.H.S $= \frac{\sqrt{a^2+c^2}}{\sqrt{b^2+d^2}} = \frac{\sqrt{b^2k^2+d^2k^2}}{\sqrt{b^2+d^2}}$

$= \frac{b^2k^2}{b^2}$ $= \frac{\sqrt{k^2(b^2+d^2)}}{\sqrt{b^2+d^2}}$

$= k$ ……(ii) $= \frac{k\sqrt{b^2+d^2}}{\sqrt{b^2+d^2}}$

$= k$ ……(iii)

From (ii) and (iii) we have

L.H.S. = R.H.S

Hence $\frac{a^2}{b^2} = \frac{\sqrt{a^2+c^2}}{\sqrt{b^2+d^2}}$

(iv) $|\mathbf{a}^2 + \mathbf{c}^2| \cdot |\mathbf{b}^2 + \mathbf{d}^2| = |\mathbf{a}\mathbf{b} + \mathbf{c}\mathbf{d}|^2 + |\mathbf{b}\mathbf{c} - \mathbf{a}\mathbf{d}|^2$

Solution:

Given $a, b \neq c, d$

Let $\frac{a}{b} = \frac{c}{d} = k$

$\Rightarrow \frac{a}{b} = k$ and $\frac{c}{d} = k$

$a = bk$ and $c = dk$

L.H.S. $= |a^2 + c^2| \cdot |b^2 + d^2| = |a^2b^2 + b^2d^2 + a^2c^2 + b^2d^2|$

$= |a^2b^2 + a^2c^2 + b^2d^2 + b^2d^2|$

$= |a^2(b^2 + c^2) + b^2(c^2 + d^2)|$

$= |a^2(b^2 + c^2) + b^2(c^2 + d^2)|$

$= a^2(b^2 + c^2) + b^2(c^2 + d^2)$ ……(i)

R.H.S. $= |ab + cd|^2 + |bc - ad|^2$

$= |ab + cd|^2 + |bc - ad|^2$

$= a^2b^2 + c^2d^2 + 2abcd + b^2c^2 + a^2d^2 - 2abcd$

$= a^2b^2 + c^2d^2 + b^2c^2 + a^2d^2$

$= a^2(b^2 + c^2) + d^2(c^2 + b^2)$

$= a^2(b^2 + c^2) + b^2(c^2 + d^2)$ ……(ii)

From (i) and (ii) we have

L.H.S. = R.H.S

Hence $|a^2 + c^2| \cdot |b^2 + d^2| = |ab + cd|^2 + |bc - ad|^2$

$$\frac{b^2a^2 + a^2c^2}{b^2 + c^2} = \frac{(b^2c^2)(a^2)}{b^2c^2}$$
$$\frac{b^2(a^2 + c^2)}{b^2 + c^2} = \frac{(b^2d^2)(a^2)}{b^2d^2}$$
$$= b^2 \quad \ldots(i) \qquad = b^2 \quad \ldots(ii)$$

From (i) and (ii) we have

L.H.S. = R.H.S

Hence $\frac{a^2 + c^2}{b^2 + c^2} = \frac{a^2}{b^2}$

(v) $\mathbf{p}(\mathbf{x}+\mathbf{b}) + \mathbf{q}\mathbf{b} \cdot \mathbf{p}(\mathbf{c}+\mathbf{d}) + \mathbf{q}\mathbf{d} = \mathbf{a} : \mathbf{c}$

Solution:

Given $a, b \neq c, d$

Let $\frac{a}{b} = \frac{c}{d} = k$

$\Rightarrow \frac{a}{b} = k$ and $\frac{c}{d} = k$

$a = bk$ and $c = dk$

L.H.S. $= p(a+b) + qb \cdot p(c+d) + qd = a : c$

$= \frac{p(a+b)+qb}{p(c+d)+qd}$

$= \frac{p(bk+b)+qb}{p(dk+d)+qd}$

$= \frac{pb(k+1)+qb}{pd(k+1)+qd}$

$= \frac{b[p(k+1)+q]}{d[p(k+1)+q]}$

$= \frac{b}{d}$ ……(i)

From (i) and (ii) we have

L.H.S. = R.H.S

Hence $p(a+b) + qb \cdot p(c+d) + qd = a : c$

(vi) $\frac{a^2+b^2}{ab} : \frac{c^2+d^2}{cd} = \frac{a^2-b^2}{a+b} : \frac{c^2-d^2}{c+d}$

Solution:

Given $a, b \neq c, d$

Let $\frac{a}{b} = \frac{c}{d} = k$

$\Rightarrow \frac{a}{b} = k$ and $\frac{c}{d} = k$

$a = bk$ and $c = dk$

L.H.S. $= a^2 + b^2 : \frac{a^2}{a+b}$ R.H.S $= c^2 + d^2 : \frac{c^2}{c+d}$

$= a^2 + b^2 : \frac{a^2}{a+b}$ $= c^2 + d^2 : \frac{c^2}{c+d}$

$= a^2 + b^2 : \frac{a^2}{a+b}$ $= c^2 + d^2 : \frac{c^2}{c+d}$

$= (a^2 + b^2) : \frac{(a+b)a}{a}$ $= (c^2 + d^2) : \frac{(c+d)c}{c}$

$= (a^2 + b^2) : \frac{b(a+b)}{a}$ $= (c^2 + d^2) : \frac{d(c+d)}{c}$

$= (a^2 + b^2) : \frac{b(bk+b)}{a}$ $= (c^2 + d^2) : \frac{d(dk+d)}{c}$

$= (a^2 + b^2) : \frac{b^2(k+1)}{a}$ $= (c^2 + d^2) : \frac{d^2(k+1)}{c}$

$= (a^2 + b^2) : \frac{b^2(k+1)}{bk}$ $= (c^2 + d^2) : \frac{d^2(k+1)}{dk}$

$= (a^2 + b^2) : \frac{b(k+1)}{k}$ $= (c^2 + d^2) : \frac{d(k+1)}{k}$ ……(ii)

From (i) and (ii) we have

L.H.S. = R.H.S

Hence $a^2 + b^2 : \frac{a^2}{a+b} = c^2 + d^2 : \frac{c^2}{c+d}$

(vii) $\frac{a}{a-b} : \frac{a+b}{b} = \frac{c}{c-d} : \frac{c+d}{d}$

Solution:

Given $a, b \neq c, d$

Let $\frac{a}{b} = \frac{c}{d} = k$

$\Rightarrow \frac{a}{b} = k$ and $\frac{c}{d} = k$

$a = bk$ and $c = dk$

L.H.S. $= \frac{a}{a-b} : \frac{a+b}{b}$ R.H.S $= \frac{c}{c-d} : \frac{c+d}{d}$

$= \frac{bk}{bk-b} : \frac{bk+b}{b}$ $= \frac{dk}{dk-d} : \frac{dk+d}{d}$

$= \frac{bk}{b(k-1)} : \frac{b(k+1)}{b}$ $= \frac{dk}{d(k-1)} : \frac{d(k+1)}{d}$

$= \frac{k}{k-1} : \frac{k+1}{1}$ $= \frac{k}{k-1} : \frac{k+1}{1}$

$= \frac{k}{k-1} : (k+1)$ ……(i) $= \frac{k}{k-1} : (k+1)$ ……(ii)

From (i) and (ii) we have

L.H.S. = R.H.S

Hence $\frac{a}{a-b} : \frac{a+b}{b} = \frac{c}{c-d} : \frac{c+d}{d}$

2. If $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} (a, b, c, d, e, f \neq 0)$ **then show that**

(i) $\frac{3a}{2b} = \frac{\sqrt{3a^2+5c^2+7e^2}}{\sqrt{2b^2+5d^2+7f^2}}$

Solution:

Let $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = k$

$\Rightarrow \frac{a}{b} = k$ and $\frac{c}{d} = k$ and $\frac{e}{f} = k$

$a = bk$ $c = dk$ $e = fk$

L.H.S. $= \frac{3a}{2b}$

$= \frac{3bk}{2b}$ ……(i)

R.H.S. $= \frac{\sqrt{3a^2+5c^2+7e^2}}{\sqrt{2b^2+5d^2+7f^2}}$

$= \frac{\sqrt{3b^2k^2+5d^2k^2+7f^2k^2}}{\sqrt{2b^2+5d^2+7f^2}}$

$= \frac{\sqrt{k^2(3b^2+5d^2+7f^2)}}{\sqrt{2b^2+5d^2+7f^2}}$

$= \frac{k\sqrt{3b^2+5d^2+7f^2}}{\sqrt{2b^2+5d^2+7f^2}}$

$= \frac{3bk}{2b}$ ……(ii)

From (i) and (ii) we have

L.H.S. = R.H.S

Hence $\frac{3a}{2b} = \frac{\sqrt{3a^2+5c^2+7e^2}}{\sqrt{2b^2+5d^2+7f^2}}$

(ii) $\frac{ace}{bdf}\left(\frac{a^2}{bd} + \frac{c^2}{df} + \frac{e^2}{fb}\right)^3 = \left(\frac{a^2}{bd}\right)^3 + \left(\frac{c^2}{df}\right)^3 + \left(\frac{e^2}{fb}\right)^3$

Solution:

Let $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = k$

$\Rightarrow \frac{a}{b} = k$ and $\frac{c}{d} = k$ and $\frac{e}{f} = k$

$a = bk$ $c = dk$ $e = fk$

L.H.S. $= \frac{ace}{bdf}\left(\frac{a^2}{bd} + \frac{c^2}{df} + \frac{e^2}{fb}\right)^3$

$= \frac{(bk)(dk)(fk)}{bdf}\left(\frac{(bk)^2}{bd} + \frac{(dk)^2}{df} + \frac{(fk)^2}{fb}\right)^3$

$= \frac{bdfk^3}{bdf}\left(\frac{b^2k^2}{bd} + \frac{d^2k^2}{df} + \frac{f^2k^2}{fb}\right)^3$

$= k^3\left(\frac{bk}{d} + \frac{dk}{f} + \frac{fk}{b}\right)^3$

$= k^3\left(k\left(\frac{b}{d} + \frac{d}{f} + \frac{f}{b}\right)\right)^3$

$= k^3 \cdot k^3\left(\frac{b}{d} + \frac{d}{f} + \frac{f}{b}\right)^3$

$= k^6\left(\frac{b}{d} + \frac{d}{f} + \frac{f}{b}\right)^3$ ……(i)

R.H.S. $= \left(\frac{a^2}{bd}\right)^3 + \left(\frac{c^2}{df}\right)^3 + \left(\frac{e^2}{fb}\right)^3$

$= \left(\frac{(bk)^2}{bd}\right)^3 + \left(\frac{(dk)^2}{df}\right)^3 + \left(\frac{(fk)^2}{fb}\right)^3$

$= \left(\frac{bk^2}{d}\right)^3 + \left(\frac{dk^2}{f}\right)^3 + \left(\frac{fk^2}{b}\right)^3$

$= \left(k\left(\frac{b}{d}\right)\right)^3 + \left(k\left(\frac{d}{f}\right)\right)^3 + \left(k\left(\frac{f}{b}\right)\right)^3$

$= k^3\left(\frac{b}{d}\right)^3 + k^3\left(\frac{d}{f}\right)^3 + k^3\left(\frac{f}{b}\right)^3$

$= k^3\left(\frac{b}{d} + \frac{d}{f} + \frac{f}{b}\right)^3$ ……(ii)

From (i) and (ii) we have

L.H.S. = R.H.S

Hence $\frac{ace}{bdf}\left(\frac{a^2}{bd} + \frac{c^2}{df} + \frac{e^2}{fb}\right)^3 = \left(\frac{a^2}{bd}\right)^3 + \left(\frac{c^2}{df}\right)^3 + \left(\frac{e^2}{fb}\right)^3$

L.H.S. $= \frac{ma}{bd} + \frac{nc}{df} + \frac{pe}{fb}$

$= \frac{(bk)(dk)}{bd} + \frac{(dk)(fk)}{df} + \frac{(fk)(bk)}{fb}$

$= \frac{bdk^2}{bd} + \frac{dfk^2}{df} + \frac{fbk^2}{fb}$

$= k^2 + k^2 + k^2$

$= 3k^2$ ……(i)

R.H.S. $= \frac{ma}{mb - nd} + \frac{nc}{nd - pf} + \frac{pe}{pf - mb}$

$= \frac{mbk}{mbk - nd} + \frac{ndk}{ndk - pf} + \frac{pfk}{pfk - mbk}$

$= k^2 + k^2 + k^2$

$= 3k^2$ ……(ii)

From (i) and (ii) we have

L.H.S. = R.H.S

Hence $\frac{ma}{mb - nd} + \frac{nc}{nd - pf} + \frac{pe}{pf - mb} = k^2 + k^2 + k^2$

Exercise 3.7

1. The surface area A of a cube varies directly as the square of the length l of an edge and A = 27 square units when l = 3 units.

Find

(i) A when l = 4 units

(ii) l when A = 12 sq. units.

Solution:

Given that $A \propto l^2$

$\Rightarrow A = kl^2$ ……(1)

Put $A = 27$ and $l = 3$ in eq.(1), we get

$27 = k(3)^2$

$27 = 9k$

$9k = 27$

$k = \frac{27}{9} = 3$

Put $k = 3$ in eq.(1), we get

$A = 3l^2$ ……(2)

(i) Put $l = 4$ in eq.(2), we get

$A = 3(4)^2$

$= 3(16) = 48$ sq.units

(ii) Put $A = 12$ in eq.(2), we get

$12 = 3l^2$

or $3l^2 = 12$

$\Rightarrow l^2 = 4$

$l = 2$

2. The surface area S of the sphere varies directly as the square of radius r, and S = 14π when r = 2. Find r when S = 36π.

Solution:

Given that $S \propto r^2$

$\Rightarrow S = kr^2$ ……(i)

Put $S = 14\pi$ and $r = 2$ in eq.(i), we get

$14\pi = k(2)^2$

$14\pi = 4k$

$4k = 14\pi$

$k = 4\pi$

Put $k = 4\pi$ in eq.(i), we get

$S = 4\pi r^2$ ……(ii)

Put $S = 36\pi$ in eq.(ii), we get

$36\pi = 4\pi r^2$

or $4\pi r^2 = 36\pi$

$r^2 = \frac{36\pi}{4\pi}$

$r^2 = 9$

$r = 3$

3. In Hook's law the force F applied to stretch a spring varies directly as the amount of elongation S and F = 32 lb when s = 1.4 in.

Find

(i) S when F = 80 lb

(ii) F when S = 0.8 in.

Solution:

Given that $F \propto S$

$\Rightarrow F = kS$ ……(1)

Put $F = 32$ and $S = 1.4$ in eq.(1), we get

$32 = k(1.4)$

or $1.4k = 32$

$k = \frac{32}{1.4} = 20$

Put $k = 20$ in eq.(1), we get

$F = 20S$ ……(2)

(i) Put $F = 50$ in eq.(2), we get

$50 = 20S$

$\Rightarrow S = 2.5$ in

(ii) Put $S = 0.8$ in eq.(2), we get

$F = 20(0.8) = 16$ lb

4. The intensity I of light from a given source varies inversely as the square of the distance d from it. If the intensity is 20 candlepower at a distance of 12ft. from the source, find the intensity at a point 8ft. from the source.

Solution:

Given that $I \propto \frac{1}{d^2}$

$\Rightarrow I = \frac{k}{d^2}$ ……(i)

Put $I = 20$ and $d = 12$ in eq.(i), we get

$20 = \frac{k}{(12)^2}$

or $20 = \frac{k}{144}$

$k = 20 \times 144 = 2880$

Put $k = 2880$ in eq.(i), we get

$I = \frac{2880}{d^2}$ ……(ii)

Put $d = 8$ in eq.(ii), we get

$I = \frac{2880}{(8)^2}$

$= \frac{2880}{64} = 45$

5. The pressure P in a body of fluid varies directly as the depth d. If the pressure exerted at the bottom of a tank by a column of fluid 8ft. high is 2.25 lb./sq. in, how deep must the fluid be to exert a pressure of 9 lb./sq. in?

Solution:

Given that $P \propto d$

$\Rightarrow P = kd$ ……(i)

Put $P = 2.25$ and $d = 3$ in eq.(i), we get

$2.25 = k(3)$

or $3k = 2.25$

$\Rightarrow k = \frac{2.25}{3}$

$= \frac{225}{300} = \frac{9}{20}$

Put $k = \frac{9}{20}$ in eq.(i), we get

$P = \frac{9}{20}d$ ……(ii)

Put $P = 9$ in eq.(ii), we get

$9 = \frac{9}{20}d$

or $d = \frac{9}{\frac{9}{20}} = 20$

$d = 20$ ft

6. Labor costs c varies jointly as the number of workers n and the coverage number of acres d. If the cost of 800 workers for 13 days is Rs. 286000, then find the labor cost of 600 workers for 18 days.

Solution:

Given that $c \propto nd$

$\Rightarrow c = knd$ ……(i)

Put $c = 800$ and $d = 13$ and $c = 286000$ in eq.(i), we get

$286000 = k(800)(13)$

$10400k = 286000$

$k = \frac{286000}{10400} = \frac{55}{2}$

Put $k = \frac{55}{2}$ in eq.(i), we get

$c = \frac{55}{2}nd$ ……(ii)

Put $n = 600$ and $d = 18$ in eq.(ii), we get

$c = \frac{55}{2} \times 600 \times 18$

$c = 297000$

7. The supporting load c of a pillar varies as the fourth power of its diameter d and inversely as the square of its length l. A pillar of diameter 6 inch and of height 20 feet will support a load of 63 tons. How high a 4 inch pillar must be to support a load of 28 tons?

Solution:

Given that $c \propto \frac{d^4}{l^2}$

$\Rightarrow c = k\frac{d^4}{l^2}$ ……(i)

Put $c = 6$, $l = 20$ and $c = 63$ in eq.(i), we get

$63 = k\frac{(6)^4}{(20)^2}$

$63 = k\frac{1296}{400}$

or $k = 63 \times \frac{400}{1296}$

$k = \frac{175}{4}$

Put $k = \frac{175}{4}$ in eq.(i), we get

$c = \frac{175d^4}{4l^2}$ ……(ii)

Put $d = 4$ and $c = 28$ in eq.(ii), we get

$28 = \frac{175(4)^4}{4l^2}$

$l^2 = \frac{175 \times 256}{4 \times 28} = 400$

$l = 20$ ft

8. The time T required for an elevator to lift a weight varies jointly as the height h and the lifting depth d varies and inversely as the power p of the motor. If 25 sec. are required for a 4 hp motor to lift 500 lb through 40 ft. what power is required to lift 900 lb. through 120 ft in 40 sec.?

Solution:

Given that $T \propto wd$

Also given that $T \propto \frac{1}{p}$

In joint variation, we can write

$T \propto \frac{wd}{p}$

$\Rightarrow T = k\frac{wd}{p}$ ……(i)

Put $T = 25$, $p = 4$, $w = 500$ and $d = 40$ in eq.(i), we get

$25 = k\frac{4 \times 500 \times 40}{4}$

$k = \frac{25 \times 4}{500 \times 40}$

or $k = \frac{1}{200}$

Put $k = \frac{1}{200}$ in eq.(i), we get

$T = \frac{wd}{200p}$ ……(ii)

Put $w = 800$, $d = 120$ and $T = 40$ in eq.(ii), we get

$40 = \frac{800 \times 120}{200p}$

$p = \frac{800 \times 120}{200 \times 40}$

$p = 12$hp

9. The kinetic energy (K.E.) of a body varies jointly as the mass "m" of the body and the square of its velocity "v". If the kinetic energy is 4320 ft/lb when the mass is 45 lb and the velocity is 24 ft/sec. Determine the kinetic energy of a 3000 lb automobile travelling 44 ft/sec.

Solution:

Given that $K.E. \propto mv^2$

$\Rightarrow K.E. = kmv^2$ ……(i)

Put $K.E. = 4320$, $m = 45$, $v = 24$ in eq.(i), we get

$4320 = k(45)(24)^2$

$k = \frac{4320}{45 \times 576}$

or $k = \frac{1}{6}$

Put $k = \frac{1}{6}$ in eq.(i), we get

$K.E. = \frac{1}{6}mv^2$ ……(ii)

Put $m = 3000$ and $v = 44$ in eq.(ii), we get

$K.E. = \frac{1}{6}(3000)(44)^2$

$P = 968000$

Miscellaneous Exercise 3

1. Multiple Choice Questions

Four possible answers are given for the following questions. Tick (✓) the correct answer.

(i) If a:b=3a:4, b is called

(a) relation (b) a fourth proportion

(c) a mean proportion (d) a third of three

(ii) If a:b=b:c, c is called

(a) relation (b) a fourth proportion

(c) a mean proportion (d) a third of three

(iii) In a proportion a : b = c : d, a and d are called,

(a) means (b) extremes

(c) third proportionals (d) fourth proportionals

(iv) In a proportion a : b = c : d, b and c are called

(a) means (b) extremes

(c) fourth proportionals (d) terms of four

(v) A continued proportion a : b = b : c, in "a", b is said to be _____

(a) relation (b) a fourth

(c) a means (d) a third of three

(vi) A continued proportion a : b = b : c, c is said to be _____ proportional to a and b.

(a) third (b) fourth

(c) a means (d) a third of three

(vii) Find x in proportion 4 : x :: 9 : 18

(a) $\frac{1}{2}$ (b) $\frac{2}{3}$

(c) $\frac{3}{2}$ (d) $\frac{2}{1}$

(viii) If 4 : 5 :: 7 : then

(a) $x = y^2$ (b) $x = 4y^2$

(c) $x = \frac{y}{4}$ (d) $x = y$

(ix) If $x^2 : 5 = 7 : 35$ then

(a) $x = \frac{1}{2}$ (b) $x = \pm\sqrt{7}$

(c) $x^2 = 49$ (d) $x^2 = 14^2$

(x) If $\frac{x}{y} = \frac{5}{7}$, x + 5 then

(a) $x = y$ (b) $x = \frac{y}{5}$

(c) $x = y$ (d) $x = \pm 5$

(xi) The third proportional of x^2 and x^3 is

(a) $\frac{x^5}{x^2}$ (b) x^5

(c) $\frac{x^2}{x^3}$ (d) $\frac{x^3}{x^2}$

(xii) The fourth proportional of a : c :: b : d is

(a) $\frac{ab}{c}$ (b) $\frac{c}{ab}$

(c) $\frac{a}{b}$ (d) $\frac{b}{a}$

(xiii) If a : b = c : d then alternando property is

(a) $\frac{a}{b} = \frac{c}{d}$ (b) $\frac{a}{c} = \frac{b}{d}$

(c) $\frac{a+b}{b} = \frac{c+d}{d}$ (d) $\frac{a-b}{b} = \frac{c-d}{d}$

(xiv) If a : b = c : d then componendo property is

(a) $\frac{a}{b} = \frac{c}{d}$ (b) $\frac{a}{c} = \frac{b}{d}$

(c) $\frac{a+b}{b} = \frac{c+d}{d}$ (d) $\frac{a-b}{b} = \frac{c-d}{d}$

(xv) If $\frac{a}{b} = \frac{c}{d}$ then componendo dividendo property is

(a) $\frac{a+b}{a-b} = \frac{c+d}{c-d}$ (b) $\frac{a+b}{c+d} = \frac{a-b}{c-d}$

(c) $\frac{a+b}{b} = \frac{c+d}{d}$ (d) $\frac{a-b}{b} = \frac{c-d}{d}$

Answers:

(i)	b	(ii)	b	(iii)	b	(iv)	a	(v)	b
(vi)	b	(vii)	c	(viii)	b	(ix)	d	(x)	b
(xi)	b	(xii)	b	(xiii)	b	(xiv)	c	(xv)	b

2. Write short answers of the following questions.

(i) Define ratio and give one example.

Ans: Ratio

A relation between two quantities of the same kind (measures) in some unit is called ratio. If a and b are the two quantities of the same kind and in a unit, then the ratio of a and b is written as a : b or in fraction $\frac{a}{b}$.

e.g. If a honesty team wins a games and loses b then the ratio of the games won to games lost is a : b or in fraction $\frac{a}{b}$.

(ii) Define proportion.

Ans: Proportion

A proportion is a statement, which is expressed as equivalence of two ratios.

(iii) Define direct variation.

Ans: Direct variation:

If two quantities are related in such a way that increase (decrease) in one quantity increases (decrease) in the other proportionally, then this variation is called direct variation.

(iv) Define inverse variation.

Ans: Inverse Variation:

If two quantities are related in such a way that when one quantity increases, the other decreases, then this variation is called inverse variation.

(v) State theorem of componendo dividendo.

Ans: The theorem of componendo dividendo is:

If a : b :: c : d, then

a + b : b :: c + d : c – d

(vi) Find x, if x : 3 :: 5 : 3.

Ans: x : 3 :: 5 : 3

Product of means = Product of extremes

3x = 15

$x = \frac{15}{3}$

$x = 5$

(vii) If x and y varies directly, and x = 27 when y = 9. Find the value of y when x = 9.

Solution:

Given that $x \propto y$

$x = ky$ ……(i)

Put x = 27 and y = 9 in eq (i), we get

$27 = k(9)$

$\frac{27}{9} = \frac{9k}{9}$

$3 = k$

$k = 3$

Put k = 3 in eq (i), we get

$x = 3y$ ……(ii)

Put x = 9 in eq (ii), we get

$9 = 3y$

$\frac{9}{3} = \frac{3y}{3}$

$3 = y$

$y = 3$

(viii) If y varies inversely, and y = 5, when x = 3. Find x when y = 12.

Solution:

Given that $x \propto \frac{1}{y}$

$x = \frac{k}{y}$ ……(i)

Put x = 3 and y = 5 in eq (i), we get

$3 = \frac{k}{5}$

$k = 15$

Put k = 15 in eq (i), we get

$x = \frac{15}{y}$ ……(ii)

Put x = 12 in eq (ii), we get

$12 = \frac{15}{y}$

$12y = 15$

$\frac{12y}{12} = \frac{15}{12}$

$y = \frac{5}{4}$

(ix) Find the fourth proportional to 6, 7, x.

Solution:

x, y, z to be fourth proportional, then

a : b :: c : d

Product of means = Product of terms

$7x = 6y$

$\frac{7x}{7} = \frac{6y}{7}$

$x = \frac{6y}{7}$

(x) Find a mean proportional.

Solution:

a, x, b to be mean proportional, then

a : x :: x : b

Product of means = Product of terms

$x \times x = a \times b$

$x^2 = ab$

$\sqrt{x^2} = \sqrt{ab}$

$x = \pm\sqrt{ab}$, Ans

(xi) Find a third proportional to 20 and 4.

Solution:

a, b, c to be third proportional, then

20 : 4 :: b : c

Product of means = Product of terms

$4b(c) = 20(4)$

$\frac{4b}{4} = \frac{80}{4}$

$b = \frac{80}{4}$

$b = 20$

(xii) If $x \propto \frac{1}{y^2}$ and x = 36 when x = 2. Find the f.

Solution:

Given that $x \propto \frac{1}{y^2}$

$x = \frac{k}{y^2}$ ……(i)

Put x = 2 and y = 3 in eq (i), we get

$2 = \frac{k}{3^2}$

$2 = \frac{k}{9}$

$\frac{2}{1} = \frac{k}{9}$

$k = 18$

Put k = 18 in eq (i), we get

$x = \frac{18}{y^2}$ ……(ii)

(xiii) If x ∝ xy and x = 36, when x = 2, y = 3. Find the f.

Solution:

Given that $x \propto xy$

$x = kxy$ ……(i)

Put x = 2 and y = 3 in eq (i), we get

$2 = k(3)(2)$

$2 = 6k$

$\frac{2}{6} = \frac{6k}{6}$

$\frac{1}{3} = k$

$k = \frac{1}{3}$

Put $k = \frac{1}{3}$ in eq (i), we get

$x = \frac{1}{3}xy$ ……(ii)

(xiv) If $x^2 \propto \frac{1}{y}$ and x = 36 when x = 2. Find the f.

Solution:

Given that $x^2 \propto \frac{1}{y}$

$x^2 = \frac{k}{y}$ ……(i)

Solution:

Given $x^2 = \frac{k}{y}$

$x = \frac{k}{y}$ ……(i)

Put x = 2 and y = 3 in eq (i), we get

$2 = \frac{k}{3}$

$6 = k$

$k = 6$

Put k = 6 in eq (i), we get

$x = \frac{6}{y}$

or $xy = 6$

(xv) If x ∝ y and x = 36 when y = 9. Find the f.

Solution:

Given that $x \propto y$

$x = ky$ ……(i)

Put x = 36 and y = 9 in eq (i), we get

$36 = k(9)$

$\frac{36}{9} = \frac{9k}{9}$

$4 = k$

$k = 4$

Put k = 4 in eq (i), we get

$x = 4y$

or $\frac{x}{y} = 4$

(xvi) If $x^2 \propto y$ and x = 3 when y = 9. Find the f.

Solution:

(i) If a' is mother-in-law(s) to a". Then a is _____.

(ii) If P is proportional to" Q" and R is _____.

(iii) If P is proportional to" Q" and R is _____.

(iv) The third proportional of a and b is _____.

(v) The continued proportion of a, b, c and d is _____.

Answers:

	(i)	(ii)	INTRODUCENT	(iii)	CONSEQUENT
(iv)	Sonship	(v)	means	(vi)	$p = \frac{1}{k}$
(vii)	$k = 4$	(viii)	$\frac{a}{b}$	(ix)	$\frac{a}{b} = \frac{c}{d}$
(x)	$y^2 x$	(xi)	(xii)	(xiii)	1
(xiv)	2 SHOUT	(xv)	$a^2 = ab$		

Summary

• A relation between two quantities of the same kind is called ratio.

• A proportion is a statement, which is expressed as equivalence of two ratios.

• If two ratios a:b and c:d are equal, then we can write a : b = c : d

• If two quantities are related in such a way that increase (decrease) in one quantity increases (decrease) in the other quantity, then the variation is called direct variation.

• If two quantities are related in such a way that when one quantity increases, the other decreases, then the variation is called inverse variation.

(i) Theorem of componendo

If a : b :: c : d, then (a + b) : (a + b)

(ii) Theorem of dividendo

If a : b :: c : d, then a – b :: c – d

(iii) Theorem of componendo dividendo

If a : b :: c : d, then

a + b : a – b :: c + d : c – d

and (a + b) : a – b) :: c + d : c – d

(iv) Theorem of Dividendo

If a : b :: c : d, then

(a + b) : b :: (c + d) : d

(a – b) : b :: (c – d) : d

(v) Theorem of Componendo dividendo

If a : b :: c : d, then

a + b : a – b :: c + d : c – d

• Ratio and proportion are used in solving problems of one or more than one variables (joint variation).

• To Method

(i) $\frac{a+b}{a-b} = \frac{c+d}{c-d}$

Then $\frac{a}{b} = \frac{c}{d}$ or $\frac{a}{c} = \frac{b}{d}$ or $\frac{b}{a} = \frac{d}{c}$ or $\frac{b}{d} = \frac{a}{c}$

(ii) $\frac{a^2+b^2}{ab} = \frac{c^2+d^2}{cd}$

Then $\frac{a}{b} = \frac{c}{d}$ or $\frac{a}{c} = \frac{b}{d}$ or $\frac{b}{a} = \frac{d}{c}$ or $\frac{b}{d} = \frac{a}{c}$

Here $a^2 + b^2 = (a+b)^2 - 2ab$

and $c^2 + d^2 = (c+d)^2 - 2cd$