

Exercise 8.1

Q1. Given $m\overline{AC} = 1\text{ cm}$, $m\overline{BC} = 2\text{ cm}$, $m\angle C = 120^\circ$. Compute the length AB and the area of $\triangle ABC$.

Hint: $(\overline{AB})^2 = (\overline{AC})^2 + (\overline{BC})^2 + 2m\overline{AC} \cdot m\overline{CD}$
Where $(m\overline{CD}) = (m\overline{BC})\cos(180^\circ - m\angle C)$ (Use theorem I)

Solution:

$$m\overline{AC} = 1\text{ cm}; m\overline{BC} = 2\text{ cm}; m\angle C = 120^\circ$$

$$\text{Required: } m\overline{AB} = ?$$

and Area of $\triangle ABC = ?$

$$\begin{aligned} m\overline{AB}^2 &= m\overline{AC}^2 + m\overline{BC}^2 + 2m\overline{AC} \cdot m\overline{CD} \\ &= (1)^2 + (2)^2 + 2(1)(\overline{CD}) \\ &= 1 + 4 + 2\overline{CD} \\ &= 1 + 4 + 2\overline{CD} \quad \text{--- (i)} \end{aligned}$$

In $\triangle BCD$,

$$m\angle BCD = 60^\circ$$

$$\text{and } m\angle CBD = 30^\circ$$

The side opposite to $\angle 30^\circ$ is $\overline{CD}$ which is


$\frac{1}{2}\overline{CB}$, the hypotenuse of right $\triangle CDB$.

$$\overline{CD} = 1\text{ cm}$$

By putting the value of $\overline{CD}$ in eq.(1)

$$m\overline{AB}^2 = 5 + 2(1)(1) = 5 + 2 = 7$$

$$m\overline{AB}^2 = 7 \Rightarrow m\overline{AB} = \sqrt{7}\text{ cm} = 2.646\text{ cm}$$



$$m\overline{CB}^2 = m\overline{CD}^2 + m\overline{BD}^2$$

$$2^2 = 1^2 + m\overline{BD}^2$$

$$m\overline{BD}^2 = 4 - 1 = 3$$

$$h = m\overline{BD} = \sqrt{3}$$


Area of $\triangle ABC$

$$= \frac{1}{2} \text{ base} \times \text{height}$$

$$= \frac{1}{2} m\overline{AC} \times m\overline{BD}$$

$$= \frac{1}{2} \times 1 \times \sqrt{3}$$

$$\text{Area of } \triangle ABC = \frac{\sqrt{3}}{2} \text{ sq cm}$$



Q2. Find $m\overline{AC}$ if in $\triangle ABC$ $m\overline{BC} = 6\text{ cm}$, $m\overline{AB} = 4\sqrt{2}\text{ cm}$ and $m\angle ABC = 135^\circ$.

Solution:

$$\text{Let } m\overline{BD} = x$$


In $\triangle ABD$, we have

$$\cos 45^\circ = \frac{\overline{BD}}{\overline{AB}}$$

$$\frac{1}{\sqrt{2}} = \frac{x}{4\sqrt{2}}$$

$$\sqrt{2} x = 4\sqrt{2}$$

$$x = 4\text{ cm}$$



We know that

To Prove

$$\begin{aligned} (m\overline{AC})^2 &= (m\overline{CB})^2 + (m\overline{AB})^2 + 2 \times m\overline{CB} \times m\overline{BD} \\ &= (6)^2 + (4\sqrt{2})^2 + 2 \times 6 \times 4 \end{aligned}$$

$$= 36 + 32 + 48$$

$$= 116$$

$$\Rightarrow m\overline{AC} = \sqrt{116} = \sqrt{4 \times 29} = 2\sqrt{29}\text{ cm}$$

EXERCISE 8.2


Q1. In a $\triangle ABC$ calculate $m \overline{BC}$ when $m \overline{AB} = 6\text{cm}$, $m \overline{AC} = 4\text{cm}$ and $m\angle A = 60^\circ$.

Solution:

Given: $mAB = 6\text{cm}$; $mAC = 4\text{cm}$;

$m\angle A = 60^\circ$.

Required: $m \overline{CB} = ?$



In $\triangle ABC$, we have


$$\begin{aligned}(\overline{BC})^2 &= (\overline{AB})^2 + (\overline{AC})^2 - 2(\overline{AB})(\overline{AD}) \\&= (6)^2 + (4)^2 - 2(6)(x) \\&= 36 + 16 - 2(6)(2) \\&= 52 - 24 \\&= 28 \\m\overline{BC} &= \sqrt{28} \\&= 2\sqrt{7} \text{ cm} \Rightarrow 5.29 \text{ cm}\end{aligned}$$

$$\begin{aligned}\because \cos 60^\circ &= \frac{x}{4} \\ \frac{1}{2} &= \frac{x}{4} \\ 2x &= 4 \\ \Rightarrow x &= 2\end{aligned}$$

Q2. In a $\triangle ABC$, $m \overline{AB} = 6\text{cm}$, $m \overline{BC} = 8\text{cm}$, $m \overline{AC} = 9\text{cm}$ and D is the midpoint of side $\overline{AC}$. Find length of the median $\overline{BD}$.

Solution:

According to the figure, we have



$$\begin{aligned}m\overline{AD} &= \overline{DC} \\ \text{and } m\overline{AC} &= m\overline{AD} + m\overline{DC} \\ m\overline{AC} &= m\overline{AD} + m\overline{AD} \\ 9 &= 2m\overline{AD} \\ \text{Or } 2m\overline{AD} &= 9\end{aligned}$$

$$m\overline{AD} = \frac{9}{2} = 4.5 \text{ cm}$$


We know that

$$\begin{aligned}(\overline{AC})^2 + (\overline{BC})^2 &= 2[(\overline{AD})^2 + (\overline{BD})^2] \\ (6)^2 + (8)^2 &= 2[(4.5)^2 + (\overline{BD})^2] \\ 36 + 64 &= 2(4.5)^2 + 2(\overline{BD})^2 \\ 100 &= 40.5 + 2(\overline{BD})^2 \\ \text{Or } 2(\overline{BD})^2 &= 100 - 40.5 \\ 2\overline{BD}^2 &= 59.5 \\ \Rightarrow \overline{BD}^2 &= 29.75 \\ \Rightarrow \overline{BD} &= \sqrt{29.75} = 5.45 \text{ cm}\end{aligned}$$

Q3. In a parallelogram $ABCD$ prove that $(\overline{AC})^2 + (\overline{BD})^2 = 2[(\overline{AB})^2 + (\overline{BC})^2]$

Solution:

$$\begin{aligned}(\overline{BD})^2 &= (\overline{CD})^2 + (\overline{BC})^2 + 2(\overline{BC})(\overline{CE}) \quad (1) \\ (\overline{AC})^2 &= (\overline{AB})^2 + (\overline{BC})^2 - 2(\overline{BC})(\overline{BF}) \quad (2)\end{aligned}$$



Adding (1) and (2), we get

$$\begin{aligned}(\overline{AC})^2 + (\overline{BD})^2 &= (\overline{CD})^2 + (\overline{BC})^2 + 2(\overline{BC})\overline{CE} + (\overline{AB})^2 + (\overline{BC})^2 - 2(\overline{BC})(\overline{BF}) \\ &= (\overline{AB})^2 + (\overline{CD})^2 + 2(\overline{BC})^2 + 2(\overline{BC})(\overline{CE}) - 2(\overline{BC})(\overline{BF})\end{aligned}$$

In parallelogram opposite sides are congruent, so

$AB = DC$, $AD = BC$, and $BF = CE$

$$\begin{aligned}(\overline{AC})^2 + (\overline{BD})^2 &= 2(\overline{AB})^2 + 2(\overline{BC})^2 + 2(\overline{CE}) - 2(\overline{BC})\overline{CE} \\ (\overline{AC})^2 + (\overline{BD})^2 &= 2(\overline{AB})^2 + 2(\overline{BC})^2\end{aligned}$$

$$(\overline{AC})^2 + (\overline{BD})^2 = 2[(\overline{AB})^2 + (\overline{BC})^2]$$

Hence Proved.

THEOREM 1

8.1 (I) In an obtuse angled triangle, the square on the side opposite to the obtuse angle is equal to the sum of the squares on the sides containing the obtuse angle together with twice the rectangle contained by one of the sides, and the projection on it of the other.

Given:


ABC is a triangle having an obtuse angle

BAC at A. Draw CD perpendicular on BA produced.

So that AD is the projection of AC on BA produced.

Take mBC = a, mCA = b, mAB = c,

mAD = x and mCD = h.



To prove

$$\left(\overline{BC} \right)^2 = \left(\overline{AC} \right)^2 + \left(\overline{AB} \right)^2 + 2 \left(m\overline{AB} \right) \left(m\overline{AD} \right)$$

i.e., $a^2 = b^2 + c^2 + 2cx$


Proof:

Statements	Reasons
In $\angle \text{rt} \square CDA$, $m\angle CDA = 90^\circ$	Given
$\therefore \left(\overline{AC} \right)^2 = \left(\overline{AD} \right)^2 + \left(\overline{CD} \right)^2$	Pythagoras Theorem
or $b^2 = x^2 + h^2$ ____ (i)	
In $\angle \text{rt} \square CDB$, $m\angle CDB = 90^\circ$	Given
$\therefore \left(\overline{BC} \right)^2 = \left(\overline{BD} \right)^2 + \left(\overline{CD} \right)^2$	Pythagoras Theorem
or $a^2 = \left(c + x \right)^2 + h^2$ $= c^2 + 2cx + x^2 + h^2$ ____ (ii)	$\overline{BD} = \overline{BA} + \overline{AD}$
Hence, $a^2 = c^2 + 2cx + b^2$ i.e., $a^2 = b^2 + c^2 + 2cx$	Using (i) and (ii)
or $\left(\overline{BC} \right)^2 = \left(\overline{AC} \right)^2 + \left(\overline{AB} \right)^2 + 2 \left(m\overline{AB} \right) \left(m\overline{AD} \right)$	

Example

In a $\square ABC$ with obtuse angle at A, if CD is an altitude on BA produced and in m AC = m AB .

Then prove that (BC)² = 2 (AB)(BD)



Given:

In a $\square ABC$, mZA is obtuse mAC = mAB and CD being altitude on BA produced.

To prove:

$$\left(\overline{BC} \right)^2 = 2 \left(m\overline{AB} \right) \left(m\overline{BD} \right)$$

Proof:

In a $\square ABC$, having obtuse angle BAC at A.

Statements	Reasons
$\left(\overline{BC} \right)^2 = \left(\overline{BA} \right)^2 + \left(\overline{AC} \right)^2 + 2 \left(m\overline{BA} \right) \left(m\overline{AD} \right)$	By Theorem I
$= \left(\overline{AB} \right)^2 + \left(\overline{AB} \right)^2 + 2 \left(m\overline{AB} \right) \left(m\overline{AD} \right)$	Given
$= 2 \left(\overline{AB} \right)^2 + 2 \left(m\overline{AB} \right) \left(m\overline{AD} \right)$	
$\left(\overline{BC} \right)^2 = 2m\overline{AB} \left(m\overline{AB} + m\overline{AD} \right)$	On the line segment BD
$= 2m\overline{AB} \left(m\overline{BD} \right)$	Point A is between B and D

THEOREM 2

8.1 (ii) In any triangle, the square on the side opposite to acute angle is equal to sum of the squares on the sides containing that acute angle diminished by twice the rectangle contained by one of those sides and the projection on it of the other.

Given:

ΔABC with an acute angle CAB at A


Take mBC = a, mCA = b and mAB = c

Draw CD ⊥ AB so that AD is projection of AC on AB

Also, mAD = x and mCD = h

To Prove:

$(\overline{BC})^2 = (\overline{AC})^2 + (\overline{AB})^2 - 2(\overline{mAB})(\overline{mAD})$ i.e., $a^2 = b^2 + c^2 - 2cx$



Proof:

Statements	Reasons
In ΔCDA, m∠CDA = 90° $(\overline{AC})^2 = (\overline{AD})^2 + (\overline{CD})^2$ or $b^2 = x^2 + h^2$ ____ (i)	Given Pythagoras Theorem
In ΔCDB, m∠CDB = 90° $(\overline{BC})^2 = (\overline{BD})^2 + (\overline{CD})^2$ $a^2 = (c - x)^2 + h^2$ or $a^2 = c^2 - 2cx + x^2 + h^2$ ____ (ii) $a^2 = c^2 - 2cx + b^2$	Given Pythagoras Theorem From the figure
Hence $a^2 = b^2 + c^2 - 2cx$ i.e., $(\overline{BC})^2 = (\overline{AC})^2 + (\overline{AB})^2 - 2(\overline{mAB})(\overline{mAD})$	Using (i) and (ii)

MISCELLANEOUS EXERCISE 8

Q1. In a $\triangle ABC$, $m\angle A = 60^\circ$, prove that $(\overline{BC})^2 = (\overline{AB})^2 + (\overline{AC})^2 - m\overline{AB} \cdot \overline{AC}$.

Solution:

Given:

In $\triangle ABC$, $m\angle A = 60^\circ$

Required:

$$(\overline{BC})^2 = (\overline{AB})^2 + (\overline{AC})^2 - \overline{AB} \cdot \overline{AC}$$

Construction:

Draw $\overline{CD} \perp \overline{AB}$, so that $\overline{AD}$ the Projection of $\overline{AC}$ on $\overline{AB}$.

Proof:

In right angle $\triangle ACD$

$$\cos 60^\circ = \frac{m\overline{AD}}{m\overline{AC}}$$


$$\frac{1}{2} = \frac{m\overline{AD}}{m\overline{AC}} \quad \left(\cos 60^\circ = \frac{1}{2} \right)$$

$$m\overline{AD} = \frac{1}{2} m\overline{AC}$$

Now, according to the theorem, we have

$$\overline{BC}^2 = \overline{AB}^2 + \overline{AC}^2 - 2\overline{AB} \cdot \overline{AD}$$

$$\Rightarrow \overline{BC}^2 = \overline{AB}^2 + \overline{AC}^2 - \overline{AB} \cdot \overline{AC} \quad \because [2\overline{AD} = \overline{AC}]$$



Q2. In a $\triangle ABC$, $m\angle A = 45^\circ$, prove that $(\overline{BC})^2 = (\overline{AB})^2 + (\overline{AC})^2 - \sqrt{2} m\overline{AB} \cdot \overline{AC}$.

Solution:

Given:

In $\triangle ABC$, $m\angle A = 45^\circ$.

Required:

$$(\overline{BC})^2 = (\overline{AB})^2 + (\overline{AC})^2 - \sqrt{2} \overline{AB} \cdot \overline{AC}$$

Construction:

Draw $CD \perp AB$, so that AD is the projection of AC on AB .

Proof:

In right angle $\triangle ACD$

$$\cos 45^\circ = \frac{m\overline{AD}}{m\overline{AC}}$$

$$\frac{\sqrt{2}}{2} = \frac{m\overline{AD}}{m\overline{AC}}$$


$$2 m\overline{AD} = \sqrt{2} m\overline{AC}$$

$\triangle ABC$ is acuted angled at A, so according to the theorem, we have

$$\overline{BC}^2 = \overline{AB}^2 + \overline{AC}^2 - 2\overline{AB} \cdot \overline{AD}$$

$$\Rightarrow \overline{BC}^2 = \overline{AB}^2 + \overline{AC}^2 - \sqrt{2} \overline{AB} \cdot \overline{AC} \quad \because [2\overline{AD} = \sqrt{2} \overline{AC}]$$

Hence proved.



Q3. In a $\triangle ABC$, calculate $m\overline{BC}$ when $m\overline{AB} = 5$ cm, $m\overline{AC} = 4$ cm, $m\angle A = 60^\circ$.

Solution:

We know that when $m\angle A = 60^\circ$ then,


$$\overline{BC}^2 = \overline{AB}^2 + \overline{AC}^2 - \overline{AB} \cdot \overline{AC}$$

$$= 5^2 + 4^2 - 5 \cdot 4$$

$$= 25 + 16 - 20$$

$$= 21$$

$$m\overline{BC} = \sqrt{21} = 4.58 \text{ cm}$$



Q4. In a $\triangle ABC$, calculate $m\overline{AC}$ when $m\overline{AB} = 5$ cm, $m\overline{BC} = 4\sqrt{2}$ cm, $m\angle B = 45^\circ$.

Solution:

We know that when $m\angle B = 45^\circ$ then,


$$\overline{AC}^2 = \overline{AB}^2 + \overline{BC}^2 - \sqrt{2} \overline{AB} \cdot \overline{BC}$$

$$= (5)^2 + (4\sqrt{2})^2 - \sqrt{2} (5) (4\sqrt{2})$$

$$= 25 + 32 - 40$$

$$= 57 - 40 = 17$$

$$m\overline{AC} = \sqrt{17} \text{ cm} = 4.123 \text{ cm}$$



Q5. In a triangle ABC, $m\overline{BC} = 21$ cm, $m\overline{AC} = 17$ cm, $m\overline{AB} = 10$ cm.

Measure the length of projection of $\overline{AC}$ upon $\overline{BC}$.

Solution:

$c = 10$ cm, $a = 21$ cm, $b = 17$ m, $x = ?$

We know that

$$c^2 = a^2 + b^2 - 2(a)(x)$$

$$(10)^2 = (21)^2 + (17)^2 - 2(21)(x)$$


$$100 = 441 + 189 - 42x$$

$$42x = 441 + 189 - 100$$

$$42x = 730 - 100$$

$$42x = 630$$

$$x = \frac{630}{42} = 15 \text{ cm}$$



Q6. In a triangle ABC, $m\overline{BC} = 21$ cm, $m\overline{AC} = 17$ cm, $m\overline{AB} = 10$ cm.

Calculate the projection of $\overline{AB}$ upon $\overline{BC}$.

Solution:

Q7. In a $\triangle ABC$, $a = 17$ cm, $b = 15$ cm and $c = 8$ cm find $m\angle A$.

Solution:

Given:

In $\triangle ABC$, $a = 17$ cm, $b = 15$ cm and, $c = 8$ cm

Required:

$m\angle A = ?$

By Pythagoras theorem.

$$a^2 = b^2 + c^2$$


$$(17)^2 = (15)^2 + (8)^2$$

$$289 = 225 + 64$$

$$289 = 289$$

So, it satisfied, that given values are the sides of a right angled triangle.

$$\therefore m\angle A = 90^\circ$$



Q8. In a $\triangle ABC$, $a = 17$ cm, $b = 15$ cm and $c = 8$ cm find $m\angle B$.

Solution:

Given:

In $\triangle ABC$; $a = 17$ cm, $b = 15$ cm and $c = 8$ cm


Required:

$m\angle B = ?$

We know that it is right angled triangle.

$$\sin(m\angle B) = \frac{b}{a} = \frac{15}{17} = 0.882$$

$$m\angle B = \sin^{-1}(0.882) = 61.90^\circ$$



Q9. Whether the triangle with sides 5 cm, 7 cm, 8 cm is acute, obtuse or right angled.

Solution:

Given:

$a = 5$ cm; $b = 7$ cm; $c = 8$ cm

Case I:

$$c^2 = a^2 + b^2$$

$$(8)^2 = (5)^2 + (7)^2$$

$$64 = 25 + 49$$

$$64 \neq 74$$

It is not right angled triangle.

Also

$$74 > 64 \text{ i.e.}$$

$$a^2 + b^2 > c^2$$

The result shows that the given triangle is an acute angled triangle.

Q10. Whether the triangle with sides 8 cm, 15 cm, 17 cm is acute, obtuse or right angled.

Solution:

Given:

$a = 8$; $b = 15$; $c = 17$

Case I:

Q11. Whether the triangle with sides 8 cm, 15 cm, 17 cm is acute, obtuse or right angled.

Solution:

Given:

$a = 8$; $b = 15$; $c = 17$

Case I:

$$c^2 = a^2 + b^2$$

$$(17)^2 = (8)^2 + (15)^2$$


$$289 = 64 + 225$$

$$289 = 289$$


Hence, It is right angled triangle.

Summary

✓ The projection of a given point on a line segment is the foot $\perp$ of drawn from the point on that line segment. If $\overline{CD} \perp \overline{AB}$, then evidently D is the foot of perpendicular $\overline{CD}$ from point C on the line segment AB.



✓ The projection of a line segment $\overline{CD}$ on a line segment AB is the portion $\overline{ED}$ of the latter intercepted between feet of the perpendiculars drawn from C and D. However, projection of a vertical line segment $\overline{CD}$ on a line segment AB is a point on $\overline{AB}$ which is of zero dimension.



✓ In an obtuse-angled triangle, the square on the side opposite to the obtuse angle is equal to the sum of the squares on the sides containing the obtuse angle together with twice the rectangle contained by one of the sides, and the projection on it of the other.

✓ In any triangle, the square on the side opposite to an acute angle is equal to the sum of the squares on the sides containing that acute angle

Q12. Whether the triangle with sides 8 cm, 15 cm, 17 cm is acute, obtuse or right angled.

Solution:

Given:

$a = 8$; $b = 15$; $c = 17$

Case I:

$$c^2 = a^2 + b^2$$

$$(17)^2 = (8)^2 + (15)^2$$

$$289 = 64 + 225$$

$$289 = 289$$

Hence, It is right angled triangle.

diminished by twice the rectangle contained by one of those sides and the projection on it of the other.

✓ In any triangle, the sum of the squares on any two sides is equal to twice the square on half the third side together with twice the square on the median which bisects the third side (Apollonius' Theorem).